


# NEVADA TEST SITE WETLANDS ASSESSMENT

**RECEIVED**

DEC 24 1997

**OSTI**



May 1997

Submitted to

U.S. Department of Energy  
Nevada Operations Office  
Environmental Protection Division  
P.O. Box 98518

Prepared by  
***Bechtel Nevada***

Ecological Services  
P.O. Box 98521  
Las Vegas, NV 89193-8521

## DISCLAIMER NOTICE

“This Report was prepared as an account of work sponsored by an agency of the U.S. Government. Neither the U.S. Government nor any agency thereof, nor any of their employees, makes any warranty or representation, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the U.S. Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the U.S. Government or any agency thereof.”

This report has been reproduced from the best available copy.

Available to DOE and DOE contractors from the Office of Scientific and Technical Information, Post Office Box 62, Oak Ridge, Tennessee 37831. (423) 576-8401:

Available to the public from the National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Road, Springfield, VA 22161. (703) 487-4650.

## **DISCLAIMER**

**Portions of this document may be illegible in electronic image products. Images are produced from the best available original document.**

# NEVADA TEST SITE WETLANDS ASSESSMENT

By

Dennis J. Hansen  
Paul D. Greger  
Cathy A. Wills  
W. Kent Ostler

May 1997

WORK PERFORMED UNDER  
CONTRACT NO. DE-AC08-96NV11718

Submitted to

U.S. Department of Energy  
Nevada Operations Office  
Environmental Protection Division  
P.O. Box 98518

## **ACKNOWLEDGMENTS**

The authors wish to thank the following individuals who made significant contributions to this report. Dan C. Steen and Derek B. Hall produced the GIS-generated figures in this report. Dodie M. Witham assisted with the literature review and data management.

# CONTENTS

EXECUTIVE SUMMARY .....	xi
1.0 INTRODUCTION .....	1
1.1 Definition of a Wetland .....	1
1.2 Ecological Importance of Wetlands .....	1
1.4 Need and Purpose for Wetlands Survey on the NTS .....	2
1.5 Goals and Objectives of the NTS Wetlands Survey and Report .....	3
2.0 STUDY AREA .....	5
3.0 METHODS .....	9
3.1 Literature Search to Identify Study Sites and Historical Use .....	9
3.2.1 Determining Wetland Site Coordinates .....	10
3.2.2 Selecting Wetland Delineation Methods .....	11
3.2.3 Delineating Wetland Boundaries .....	12
3.2.4 Characterizing Wetland Vegetation .....	14
3.2.5 Characterizing Wetland Hydrology .....	15
3.2.6 Characterizing Hydric Soils .....	16
3.2.7 Delineating Jurisdictional Wetlands .....	18
3.2.8 Describing Historical Use .....	19
3.2.9 Characterizing Wildlife Use .....	19
3.2.10 Photographing Wetlands .....	19
3.2.11 Creating a Wetlands Geospatial Database .....	20
4.0 RESULTS .....	21
4.1 Previous Studies .....	21
4.1.1 Hydrology .....	21
4.1.2 Water Quality .....	22
4.1.3 Vegetation .....	22
4.1.4 Wildlife Use .....	23
4.1.5 Historical Use .....	23
4.2 Description of Study Sites .....	24
4.2.1 Ammonia Tanks .....	24
4.2.1.1 Site Description and Historical Use .....	24
4.2.1.2 Hydrophytic Vegetation .....	24
4.2.1.3 Hydrology .....	24
4.2.1.4 Hydric Soils .....	27

## CONTENTS

4.2.1.5	Determination of Jurisdictional Status	27
4.2.1.6	Wildlife Use	27
4.2.2	Cane Spring	28
4.2.2.1	Site Description and Historical Use	28
4.2.2.2	Hydrophytic Vegetation	28
4.2.2.3	Wetland Hydrology and Water Quality	33
4.2.2.4	Hydric Soils	33
4.2.2.5	Determination of Jurisdictional Status	34
4.2.2.6	Wildlife Use	34
4.2.3	Captain Jack Spring	34
4.2.3.1	Site Description and Historical Use	34
4.2.3.2	Hydrophytic Vegetation	36
4.2.3.3	Wetland Hydrology and Water Quality	36
4.2.3.4	Hydric Soils	39
4.2.3.5	Determination of Jurisdictional Status	39
4.2.3.6	Wildlife Use	39
4.2.4	Cottonwood Spring	40
4.2.4.1	Site Description and Historical Use	40
4.2.4.2	Hydrophytic Vegetation	40
4.2.4.3	Wetland Hydrology and Water Quality	43
4.2.4.4	Hydric Soils	45
4.2.4.5	Determination of Jurisdictional Status	45
4.2.4.6	Wildlife Use	45
4.2.5	Coyote Spring	45
4.2.5.1	Site Description and Historical Use	45
4.2.5.2	Hydrophytic Vegetation	45
4.2.5.3	Wetland Hydrology and Water Quality	48
4.2.5.4	Hydric Soils	48
4.2.5.5	Determination of Jurisdictional Status	48
4.2.5.6	Wildlife Use	49
4.2.6	Fortymile Canyon Tanks	49
4.2.6.1	Site Description and Historical Use	49
4.2.6.2	Hydrophytic Vegetation	49
4.2.6.3	Hydrology	49
4.2.6.4	Hydric Soils	52

## CONTENTS

4.2.6.5	Jurisdictional Wetland Determination .....	52
4.2.6.6	Wildlife Use .....	52
4.2.7	Gold Meadows Spring .....	53
4.2.7.1	Site Description and Historical Use .....	53
4.2.7.2	Hydrophytic Vegetation .....	53
4.2.7.3	Wetland Hydrology and Water Quality .....	53
4.2.7.4	Hydric Soils .....	53
4.2.7.5	Determination of Jurisdictional Status .....	56
4.2.7.6	Wildlife Use .....	56
4.2.8	John's Spring .....	56
4.2.8.1	Site Description and Historical Use .....	56
4.2.8.2	Hydrophytic Vegetation .....	59
4.2.8.3	Wetland Hydrology and Water Quality .....	59
4.2.8.4	Hydric Soils .....	59
4.2.8.5	Determination of Jurisdictional Status .....	60
4.2.8.6	Wildlife Use .....	60
4.2.9	Oak Spring .....	60
4.2.9.1	Site Description and Historical Use .....	60
4.2.9.2	Hydrophytic Vegetation .....	61
4.2.9.3	Wetland Hydrology and Water Quality .....	64
4.2.9.4	Hydric Soils .....	64
4.2.9.5	Determination of Jurisdictional Status .....	64
4.2.9.6	Wildlife Use .....	65
4.2.10	Pavits Spring .....	65
4.2.10.1	Site Description and Historical Use .....	65
4.2.10.2	Hydrophytic Vegetation .....	65
4.2.10.3	Wetland Hydrology And Water Quality .....	65
4.2.10.4	Hydric Soils .....	65
4.2.10.5	Determination of Jurisdictional Status .....	69
4.2.10.6	Wildlife Use .....	69
4.2.11	Rainier Spring .....	69
4.2.11.1	Site Description and Historical Use .....	69
4.2.11.2	Hydrophytic Vegetation .....	72
4.2.11.3	Hydrology .....	72
4.2.11.4	Hydric Soils .....	73

## CONTENTS

4.2.11.5	Jurisdictional Wetland Determination	73
4.2.11.6	Wildlife Use	73
4.2.12	Reitmann Seep	73
4.2.12.1	Site Description and Historical Use	73
4.2.12.2	Hydrophytic Vegetation	73
4.2.12.3	Wetland Hydrology and Water Quality	76
4.2.12.4	Hydric Soils	76
4.2.12.5	Determination of Jurisdictional Status	77
4.2.12.6	Wildlife Use	77
4.2.13	Rock Valley Tank	77
4.2.13.1	Site Description and Historical Use	77
4.2.13.2	Hydrophytic Vegetation	77
4.2.13.3	Hydrology	77
4.2.13.4	Hydric Soils	80
4.2.13.5	Jurisdictional Wetland Determination	80
4.2.13.6	Wildlife Use	80
4.2.14	Tippipah Spring	80
4.2.14.1	Site Description and Historical Use	80
4.2.14.2	Hydrophytic Vegetation	83
4.2.14.3	Wetland Hydrology and Water Quality	83
4.2.14.4	Hydric Soils	88
4.2.14.5	Determination of Jurisdictional Status	88
4.2.14.6	Wildlife Use	89
4.2.15	Tongue Wash Tank	89
4.2.15.1	Site Description and Historical Use	89
4.2.15.2	Hydrophytic Vegetation	89
4.2.15.3	Hydrology	89
4.2.15.4	Hydric Soils	89
4.2.15.5	Jurisdictional Wetland Determination	89
4.2.15.6	Wildlife Use	92
4.2.16	Topopah Spring	92
4.2.16.1	Site Description and Historical Use	92
4.2.16.2	Hydrophytic Vegetation	92
4.2.16.3	Wetland Hydrology and Water Quality	98
4.2.16.4	Hydric Soils	98

## CONTENTS

4.2.16.5	Determination of Jurisdictional Status	98
4.2.16.6	Wildlife Use	98
4.2.17	Tub Spring	98
4.2.17.1	Site Description and Historical Use	98
4.2.17.2	Hydrophytic Vegetation	101
4.2.17.3	Wetland Hydrology and Water Quality	101
4.2.17.4	Hydric Soils	102
4.2.17.5	Determination of Jurisdictional Status	102
4.2.17.6	Wildlife Use	102
4.2.18	Tupapa Seep	102
4.2.18.1	Site Description and Historical Use	102
4.2.18.2	Hydrophytic Vegetation	102
4.2.18.3	Wetland Hydrology and Water Quality	102
4.2.18.4	Hydric Soils	105
4.2.18.5	Determination of Jurisdictional Status	105
4.2.18.6	Wildlife Use	105
4.2.19	Twin Spring	106
4.2.19.1	Site Description and Historical Use	106
4.2.19.2	Hydrophytic Vegetation	106
4.2.19.3	Wetland Hydrology and Water Quality	109
4.2.19.4	Hydric Soils	110
4.2.19.5	Determination of Jurisdictional Status	110
4.2.19.6	Wildlife Use	110
4.2.20	Wahmonie Seep 1	110
4.2.20.1	Site Description and Historical Use	110
4.2.20.2	Hydrophytic Vegetation	110
4.2.20.3	Wetland Hydrology and Water Quality	113
4.2.20.4	Hydric Soils	113
4.2.20.5	Determination of Jurisdictional Status	114
4.2.20.6	Wildlife Use	114
4.2.21	Wahmonie Seep 2	114
4.2.21.1	Site Description and Historical Use	114
4.2.21.2	Hydrophytic Vegetation	114
4.2.21.3	Wetland Hydrology and Water Quality	114
4.2.21.4	Hydric Soils	117

4.2.22.6	Wildlife Use .....	120
4.2.23	Whiterock Spring .....	120
4.2.23.1	Site Description and Historical Use .....	120
4.2.23.2	Hydrophytic Vegetation .....	122
4.2.23.3	Wetland Hydrology and Water Quality .....	122
4.2.23.4	Hydric Soils .....	126
4.2.23.5	Determination of Jurisdictional Status .....	126
4.2.23.6	Wildlife Use .....	126
4.2.24	Yellow Rock Springs .....	126
4.2.24.1	Site Description and Historical Use .....	126
4.2.24.2	Hydrophytic Vegetation .....	126
4.2.24.3	Wetland Hydrology and Water Quality .....	129
4.2.24.4	Hydric Soils .....	129
4.2.24.5	Determination of Jurisdictional Status .....	129
4.2.24.6	Wildlife Use .....	130
4.2.25	Yucca Playa Pond .....	130
4.2.25.1	Site Description and Historical Use .....	130
4.2.25.2	Hydrophytic Vegetation .....	130
4.2.25.3	Wetland Hydrology and Water Quality .....	133
4.2.25.4	Hydric Soils .....	133
4.2.25.5	Determination of Jurisdictional Status .....	134
4.2.25.6	Wildlife Use .....	134
4.2.26	Potential Man-Induced Wetlands .....	134
4.2.27	Waters of the United States .....	135

	Methods and location coordinates of FWS natural water sources monitored in 1996 and 1997 . . . . .	7
3-1	Steps for conducting an on-site routine wetland delineation survey . . . . .	11
4-1	Ammonia Tanks wetland vegetation as surveyed on January 7, 1997 . . . . .	27
4-2	Cane Spring wetland vegetation as surveyed on June 19, 1996 . . . . .	32
4-3	Cane Spring wetland vegetation as surveyed on September 9, 1996 . . . . .	32
4-4	Captain Jack Spring wetland vegetation as surveyed on June 19, 1996 . . . . .	39
4-5	Cottonwood Spring wetland vegetation as surveyed on December 12, 1996 . . . . .	43
4-6	Coyote Spring wetland vegetation as surveyed on September 4, 1996 . . . . .	48
4-7	Fortymile Canyon Tanks wetland vegetation as surveyed on February 12, 1997 . . . . .	52

5-1	Wetland hydrology data and jurisdictional wetland status of natural water sources surveyed June 1996 – February 1997 . . . . .	139
5-2	Water quality measurements of natural water sources on the NTS surveyed June 1996 – February 1997 . . . . .	142
5-3	List of plants recorded at wetland sites on the NTS . . . . .	145
5-4	Number of vertebrate species recorded at each wetland study site on the NTS . . . . .	148

### List of Figures

2-1	Location of NTS natural water sources monitored in 1996 and 1997 . . . . .	6
3-1	Form used to document field indicators for wetlands on the NTS . . . . .	13
3-2	Form used to collect data on water quality and wildlife observations . . . . .	17
4-1	Location of Ammonia Tanks . . . . .	25
4-2	Location and sketch of Cane Spring . . . . .	29
4-3	Location of Captain Jack Spring . . . . .	35
4-4	Location and sketch of Cottonwood Spring . . . . .	41
4-5	Location and sketch of Coyote Spring, Pavits Spring, and Tupapa Seep and sketch of Coyote Spring . . . . .	46

4-14	Pool in rocky wash at Cottonwood Spring on January 8, 1997 . . . . .	44
4-15	Habitat at Coyote Spring on September 4, 1996 . . . . .	47
4-16	Closeup of vegetation at Coyote Spring on September 4, 1996 . . . . .	47

4-29	Habitat at Rainier Spring looking north on February 20, 1997	71
4-30	Closeup of vegetation at Rainier Spring on February 20, 1997	71
4-31	Habitat of Reitmann Seep looking southwest on June 2, 1989	75
4-32	Pool at Reitmann Seep on June 2, 1989	75
4-33	Habitat in the vicinity of Rock Valley Tank looking northwest on January 7, 1997	79
4-34	Tank opening (center) at Rock Valley Tank looking west on January 7, 1997	79
4-37	Tippipah Spring channel nearly dry, with drought-stressed vegetation, looking north on November 29, 1990	84
4-38	Upper reach of Tippipah Spring channel showing recovery of vegetation following drought, looking north on June 4, 1992	84
4-39	Middle reach of Tippipah Spring channel looking southwest on June 18, 1996	85
4-40	Lower reach of Tippipah Spring channel looking northwest on June 18, 1996	85
4-41	Cave opening at Tongue Wash Tank looking south on January 10, 1997	91
4-42	Tank inside cave at Tongue Wash Tank on January 10, 1997	91
4-43	Topopah Spring wetland area looking north on June 27, 1988	94
4-44	Vegetation of Topopah Spring cave pool on April 18, 1989	94
4-45	Outflow pipe at Topopah Spring on September 9, 1996	95
4-46	Hillside meadow at Topopah Spring looking north on June 20, 1996	95
4-47	Habitat at Tub Spring looking northwest on November 10, 1988	100
4-48	Vegetation around Tub Spring on August 24, 1996	100
4-49	Habitat around Tupapa Seep looking north on September 4, 1996	104
4-50	Closeup of vegetation at Tupapa Seep looking east on November 4, 1996	104
4-51	Habitat around Twin Spring looking east on January 8, 1997	108
4-52	Wetland vegetation at Twin Spring looking north on January 8, 1997	108
4-53	Wahmonie Seep 1 looking north on June 20, 1996	112
4-54	Wahmonie Seep 2 looking south on June 20, 1996	112

## CONTENTS

### List of Photographs

4-55	Wetland vegetation at Wahmonie Seep 2 looking south on June 20, 1996 . . . .	116
4-56	Wahmonie Seep 3 looking north on June 20, 1996 . . . . .	116
4-57	Whiterock Spring with grazed vegetation looking north about 1968 . . . . .	123
4-58	Whiterock Spring without grazing looking north on June 18, 1996 . . . . .	123
4-59	Emergent vegetation in the east cave of Whiterock Spring on December 7, 1990 . . . . .	124
4-60	Surface pool at Whiterock Spring looking north on December 7, 1990 . . . .	124
4-61	Habitat around Yellow Rock Spring looking southeast on January 29, 1997 . .	128
4-62	Drainage channel at Yellow Rock Spring looking southeast on January 29, 1997 . . . . .	128
4-63	Emergent vegetation at Yucca Playa Pond looking north on January 7, 1997 . . . . .	132
4-64	Drainage channel entering Yucca Playa Pond looking northwest on January 7, 1997 . . . . .	132

~~UTM~~  
UTM

~~Universal Transverse Mercator~~  
Universal Transverse Mercator

wildlife usage, and photographs. At selected sites, cursory physical and chemical water quality data were also collected.

Sixteen of the 25 NTS water sources surveyed met the three required criteria to be considered jurisdictional wetlands. They included Cane Spring, Captain Jack Spring, Cottonwood Spring, Coyote Spring, Gold Meadows Spring, John's Spring, Oak Spring, Reitmann Seep, Tippipah Spring, Topopah Spring, Twin Spring, Wahmonie Seep 1, Wahmonie Seep 2, Wahmonie Seep 3, Whiterock Spring, and the vegetated margins of Yucca Playa Pond.

wetland hydrology were Rainier Spring and Tupapa Seep. The source of water at most of the study sites is groundwater discharge from seeps and springs. Surface runoff from precipitation was found at Yucca Playa Pond and at the four tank sites.

Nineteen of the 25 study sites had field indicators for hydric soils which included saturated or inundated soils, dark-colored soils due to high organic matter content, and soil mottling. Field indicators were scarce at the 19 sites and were often inferred from site hydrology and past observations of surface water or saturated soils. Four of the six sites lacked saturated soils or other field indicators for hydric soils (Pavits Spring, Rainier Spring, Rock Valley Tank, and Tupapa Seep). Two of the six sites lacked soils altogether because they were located in bedrock (Fortymile Canyon Tanks and Tongue Wash Tank).

Eleven of the springs and seeps surveyed have surface flow of water all year long. These sites are Cane Spring, Captain Jack Spring, Cottonwood Spring, John's Spring, Oak Spring, Reitmann Seep, Tippipah Spring, Topopah Spring, Tub Spring, Twin Spring, and Whiterock Spring. The remaining 15 sites are ephemeral; they may dry up at some period of time during the year or during dry years. The sizes of the NTS wetlands are very small. With the exception of Tippipah Spring, Whiterock Spring, and Yucca Playa Pond, most were less than 300 square meters ( $m^2$ ) (3,228 square feet [ $ft^2$ ]). They varied in size from less than 1  $m^2$  (10.8  $ft^2$ ) at Reitmann Seep to approximately 3,400  $m^2$  (37,000  $ft^2$ ) along the edges of Yucca Playa Pond, based on the area of hydrophytic vegetation. Water levels were generally shallow at all study sites, ranging from 3 to 200 centimeters (cm) (1.2 to 78.7 inches [in.]). Flow rates of NTS springs measured during 1996 were very low, ranging from 0.0-3.0 liters/minute ( $l/min$ ) (0.0 to 0.80 gallons [gal]/min).

Eighty-one species of vascular plants have been recorded in or near the NTS wetland sites based on the results of the 1996 and 1997 field surveys and on past NTS studies. Most of the species in NTS wetlands are forbs (33 species, 41 percent) followed by grasses, rushes, and sedges combined (30 species, 37 percent), and trees and shrubs combined (18 species, 22 percent). This is very different from the general flora of the NTS, where forbs make up approximately 74 percent of the total number of species. Grasses, rushes, and sedges

management goals specific for NTS wetlands that incorporate the intent of existing wetlands legislation, the principles of ecosystem management, and the interests of regional land managers and other stakeholders. Specific management goals presented in this report include (1) avoiding wetland impacts whenever possible, (2) minimizing all unavoidable wetland impacts, (3) restoring the biological integrity of wetlands if degradation occurs, and (4) preserving and enhancing the natural and beneficial values of NTS wetlands. Other recommendations pertaining to the management of NTS natural water sources are discussed in the final chapter of this report.

**This Page Intentionally Left Blank**

Identifying and characterizing wetlands is important because of their value to local and regional ecosystems. Wetlands are known to have three major functions: (1) habitat maintenance and food web support; (2) short- and long-term storage of water; and (3) cycling of nutrients, removal of dissolved substances, and accumulation of inorganic sediments. The ecological and societal value of these functions include flood damage control, maintaining water quality, maintaining biodiversity, and providing habitat and forage for nongame and game species such as fish, fur-bearers, and waterfowl.

Wetlands within desert ecosystems are distinctly different from the familiar examples of large marshlands, peatlands, and river floodplains within other physiographic regions of the nation. Desert springs and seeps are often too small in size to affect local or regional surface water flow or nutrient cycling. However, they do provide wildlife habitat, free-standing water, and forage which are regionally rare. They may increase the biodiversity of desert ecosystems, provide habitat and forage to migratory species passing through desert ecosystems, or help to determine the home range of resident wildlife species and the size of resident wildlife populations dependent on drinking water.

### **1.3 Legal Status of Wetlands**

Wetlands are federally regulated throughout the nation on both private and public lands. The Clean Water Act (CWA) of 1977 (42 U.S. Code [USC] 1251, *et seq.* [amendments to the Federal Water Pollution Control Act of 1972]) was enacted to maintain and restore the chemical, physical, and biological integrity of the "waters of the United States." Section 404 of the Act authorizes the U.S. Army Corps of Engineers (USACE) to issue permits

available literature. These lesser-known natural water sources are equally rare and unique NTS habitats important to regional wildlife and localized populations of aquatic organisms and water-dependent vegetation. Their identification is essential for proper management and protection of natural resources on the NTS.

DOE/Nevada Operations Office (NV) has for the past two decades implemented an ecological monitoring program on the NTS to monitor various components of the ecosystem and to provide baseline data needed to identify and protect rare biological resources and federal- and state-protected plants and animals. The monitoring of wildlife

but they were not in identifying and delineating the natural water sources on the NTS, was conducted in 1996 and 1997 through DOE/NV's Ecological Monitoring and Compliance (EMAC) program.

The pertinent ecological data gathered on wetlands during this survey will be incorporated into an NTS resource management plan. DOE's *Land- and Facility-Use Management Policy* (O'Leary, 1994) is ". . . to manage all of its land and facilities as valuable national resources . . . based on the principles of ecosystem management and sustainable development. [DOE] will integrate mission, economic, ecologic, social, and cultural factors in a comprehensive plan for each site that will guide land and facility decisions . . . . This policy will result in land and facility uses that support the Department's critical missions, stimulate the economy, and protect the environment." The principles of ecosystem management is an approach to sustain the production of natural resources and the ecosystems on which those resources depend. This resource management plan will identify the rare and unique habitats of the NTS, such as wetlands, and how they will be managed based on ecosystem principles.

This wetlands survey is also helpful for compliance with NEPA. In the *Final Environmental Impact Statement for the Nevada Test Site and Off-Site Locations in the State of Nevada* (EIS) (DOE, 1996a), several alternative DOE activities proposed for the NTS over the next ten years were analyzed for their impacts on the existing environment, which included ten known springs and seeps. None of the proposed activities were expected to negatively affect these water sources. Information regarding all the natural water sources, however, even those that are more remote or are ephemeral, may be needed during NEPA impact analyses and siting suitability analyses for new NTS projects developed during, and well beyond, the next ten years.

Due to their federal regulatory status, this survey was designed to provide a preliminary delineation of jurisdictional wetlands that occur on the NTS. If a proposed project would impact a jurisdictional wetland, a permit from the USACE would need to be obtained before construction could begin. This survey also identifies those natural water sources which do not qualify as jurisdictional wetlands, but which may be protected as waters of the United States. Examples of waters of the United States are springs, seeps, tanks, and playas that do not support hydrophytic vegetation, but which have standing water for long periods. The permitting process for projects which may impact these waters is similar to that for jurisdictional wetlands. This survey of natural NTS water sources may expedite project siting, permitting, and construction by identifying up front those sites that may be regulated under Section 404 of the CWA.

## **1.5 Goals and Objectives of the NTS Wetlands Survey and Report**

The goals and objectives of the 1996-1997 NTS wetlands field survey and this summary report are to

- identify and summarize all previous studies of NTS natural water sources;
- describe the physical, chemical, and biological features of these water sources;

~~JURISDICTIONAL WETLANDS OF WATERS OF THE UNITED STATES, AND~~

- identify current DOE management practices related to the protection of NTS wetlands

directly related to the location of natural and man-made ponds (Greger, 1995).

A more detailed description of the physiography, geology, climate, and vegetation of the NTS has been presented in Beatley (1976) and O'Farrell and Emery (1976).




Figure 2-1 Location of NTS natural water sources monitored in 1996 and 1997

Table 2-1 Attributes and location coordinates of NTS natural water sources monitored in 1996 and 1997

Water Source	NTS Area	Elevation (m)	Habitat type <sup>a</sup>	Geology/Age <sup>b</sup>	UTM Easting	UTM Northing
Ammonia Tanks	19	1615	Sagebrush	Ash flow tuff/Miocene	562850	4110240
Cane Spring	5	1237	Blackbrush	Lava flows/Miocene	580750	4072641
Captain Jack Spring	12	1792	Pinyon-Juniper	Ash fall tuff/Miocene-Oligocene	573834	4113579
Cottonwood Spring	25	1292	Rabbitbrush	Rhyolitic lava flows-bedded tuff/Miocene	554045	4083726
Coyote Spring	27	1085	Creosote	Lava flows/Miocene	583594	4066568
Fortymile Canyon Tanks	25	1396	Rabbitbrush	Rhyolitic lavas-tuffaceous beds/Miocene	557500	4085000
Gold Meadows Spring	12	2048	Pinyon-Juniper	Ash fall tuff/Miocene-Oligocene	570415	4120398
John's Spring	15	1840	Sagebrush-Oak	Ash fall tuff/Miocene-Oligocene	582100	4122490
Oak Spring	15	1783	Sagebrush-Oak	Ash fall tuff/Miocene-Oligocene	582208	4122209
Pavits Spring	27	1203	Creosote	Lava flows/Miocene	581931	4068118
Rainier Spring	12	1890	Pinyon-Juniper	Ash fall tuff/Miocene-Oligocene	571463	4116050
Reitmann Seep	7	1402	Blackbrush	Ash fall tuff/Miocene	591278	4105578
Rock Valley Tank	25	1048	Creosote	Limestone, dolomites/Upper-Mid Cambrian	568070	4061000
Tippipah Spring	16	1585	Sagebrush	Ash fall tuff/Miocene-Oligocene	570857	4099671
Tongue Wash Tank	12	1950	Pinyon-Juniper	Ash fall tuff/Miocene	571360	4113050
Topopah Spring	29	1774	Blackbrush	Ash flow tuff/Miocene	564973	4088339
Tub Spring	15	1594	Sagebrush	Ash fall tuff/Miocene-Oligocene	584925	4121850
Tupapa Seep	27	1140	Creosote	Lava flows/Miocene	582129	4066459
Twin Spring	29	1310	Rabbitbrush	Rhyolitic lavas/Miocene	555484	4089984
Wahmonie Seep 1	26	1286	Blackbrush	Lava flows/Miocene	577679	4073923
Wahmonie Seep 2	26	1347	Blackbrush	Lava flows/Miocene	577471	4073319
Wahmonie Seep 3	26	1341	Blackbrush	Lava flows/Miocene	577044	4073349

Table 2-1 (continued)

Water Source	NTS Area	Elevation (m)	Habitat type <sup>a</sup>	Geology/Age <sup>b</sup>	UTM Easting	UTM Northing
Whiterock Spring	12	1539	Blackbrush	Ash fall tuff/ Miocene-Oligocene	577099	4117282
Yellow Rock Springs	30	1298	Blackbrush	Rhyolitic lava flow/Miocene	555979	4091944
Yucca Playa Pond	6	1189	Salt cedar	Alluvium/Holocene-Pliocene	584805	4090584

<sup>a</sup> Dominant perennial vegetation surrounding each site.

<sup>b</sup> Taken from Frizzell and Shulters (1990).

quality, wildlife use, species inventories, and documented anthropogenic disturbances. These data were summarized and presented in this report, along with the results of the field surveys conducted in 1996-1997.

Spring, Fortymile Canyon Tanks, John's Spring, Rainier Spring, Twin Spring, and Yellow Rock Springs. In January 1997, Ammonia Tanks, Rock Valley Tank, Yellow Rock Springs, and Yucca Playa Pond were visited, and in February 1997, Fortymile Canyon Tanks, John's Spring, and Rainier Spring were visited.

Data were collected at each site to (1) identify the location and size of each wetland (i.e., delineate wetland boundaries); (2) describe the vegetation, hydrology, and soils to delineate jurisdictional wetland boundaries; (3) collect cursory physical and chemical water quality data; (4) document wildlife usage; and (5) document wetland habitat with photographs.

### **3.2.1 Determining Wetland Site Coordinates**


The coordinates identifying the field location of each study site (Table 2-1) were recorded using hand-held Global Positioning System (GPS) units (Magellan™ ProMark V). Coordinates were recorded as Universal Transverse Mercator (UTM) system values (Zone 11, North American datum) based on an average of 20 GPS readings and an average PDOP for all 20 readings of 10 or less. PDOP is defined as the precision dilution of perception and refers to the three-dimensional spacing of satellites. If satellites are properly spaced (indicated by a low PDOP, where PDOPs of six or less are recommended), the resulting coordinate readings are more accurate. The GPS coordinates for each site were not corrected for degradation by the U.S. Department of Defense and are believed to be accurate to within 100 m (328 ft) of the true location. The UTM coordinates of each NTS wetland site were then cross-checked with coordinates from the appropriate USGS 7.5-Minute Series quadrangle maps.

Table S-1 Steps for conducting an on-site routine wetland delineation survey

- Determine whether an atypical situation exists
- Identify the plant community types
- Determine whether normal environmental conditions are present
- Select representative observation points
- Characterize each plant community type
- Record indicator status of dominant species
- Determine whether hydrophytic vegetation occurs
- Apply wetland hydrologic indicators
- Determine whether wetland hydrology is present
- Determine whether soils must be characterized
- Dig a soil pit
- Apply hydric soil indicators \*
- Determine whether hydric soils are present
- Make wetlands determination
- Determine wetland-nonwetland boundary
- Sample other transects and synthesize data

---

\* Munsell color chart classifications were not applied in describing soils; only the presence or absence of mottling was recorded. Table is cited from the methods described in *U.S. Army Corps of Engineers Wetlands Delineation Manual* (Environmental Laboratory, 1987).



# Routine Jurisdictional Wetland Determination

Name: \_\_\_\_\_ Wetland Unit: \_\_\_\_\_  
 Location: \_\_\_\_\_ UTM Coordinates Easting: \_\_\_\_\_ Northing: \_\_\_\_\_  
 Date: \_\_\_\_\_

## Hydrology

Type: Seep \_\_\_\_\_ Spring \_\_\_\_\_ Pond \_\_\_\_\_ Detention basin \_\_\_\_\_ Stream \_\_\_\_\_ Mechanically contained \_\_\_\_\_  
 Source: Natural \_\_\_\_\_ Man-enhanced \_\_\_\_\_ Man-made \_\_\_\_\_ Ephemeral \_\_\_\_\_ Permanent \_\_\_\_\_ Temporary \_\_\_\_\_  
 Date of construction/Period of flow: \_\_\_\_\_  
 Disturbance type (if any) and date: \_\_\_\_\_  
 Inundated: Yes \_\_\_\_\_ No \_\_\_\_\_ Depth of standing water \_\_\_\_\_ Saturated: Yes \_\_\_\_\_ No \_\_\_\_\_ Depth to saturation \_\_\_\_\_  
 Other field indicators: \_\_\_\_\_  
 Atypical situation: Yes \_\_\_\_\_ No \_\_\_\_\_ Wetland hydrology: Yes \_\_\_\_\_ No \_\_\_\_\_  
 Basis: \_\_\_\_\_

## Vegetation

List 3 dominant species, percent cover in bold, in each vegetation layer (5 if only 1 or 2 layers are present)

	Species	Indicator Status	% Cover
<b>Trees</b>			
1.	_____	_____	_____
2.	_____	_____	_____
3.	_____	_____	_____
4.	_____	_____	_____
5.	_____	_____	_____
<b>Shrubs</b>			
1.	_____	_____	_____
2.	_____	_____	_____
3.	_____	_____	_____
4.	_____	_____	_____
5.	_____	_____	_____
<b>Herbs</b>			
1.	_____	_____	_____
2.	_____	_____	_____
3.	_____	_____	_____
4.	_____	_____	_____
5.	_____	_____	_____
6.	_____	_____	_____
7.	_____	_____	_____
8.	_____	_____	_____
9.	_____	_____	_____
10.	_____	_____	_____

Other field indicators: \_\_\_\_\_  
 Percentage of species that are OBL, FACW, and/or FAC: \_\_\_\_\_%; Hydrophytic vegetation: Yes \_\_\_\_\_ No \_\_\_\_\_  
 Basis: \_\_\_\_\_

## Hydric Soils

Field indicators: \_\_\_\_\_  
 Hydric Soils: Yes \_\_\_\_\_ No \_\_\_\_\_  
 Jurisdictional Wetland Determination : Wetland \_\_\_\_\_ Nonwetland \_\_\_\_\_

## NOTES:

Figure 3-1 Form used to document field indicators for wetlands on the NTS

wetland (non-upland) areas was required. The wetland area at each site was subdivided into zones. Areas with standing water, or evidence of historic standing water (e.g., presence of dried algae or waterlines), were distinguished as one zone. Transitional areas between the very wet zone and the dry uplands were divided into one or more zones based on changes in species composition. The NTS wetlands were very small, often less than 0.4 ha (1 ac), and many only a few square meters in size. Therefore, usually only one wetland vegetation zone was identified per site. Within each major zone, a sample plot (also referred to as an "observation point") of approximately 1.5 m (5 ft) radius was selected. Plot size was smaller for small wetlands having less area. Tippipah Spring and Whiterock Spring were large wetlands where a mosaic of mesic habitats occurred or where a long linear wetland habitat occurred which had different vegetation characteristics downslope from the springhead. Within such larger wetland mosaics, boundaries of wetland zones that were patchy, discontinuous, or represented a changing linear corridor were sketched on a site map showing the relative position of site features. Approximate dimensions of each distinct wetland zone were recorded. The exact location of boundaries was difficult to document in the field because of a lack of aerial photos or maps with sufficient resolution or detail to permit determining their precise spacial locations.

### 3.2.4 Characterizing Wetland Vegetation

At each NTS site, one representative observation point per wetland zone was selected to examine field indicators for hydrophytic vegetation. The presence of hydrophytic vegetation is one of three parameters used to define a jurisdictional wetland. Vascular plants were identified and, using an ocular estimate (a visual projection of the plant canopy to the ground surface), were assigned an absolute percent cover estimate in each vegetation layer present (i.e., tree, shrub, herb). Plants in each vegetation layer with a 10 percent or greater cover value were defined as dominant species. All plants were classified as to their wetland status. To classify them, the *National List of Plant Species That Occur in Wetlands Intermountain (Region 8)* (Reed, 1996) was used. The plant names used are according to the National Plant Database (NRCS, 1996a). This list has assigned species to one of the following classes. A suffix of "-" or "+" is often used with the following codes to indicate that the plant is found on either the lower or higher ends of the range of probabilities that define each class. An asterisk (\*) following a regional indicator identifies tentative assignments based on limited information from which to determine the indicator status.

- UPL – Upland plants, occur almost exclusively in upland environments
- FACU – Facultative upland plants, occur in wetlands less than 33 percent of the time

- FAC – Facultative wetland species, occur in wetlands between 33 to 67 percent of the time
- FACW – Facultative wetland plants, occur in wetlands between 67 to 99 percent of the time
- OBL – Obligatory wetland species, occur in wetlands greater than 99 percent of the time
- NI – No indicator, or not sufficient information to classify this species at this time
- NL – Not listed in the *National List of Plant Species That Occur In Wetlands: Intermountain (Region 8)* because they occur in wetlands less than 1 percent of the time

One additional classification was assigned to some plants found at NTS study sites:


- UNKN – Unknown status because plants lacked taxonomic characteristics needed to determine the genus and species.

Those wetland zones in which greater than 50 percent of the dominant plants are classified as FAC, FACW, or OBL were considered to possess hydrophytic vegetation per the USACE guidelines (Environmental Laboratory, 1987). In situations where there were equal numbers of dominant hydrophytic and upland species (e.g., four dominant hydrophytic and four dominant upland species), an alternative method for establishing dominance, recommended in a USACE memorandum (Williams, 1992), was used. This method was presented in a 1989 interagency manual produced by the USACE, FWS, and the U.S. Department of Agriculture (USDA) and uses a comparison of the proportion of canopy cover due to hydrophytic versus upland species. If the canopy cover at a site due to hydrophytic plants is greater than 50 percent, then the site would be considered as having field indicators for hydrophytic vegetation. The presence of filamentous algae and moss, although they are not vascular plants used to define wetlands, also were recorded when observed at study sites.

To conform to USACE convention, the Latin scientific names of plants are used throughout the text of this report. The common name of each plant species is presented once in the text the first time it is discussed, and thereafter the genus and species name of each plant is used. To assist the reader, Appendix A presents a table of both the scientific and common names of all plants presented in the text or tables of this report.

### 3.2.5 Characterizing Wetland Hydrology

Wetland hydrology encompasses all hydrologic characteristics of areas that are periodically inundated or have soils saturated to the surface for some duration during the growing season, usually about 12.5 percent of the growing season (Environmental Laboratory, 1987). At the NTS, this is about 14 to 21 days depending on elevation. The presence of natural surface water or saturated soils (i.e., not derived from human intervention or construction of such things as wells or sumps) in the late summer or fall was considered sufficient evidence of wetland hydrology at each NTS study site. Because 1996 was a rather dry year with little precipitation, most of the water at the springs and seeps was assumed to be from subsurface flows rather than from runoff. Data from past visits to these sites were also used as evidence of perennial supplies of spring water.



### NTS Wetlands Water Quality and Wildlife Monitoring Form

Location: \_\_\_\_\_ Date: \_\_\_\_\_ Time: Start \_\_\_\_\_ Finish \_\_\_\_\_

Observer(s) \_\_\_\_\_ Air Temp °C: \_\_\_\_\_ Wind: \_\_\_\_\_ Weather: \_\_\_\_\_ Cloud Cover: \_\_\_\_\_

Location notes: \_\_\_\_\_

Water Temp (°C)	Dissolved Oxygen (ppm)	pH	TDS (ppm)	Cond/(µS)
1) _____	1) _____	1) _____	1) _____	1) _____
2) _____	2) _____	2) _____	2) _____	2) _____
3) _____	3) _____	3) _____	3) _____	3) _____

Max Depth of surface water: \_\_\_\_\_ Surface area of spring: \_\_\_\_\_ Spring flow rate: \_\_\_\_\_ l/min

Rel. Depth to water table: \_\_\_\_\_ Location notes: \_\_\_\_\_

Water Temp (°C)	Dissolved Oxygen (ppm)	pH	TDS (ppm)	Cond/(µS)
1) _____	1) _____	1) _____	1) _____	1) _____
2) _____	2) _____	2) _____	2) _____	2) _____
3) _____	3) _____	3) _____	3) _____	3) _____

Max depth of cave water \_\_\_\_\_ Surface area of cave pool: \_\_\_\_\_ Vegetation \_\_\_\_\_ Invertebrates \_\_\_\_\_

Filamentous algae/moss: \_\_\_\_\_

Genus species	Common Name	Total	Number Observed				Animal Sign		Condition/behavior/animal
			M	F	J	A	Track	Scat	
1									
2									
3									
4									
5									
6									

Searched riparian zone for presence of animal sign: \_\_\_\_\_. Abbreviations: M = male, F = female, J = juvenile, A = adult. Animal Sign: 1 = low abundance, 2 = moderate abundance, 3 = high abundance. Cloud cover = % visual estimates; 0-25% = low, 25-75% = moderate, 75-100% = high. Filamentous Algae: absent, minimal, abundant, or heavy growth.

Notes, UTMs, Slope, Drawings:

delineate sites as jurisdictional wetlands. It is known that the USACE regards desert springs and seeps to be sites where "atypical situations" often occur in regards to the presence of hydric soil indicators (personal communication with Nancy Kang, USACE, Reno, Nevada Office, November 21, 1996). An atypical situation, as defined by the USACE, is an occasion where one or more field indicators for wetlands (i.e., hydrophytic vegetation, hydrology, hydric soils) have been sufficiently altered by natural events or by recent human activities to preclude their presence (*Section F., Atypical Situations*, in Environmental Laboratory, 1987). NTS wetlands that support hydrophytic vegetation and have surface hydrology are considered by the USACE to possess hydric soils, even though the field indicators for hydric soils are absent. Such sites would be classified by the USACE as atypical situations and would still be classified as jurisdictional wetlands.

Soils in sites that are ponded or saturated for a long duration (seven days to one month) or a very long duration (>one month) during the growing season (NRCS, 1996b) are also defined as hydric soils according to the 1987 Manual. The growing season is defined in the 1987 Manual as the portion of the year when soil temperatures at 50 cm (19.7 in) below the soil surface are higher than biologic zero (5 degrees Celsius [ $^{\circ}\text{C}$ ]·[41 $^{\circ}$  Fahrenheit [F]) (for ease of determination, this period is usually approximated by the number of frost-free days). Because there had been only trace amounts of precipitation at the NTS prior to and during the fall of 1996, the presence of saturated soils or the presence of surface water, together with the presence of filamentous algae (which requires several weeks to grow), was interpreted as evidence to support the conclusion that the water must have persisted at least seven days and perhaps for several weeks or months. This water would have been present during a frost-free period (i.e., the growing season) and would therefore meet the criteria used to verify the presence of hydric soils.

### **3.2.7 Delineating Jurisdictional Wetlands**

After field data about field indicators were collected, a determination was made as to which area(s) within each site would be considered jurisdictional wetlands regulated by the USACE. Wetland plant community zones that were dominated by hydrophytic vegetation, hydric soils, and that had wetland hydrology were considered jurisdictional wetlands. Because of the small size of the jurisdictional wetlands on the NTS, it was not always possible to accurately depict boundaries on site sketches or aerial photographs (i.e., identifying 1 m [3.3 ft] boundaries on photos or maps that are only accurate to 10 m [32.8 ft]) or through narrative descriptions. The original intent of the wetland surveys was merely to identify NTS wetlands that had jurisdictional wetland areas that would require future field work to precisely identify boundaries. It is anticipated that if future development plans require disturbance of wetlands with areas considered jurisdictional, then the exact boundaries of the jurisdictional and nonjurisdictional areas would be surveyed and appropriate scale maps produced.

All boundaries of jurisdictional wetlands within this report should be considered approximate and managers directing future development activities within or near these areas should recognize that additional field work will be required to accurately identify jurisdictional boundaries. The size of these areas, and associated boundaries, probably

fluctuate over time along with the amount of groundwater surfacing at each site. Because detailed aerial photos or surveys were not available, the relative location of areas (plant communities) that are considered jurisdictional wetlands is communicated through site sketches or narrative descriptions. The USACE usually conducts a site visit to verify field boundaries and site conditions, and locating jurisdictional boundaries may be adjusted by the USACE at the time of the site visit.

### **3.2.8 Describing Historical Use**

Sites with prior historical use or natural events that obscure positive wetland field indicators require additional field techniques for delineating wetland boundaries. These altered sites are referred to as "atypical situations." Unique delineation procedures for these atypical situations are described in the 1987 Manual. These procedures attempt to determine site conditions prior to alteration by observing field evidence remaining at the site. Historical use at each potential wetland site was reviewed in the field and in the literature to determine if human alteration of a site occurred. Such alterations might include the removal of soils while attempting to improve water sources or while constructing stream channels. Each site was also evaluated to determine if natural events such as floods, fires, or landslides may have altered site conditions. Evidence of atypical situations, when found, is presented in the appropriate subsections of Section 4.2, "Site Description and Historical Use."

### **3.2.9 Characterizing Wildlife Use**

Biologists recorded all sightings of animals, presence of tracks, and scat observed at each wetland site. On approach to a site, biologists first observed and counted any fleeing animals. Biologists then walked the complete perimeter of the vegetated wetland zones to locate any tracks and scat. Tracks and scat were identified with the aid of Murie (1974). The spring pools were also inspected for the presence of selected aquatic invertebrate groups (e.g., snails, ostracods, copepods). After vegetation, hydrology, soils, wildlife signs, and aquatic animal data were collected, a stationary observation point was chosen where birds, attracted to the wetland site, were observed and counted for a period of 30 minutes. All animal use data from these 1996 and 1997 surveys were recorded (Figure 3-2), compiled, and added to an existing historical database of similar data collected at NTS springs and seeps from 1988 to 1994.

The common names of animals are used throughout this report. The Latin genus and species name for each animal is presented only once in the text the first time the animal is mentioned. Appendix D presents a table which includes both the common and scientific names of all animals mentioned in this report.

### **3.2.10 Photographing Wetlands**

An historical database of over 550 photographs taken on and near the NTS wetland sites between 1960 and 1996 was searched. Recent photographs which best showed the boundary of the wetlands and the characteristic vegetation and hydrology of each site were retrieved and compared with historical photographs dating to 1960. These comparisons were made to identify any trends in wetland attributes over time. Where needed,

photographs (10 × 15 cm [4 × 6 in]) were scanned at a file size ranging from 300 bytes to 3 megabytes and stored as digital image files in a tagged image file format (TIFF) format. Several GIS project files were created for the wetlands geospatial database and stored in computer files at BN's North Las Vegas Facility. Layouts (figures) were developed for each wetland site and printed on an Epson Stylus Pro XL™ printer.

discharge and uniform temperatures, generally from 24°C to 35°C (75°F to 95°F) (Winograd and Thordarson, 1975).

Discharges from springs, seeps, and aquifers in the region range from less than one to several thousands of gallons per minute (DOE, 1996a). Recorded flow rates from springs and seeps at NTS are all very low, generally less than 10 l/min (2.6 gal/min) (Ball, 1907; Moore, 1961; Thordarson and Robinson, 1971; DOE, 1988; Lyles *et al.*, 1990; Ingraham *et al.*, 1991). Moore (1961) reported estimates of discharge rates for eight springs on the NTS taken from 1958 to 1960. The highest discharge rates were observed at Cane Spring (7.6-11.4 l/min [2-3 gal/min]) and Whiterock Spring (3.8-7.6 l/min [1-2 gal/min]). The other six NTS springs (Captain Jack, Oak, Rainier, Tippipah, Topopah, and Tub) have recorded maximum discharge rates less than 1.7 l/min (0.4 gal/min) (Moore, 1961; Lyles *et al.*, 1990).

Cane and Whiterock springs are two of the most studied springs, and the variation in their flow rates over time has been examined. Between 1981 and 1988, discharges varied from 1.1 to 6.0 l/min (0.3 to 1.5 gal/min) at Cane Spring and from 0.5 to 4.4 l/min (0.1 to 1.2 gal/min) at Whiterock Spring (Ingraham *et al.*, 1991; Lyles *et al.*, 1990). Such changes in discharge rates are most likely affected by changes in the volume of local precipitation. Recharge of the perched water tables that feed NTS springs occurs by infiltration of rain or snowmelt through unsaturated, fractured rock. Two mechanisms of infiltration are

Lyles *et al.* (1990) recorded water temperature, electrical conductivity, pH, selected cations, and anions at Cane Spring and Whiterock Spring to examine how precipitation events influence temporal water chemistry of springs. Romney and Greger (1992) measured selected cations and 20 mineral elements in spring water from ten sites on the NTS. Stetzenbach (1995) also measured selected anions, cations, and 73 trace metals at three NTS springs.

Moore (1961) measured radioactivity at eight springs on the NTS. Beginning in 1965, DOE began long-term monitoring of radioactivity in four to eight springs on the NTS (Lewis *et al.*, 1965). This monitoring still occurs annually (Davis *et al.*, 1996). Taylor and Giles (1979) measured the uptake of tritium and gamma-emitting radionuclides by algae in eight NTS springs. All of these investigations have reported negligible levels of radioactivity, commensurate with natural background levels or indicative of regional and global fallout as a result of historic aboveground nuclear testing.

#### **4.1.3 Vegetation**

No systematic inventory of vascular or aquatic plants has been conducted at springs on the NTS. Past botanical surveys have been conducted at only a few of the known springs. They provide lists of species present at or near springs, but the location of hydrophytic

seven other NTS springs. Fifty-two individual species were documented, of which 29 were found at only a single spring. Allred *et al.* (1963) surveyed four NTS springs (Cane, Tippipah, Topopah, and Whiterock) and listed 33 species of vascular plants located at or near these springs. Beatley (1976) lists 35 species of plants occurring at or near six springs on the NTS.

#### 4.1.4 Wildlife Use

The most thorough surveys of selected NTS springs to record the presence of both invertebrates and vertebrates were conducted in the 1960s by researchers from Brigham Young University (Allred *et al.*, 1963; Jorgensen and Hayward, 1965). Of the four springs surveyed, Cane Spring was studied most extensively. Allred *et al.* (1963) recorded 5 species of lizards, 5 species of mammals, 18 species of insects, and 70 species of birds at Cane Spring. Hayward *et al.* (1963) reported 35 species of birds from Cane Spring and only 7, 2, and 3 species of birds from Tippipah, Topopah, and Whiterock springs, respectively. Castetter and Hill (1979 and unpublished field notes) reported 45 species of birds observed at Cane Spring from 1975 to 1977.

Giles (1976) examined eight NTS springs to assess the potential costs of improving the availability of water for wildlife at the springs and provided some anecdotal comments on wildlife use at springs. Most wildlife species which were noted as using the sites included mule deer (*Odocoileus hemionus*), mourning doves (*Zenaida macroura*), black-tailed jackrabbits (*Lepus californicus*), coyotes (*Canis latrans*), and mountain lions (*Felix concolor*). In 1987, researchers from the University of California at Los Angeles were funded by DOE and began to collect wildlife use data from nine springs on the NTS. A total of 46 vertebrate species, including reptiles, birds, and mammals were observed across all nine springs (Romney and Greger, 1992; Greger and Romney, 1994a,b). An undescribed species of aquatic snail was observed at Cane Spring during these wildlife use surveys (Greger and Romney, 1994a).

#### 4.1.5 Historical Use

Worman (1969) has provided the most extensive historical account of human activities around the NTS springs. Seven springs that occur east and north of the Yucca Mountain area have been identified as important Native American sites (Stoffle *et al.*, 1990a,b; Henton and Pippin, 1988). Also, ethnobotanical studies have been conducted on the NTS which document how Native Americans in the region used numerous wetland plant species (Stoffle *et al.*, 1989). From the available literature, it is known that over 15 NTS springs, seeps, and tanks were used by either miners, homesteaders, or Native Americans. Of these, over 11 springs have been modified in an attempt to increase or contain water flow. The site-specific information on the historical usage of each NTS natural water source, obtained from these citations, is presented in the following descriptions of each study site.

3 percent and 2 percent, respectively. The total vegetated area near the two largest tanks was estimated to be about 12 m<sup>2</sup> (108 ft<sup>2</sup>). Other plants in the surrounding upland area included big sagebrush (*Artemisia tridentata*), redstem stork's bill (*Erodium cicutarium*), mormon tea (*Ephedra viridis*), cheatgrass (*Bromus tectorum*), green rabbitbrush (*Chrysothamnus viscidifloris*), flax (*Linum* sp.), and antelope bitterbrush (*Purshia tridentata*). The lower tank had very little vegetation with no hydrophytic species of plants.

#### **4.2.1.3 Hydrology**

The site consists of two large tanks and several smaller tanks that vary in size. Two of the largest tanks were each approximately 3 m by 5 m (9.8 ft by 16.4 ft) and had estimated depths of 100 cm (39 in) (Photo 4-2). The tanks fill with water primarily from




Figure 4-1 Location of Ammonia Tanks



Photo 4-2. Lower tank at Ammonia Tanks looking northeast on January 7, 1997.  
(WS346-18.TIF)

Table 4-1 Ammonia Tanks wetland vegetation as surveyed on January 7, 1997

<b>Habitat: Wash Pool</b>			
<b>Species</b>	<b>Common Name</b>	<b>Indicator Status<sup>a</sup></b>	<b>Absolute % Cover</b>
<b>Tree Layer:</b> no species			
<b>Shrub Layer:</b> no species			
<b>Herb Layer:</b>			
<i>Artemisia ludoviciana</i>	Louisiana sagewort	FACU	<b>30</b>
<i>Encelia</i> sp.	unidentified brittlebush	UNKN	<b>2</b>
<i>Juncus balticus</i>	Baltic rush	FACW	<b>8</b>
<i>Leymus cinereus</i>	basin wildrye	FACU	<b>5</b>
Percentage of dominant species that are OBL, FACW, or FAC indicator status: <u>0</u> %.			
Dominant plant species are indicated by bold Absolute % Cover values.			
<sup>a</sup> For Region 8 indicator status codes for plants, see Section 3.2.4.			
			Hydrophytic vegetation: <u>No</u>

surface flow. They are located in a narrow, rocky wash with moderate amounts of exposed bedrock upstream. No surface flow from the tanks was observed. No water quality measurements were taken.

#### 4.2.1.4 *Hydric Soils*

Field indicators of hydric soils were restricted to small accumulations of soil fines located downslope of the tank above bedrock where soil appeared to be saturated for at least seven days during the growing season, indicating the presence of hydric soils.

#### 4.2.1.5 *Determination of Jurisdictional Status*

Because of a lack of hydrophytic vegetation at Ammonia Tanks, this site would probably not be considered a jurisdictional wetland.

#### 4.2.1.6 *Wildlife Use*

Little is known of wildlife use of the area; however, deer and coyote scat and tracks were observed near the water source, suggesting use by these species.

and blackbrush (*Coleogyne ramosissima*. Largeflower suncup (Cane Spring evening primrose) was first described from Cane Spring (Beatley, 1976). Two additional wetland




Figure 4-2 Location and sketch of Cane Spring



Photo 4-4. Cave pool and outflow at  
Cane Spring on June 19, 1996. (WS344-  
19.TIF)




Photo 4-6. Dry reservoir at Cane Spring, looking east on November 14, 1996. (WS501-01.TIF)

species were reported to occur at Cane Spring in 1988 that were not observed during the 1996 field survey: common reed (*Phragmites australis*) and western honey mesquite (*Prosopis glandulosa*) (Stoffle *et al.*, 1989). Beatley (1976) stated that three small western honey mesquite trees were apparently planted at Cane Spring and that they were not known to flower.

A wetland plant survey was conducted in the drainage channel 5 m (16.4 ft) directly down slope from the cave pool on June 19, 1996. In this area, approximately 33 percent of the dominant plants observed were hydrophytic, a value too low for the area downslope from the cave pool to be considered a jurisdictional wetland (Table 4-2). The seep site, however, had a dominance of hydrophytic plants (>51 percent) when surveyed in September 1996 (Table 4-3). These hydrophytic plants included Baltic rush, southern cattail, and Goodding's willow which were growing in saturated soils. This area of about 230 m<sup>2</sup> (2,475 ft<sup>2</sup>) (Table 5-1, Section 5.0) has field indicators positive for hydrophytic vegetation.

Table 4-2 Cane Spring wetland vegetation as surveyed on June 19, 1996

<b>Habitat: Drainage Channel Below Cave Pool</b>			
Species	Common Name	Indicator Status <sup>a</sup>	Absolute % Cover
<b>Tree Layer:</b> no species			
<b>Shrub Layer:</b>			
<i>Atriplex canescens</i>	fourwing saltbush	UPL	<b>15</b>
<b>Herb Layer:</b>			
<i>Leymus cinereus</i>	basin wildrye	FACU	<b>40</b>
<i>Rumex salicifolius</i>	willow dock	FACW*	<b>15</b>
Percentage of dominant species that are OBL, FACW, or FAC indicator status: <u>33</u> %.			
Dominant plant species are indicated by bold Absolute % Cover values.			
<sup>a</sup> For Region 8 indicator status codes for plants, see Section 3.2.4.			
Hydrophytic vegetation:			<u>No</u>

Table 4-3 Cane Spring wetland vegetation as surveyed on September 9, 1996

<b>Habitat: Seep Under Willow Trees</b>			
Species	Common Name	Indicator Status <sup>a</sup>	Absolute % Cover
<b>Tree Layer:</b>			
<i>Salix gooddingii</i>	Goodding's willow	FACW	<b>90</b>
<b>Shrub Layer:</b> no species			

had not overflowed the cave dam. In November, the water depth measured was greater than 1 m (3 ft), and its maximum depth was visually estimated to be 2 m (6 ft). Water quality data were taken at the cave pool and the flow box in June, September, and November 1996. These water quality data are presented in Table 5-2 (Section 5.0).

#### **4.2.2.4 Hydric Soils**

Several field indicators of hydric soils were found at the cave pool and the seep area inside the fence and included dark (low-chroma) organic profiles, mottling, and evidence

All undescribed species of hydrobiid snail (*Pyrgulopsis* sp.) occurs at Cane Spring. From June through November 1996, the snail was found on several algal mats in the cave pool nearest the tunnel entrance. Previous field records show that the snail occurred in the east channel (Paul Greger, unpublished data, 1988) and was collected from the wooden flow gauge box in 1992. The east channel was dry and had no snails in June 1996. Other invertebrates including both ostracods and copepods occur at this spring.

### **4.2.3 Captain Jack Spring**

#### **4.2.3.1 Site Description and Historical Use**

Captain Jack Spring (Figure 4-3) occurs in a rocky, remote area at the northeastern end of the Eleana Range on steep slopes with a southeast-facing aspect. Native American cultural sites, including two rock shelters and various other artifacts, are located at short distances from the spring (Worman, 1969). The area was occupied and named after an early Native American explorer who carried mail from Utah to the Groom Mine (Worman, 1969). Livestock was kept in the area as evidenced by the presence of an old corral. Remnants of old livestock watering tanks and old pipes occur about 30 m (98 ft)




Figure 4-3 Location of Captain Jack Spring

below the spring pool. The old pipes (not functional) apparently fed water to the tanks for livestock. After flood damage, new pipes were installed and a new watering tank was bolted to the bedrock (Smith *et al.*, 1979).

Water flows from the base of a narrow rocky box canyon which is about 5 m (16.4 ft) wide by 15 m (49.2 ft) long. The water forms a pool 61 × 76 cm (24 × 30 in) which is about 20 cm (8 in) deep (Photo 4-7). This pool drains downslope through a channel. This small stream of water was about 30 m (131 ft) long and 20 cm (8 in) wide. The drainage channel is at times thickly vegetated with aquatic plants (Photo 4-8). A small herd of about 20 feral horses occupies the area (Photo 4-9) (Greger and Romney, 1994b). Use of the spring by horses varies with the season, and heavy grazing and trampling by horses results in seasonal reductions in the absolute cover of wetland vegetation at the site (Photo 4-10).

#### **4.2.3.2 Hydrophytic Vegetation**

Captain Jack Spring occurs in typical pinyon-juniper habitat where localized patches of Gambel's oak (*Quercus gambelii*) are common around the base of rocky ledges. Upland species include Louisiana sagewort, big sagebrush, foxtail brome, cheatgrass, mormon tea, eastern Mojave buckwheat (*Eriogonum fasciculatum*), Utah juniper (*Juniperus osteosperma*), basin wildrye, singleleaf pinyon (*Pinus monophylla*), and bluegrasses (*Poa* spp.). Wetland plant species in the area around the spring and within the drainage channel include seep monkeyflower (*Mimulus guttatus*), biennial cinquefoil (*Potentilla biennis*), willow dock (*Rumex salicifolius*), water speedwell (*Veronica anagallis-aquatica*), and bridge penstemon (*Penstemon rostriflorus*).

On June 19, 1996, the spring drainage channel was nearly denuded of aquatic vegetation for most of its length coincident with heavy horse usage. On September 19, 1996, vegetation had regrown and there was extensive growth of aquatic vegetation throughout the total length of the drainage channel and the pool. This vegetated area was approximately 30 m<sup>2</sup> (323 ft<sup>2</sup>) (Photo 4-8). A wetland vegetation survey was conducted in the drainage channel of Captain Jack Spring about 25 m (82 ft) downslope of the spring pool. Results showed that 100 percent of the dominant species observed were hydrophytic plants indicating that field indicators for hydrophytic vegetation are present at Captain Jack Spring (Table 4-4).

#### **4.2.3.3 Wetland Hydrology and Water Quality**

Areas observed to have field indicators of wetland hydrology included the spring pool and the drainage channel below the pool. Water flow rate was approximately 0.9 l/min (0.2 gal/min) on September 10, 1996 (Table 5-1, Section 5.0). The total inundated area at Captain Jack Spring was about 7 m<sup>2</sup> (75 ft<sup>2</sup>) during September 1996. The spring pool is less than 0.5 m<sup>2</sup> (4.9 ft<sup>2</sup>) in surface area and contains an estimated volume of about 70 L (18.5 gal). Surface water and saturated soils were present at Captain Jack Spring on both visits during June and September 1996. Water quality data were taken during both visits and are presented in Table 5-2 (Section 5.0).

(WS340-18.TIF)

8. A swimming spring catfish at Captain Jack Spring on September 10, 1996.



Photo 4-10. Horse damage at Captain Jack Spring on November 21, 1988. (WS104-04.TIF)

where the soils appeared to be saturated for seven or more days during the growing season, indicating the presence of hydric soils.

#### **4.2.3.5 Determination of Jurisdictional Status**

Two areas at Captain Jack Spring (the spring pool and the narrow drainage channel below the pool) would probably qualify as jurisdictional wetlands because they had field indicators for all three required parameters: hydrophytic vegetation, wetland hydrology, and hydric soils.

#### **4.2.3.6 Wildlife Use**

This spring is commonly used by coyotes, feral horses, mountain lions, mule deer, and large numbers of upland game birds such as chukar, Gambel's quail, and mourning doves. Sixteen or more species of passerine birds have been recorded using the spring habitat (Table 5-4, Section 5.0). Raptors are also common in this area, including the Cooper's hawk (*Accipiter cooperii*) and the sharp-shinned hawk (*Accipiter striatus*). Tiny fresh-water crustaceans such as ostracods and copepods are common in the spring pool.

## 4.2.4 Cottonwood Spring

### 4.2.4.1 Site Description and Historical Use

Cottonwood Spring is located northwest of Calico Hills about 1 km (0.6 mi) west of Fortymile Canyon (Figure 4-4). This spring occurs in an east-facing wash at the top of a steep (40 percent) slope. It flows from fractures in rock ledges at approximately a 1,292-m (4,240-ft) elevation and is visible from a distance because three cottonwood trees (*Populus fremontii*) occur at the spring (Photo 4-11). The spring is marked on the Topopah Spring NW USGS 7.5-Minute Series quadrangle map (1961), but is not named. This spring is the only site on the NTS where a cottonwood tree exists; therefore, the spring was named "Cottonwood Spring." The spring was used by Native Americans, as evidenced by a temporary camp site which is located just above the spring on the ridge face at 1,310 m (4,300 ft). This site has three rock shelters and one rock alignment (Henton and Pippin, 1988). Prospecting and mining occurred in Fortymile Canyon near this spring. A surviving Forty-Niner emigrant claimed to have found some ore in Fortymile Canyon near a spring with a cottonwood tree close to the junction of several Native American trails (Stoffle *et al.*, 1990a). A prospector set out to find the spring in 1880 and was attacked by Native Americans and driven away (Stoffle *et al.*, 1990a).

There appears to be little evidence of human disturbance at this spring. A few pieces of iron pipe were located in the wash about 100 m (328 ft) below the spring, suggesting that water was piped down the wash. A few pieces of metal rebar were also found near the cottonwood trees.

### 4.2.4.2 Hydrophytic Vegetation

A wetland vegetation survey was conducted on December 12, 1996. Within the sampling area (observation point), 66 percent of the dominant plants were hydrophytic species indicating that hydrophytic vegetation was present at Cottonwood Spring (Table 4-5). Seep monkeyflower was the dominant species growing throughout the entire habitat; however, most of these individuals were very young plants of 1 cm (0.5 in) in height or less (Photo 4-12). Old flowering stalks of last year's plants remained. Mosses and western goldfern (*Pentagramma triangularis*) were also widely distributed throughout the habitat. Plant species along the border of the delineated wetland area included wormwood (*Artemisia dracuncululus*), Louisiana sagewort, and New Mexico thistle (*Cirsium neomexicanum*). Plants observed in the upland areas surrounding the drainage channels included green rabbitbrush, needle-leaf rabbitbrush (*Ericameria teretifolia*), eastern Mojave buckwheat, mormon tea, Cooper's heathgoldenrod (*Ericameria cooperi*), and Mexican bladdersage (*Salazaria mexicana*). An approximate 40-m<sup>2</sup> (430-ft<sup>2</sup>) area near the cottonwood trees had accumulated deep soils, but the soils were not saturated. Roots from the trees appeared to extend into the adjacent spring pool area where saturated soils were present. A small pile of runoff debris in the dry wash near the cottonwoods suggested previous surface water flow through this area. The surface area delineated by hydrophytic plants was estimated to be approximately 130 m<sup>2</sup> (1,399 ft<sup>2</sup>) (Table 5-1, Section 5.0).




Figure 4-4 Location and sketch of Cottonwood Spring





Photo 4-12. Vegetation on rock face at Cottonwood Spring on January 8, 1997. (WS347-09.TIF)

habitat were shallow, about 3 cm (1 in) deep. The seep area and two channels near the spring had surface flow in December 1996 and formed a confluence about 70 m (230 ft) below the cliff face (Figure 4-4). These channels cut through rock and formed pools of various size and depth. They vary from about 15 cm (6 in) to 2 m (6 ft) in width and 3 cm (1 in) to about 25 cm (10 in) in depth (Photo 4-14). The flow rate measured in December 1996 in the wash below the confluence of the two channels was approximately 1  $\ell$ /min (0.3 gal/min) (Table 5-1, Section 5.0). The area of surface inundation was about 90 m<sup>2</sup> (969 ft<sup>2</sup>). Water quality data were taken in January 1997 and are presented in Table 5-2 (Section 5.0).



#### **4.2.4.4 Hydric Soils**

Hydric soils appear to be present at Cottonwood Spring and were confined to the seep below the spring pool, and in pools and saturated soils within the two drainage channels. These soils appeared to be saturated for at least seven days during the growing season, indicating the presence of hydric soils.

An area with deeper soils directly under the three cottonwood trees did not appear to have hydric soils. Soils did not appear to be saturated within 61 cm (2 ft) of the surface; however, no soil pits were dug at this site.

#### **4.2.4.5 Determination of Jurisdictional Status**

Several areas around Cottonwood Spring would probably be considered jurisdictional wetlands because they have field indicators of all three required wetland parameters: hydrophytic vegetation, wetland hydrology, and hydric soils. These areas include the spring pool, the seep or saturated area below the spring pool, and intermittent pools or wet areas within the two drainage channels located about 70 m (230 ft) upslope and about 150 m (492 ft) downslope from the confluence of the drainage channels.

#### **4.2.4.6 Wildlife Use**

No previous monitoring of wildlife use has been conducted at this spring. Little is known about wildlife use of the spring. Mule deer scat was observed in the vicinity of the spring at the time of the wetland survey. Aquatic invertebrates observed in the spring pool in December 1996 included crustaceans (ostracods and copepods).

### **4.2.5 Coyote Spring**

#### **4.2.5.1 Site Description and Historical Use**

Coyote Spring (Figure 4-5) is located about 3 km (2 mi) southeast of Hampel Hill and approximately 3 to 5 km (2 to 3 mi) southwest of Frenchman Flat. The site (Photo 4-15) is within a wash in an area that is distant from any roads and shows no evidence of disturbance by man.

#### **4.2.5.2 Hydrophytic Vegetation**

A wetland vegetation survey of Coyote Spring was conducted on September 4, 1996. Coyote Spring consists of three seep areas (Figure 4-5) that either currently or previously supported wetland vegetation. In September, only one of these three areas was dominated by hydrophytic vegetation (Table 4-6, Photo 4-16). This area was the furthest downslope and was dominated by inland saltgrass (*Distichlis spicata*), a wetland plant species which covered a surface area of about 160 m<sup>2</sup> (1,722 ft<sup>2</sup>). A second seep area is about 40 m<sup>2</sup> (430 ft<sup>2</sup>) in size and is located on the adjacent hill side. It was dominated by common kochia (*Kochia scoparia*) and soils there were moist and dark-colored. No vegetation occurred on the third area which had dark-colored and moist soils and was located west of the other seep areas.




Figure 4-5 Location and sketch of Coyote Spring, Pavits Spring, and Tupapa Seep and sketch of Coyote Spring



Photo 4-16. Closeup of vegetation at Coyote Spring on September 4, 1996. (WSS04-01.TIF)

Hydrophytic vegetation: Yes

Other upland plants in the area included shadscale saltbush (*Atriplex confertifolia*), largeflower suncup (Cane Spring evening primrose), bottlebrush squirreltail (*Elymus elymoides*), and basin wildrye.

#### **4.2.5.3 Wetland Hydrology and Water Quality**

The only area observed to have field indicators of wetland hydrology was the seep area farthest downslope, although no standing water was observed at this site. Seasonal water availability at Coyote Spring is poorly understood; however, the presence of facultative wetland vegetation (inland saltgrass) and dark-colored soils are evidence of prior wetland hydrology. The unvegetated area of 6 m<sup>2</sup> (65 ft<sup>2</sup>) located in the wash also had dark-colored and wet surface soils. Observations made in September 1996 may not be representative of conditions during wetter years. No water quality data were collected at the time of the field survey.

#### **4.2.5.4 Hydric Soils**

Field indicators for hydric soils appeared to be present in the area dominated by inland saltgrass (i.e., the seep area furthest down slope). These indicators included dark-colored soils which appeared to have been saturated for periods of at least seven days during the growing season. No soil pits were dug. The other two seep areas also had dark-colored moist soils, but did not have evidence of saturated soils.

#### **4.2.5.5 Determination of Jurisdictional Status**

Coyote Spring has one area that would probably qualify as a jurisdictional wetland (the area dominated by inland saltgrass) because it had all three required parameters:

narrow section of the canyon with steep barren slopes on either side (Photo 4-17). There is no reported evidence of human occupation or use of this site.

#### **4.2.6.2 Hydrophytic Vegetation**

A wetland vegetation survey of Fortymile Canyon Tanks was conducted on February 12, 1997. Most of the tanks at this site have no vegetation associated with them. One area did have some limited soil and vegetation (Photo 4-18). Cover on this area was low at approximately 11 percent (Table 4-7). Louisiana sagewort and foxtail brome dominated the site. One other species, seep monkeyflower, occurred at the site, but averaged only 1 percent cover. No other species were found at this site. Because only 33 percent of the species identified from the observation point were classified as obligate or facultative wetland species, the site would be classified as not having hydrophytic vegetation. Plant species observed in the upland area were Stansbury cliffrose (*Purshia stansburiana*), skunkbush sumac (*Rhus trilobata*), green rabbitbrush, roundleaf rabbitbrush (*Ericameria teretifolia*), mormon tea, Nevada jointfir (*Ephedra nevadensis*), and big sagebrush.

#### **4.2.6.3 Hydrology**

This site consists of seven water tanks located in bedrock and a small seep at the head of the tanks. The tanks are located in a narrow rocky wash and they vary in size. The two largest ones were approximately 1 m by 3 m (3 ft by 9 ft), and the water in this tank was about 20 cm (8 in) deep. All tanks were filled with water at the time of the survey in February 1997. The estimated total area of surface water was 8 m<sup>2</sup> (86 ft<sup>2</sup>) (Table 5-1, Section 5.0). Bedrock is exposed about 30 m (98 ft) up both sides of the canyon, which




Figure 4-6 Location of Fortymile Canyon Tanks



Photo 4-18. Vegetation at Fortymile  
Canyon Tanks on February 12, 1997.  
(40MILE2.TIF)

survey, three golden eagles were observed soaring in the canyon above the tanks.

the pond measures about 10 × 40 m (33 × 131 ft) (Photo 4-19).

#### **4.2.7.2 *Hydrophytic Vegetation***

From the survey observation point located within the pond area, 100 percent of the dominant plants were wetland species (Baltic rush; Table 4-8) indicating that hydrophytic vegetation was present at this site. Baltic rush, a facultative wetland species, was the only plant species recorded at the Gold Meadows Spring observation point during July of 1996. This species covered an area about 45 m<sup>2</sup> (484 ft<sup>2</sup>). Plant species bordering the perimeter of the pond (then dry) were primarily big sagebrush and basin wildrye. The upland habitat was dominated by the trees Utah juniper and singleleaf pinyon.

#### **4.2.7.3 *Wetland Hydrology and Water Quality***

The only area observed to have field indicators of wetland hydrology was the ephemeral pond. During the winter and spring of most years, water in the pond is present (based on observations in 1989 to 1996). By summer, the pond usually dries up (Photo 4-20); however, in 1992 the pond remained all year. Although surface water was not present at the time of the field survey on July 22, 1996, previous field observations of ponded water and water lines on rocks in the pond area indicated that the site had wetland hydrology. No water quality data were able to be taken at the time of the 1996 survey.

#### **4.2.7.4 *Hydric Soils***

Soils at this site were shallow, about 20 cm (8 in) deep and had a dark-colored surface layer (probably because of high levels of organic matter). No evidence of soil mottling was observed. This site may have been periodically dug out by man to deepen the pond, and excavated soils were probably used to build up the existing berm. Therefore, an atypical situation occurred where hydric soils have been removed or disturbed. At the time of the survey, no saturated soils were observed (Table 5-1, Section 5.0); however, because the pond soils appeared to have been flooded or saturated for at least seven days during the growing season of each year, it seems that the site had hydric soils. This estimated area of saturated soils coincides with the area where Baltic rush was found (45 m<sup>2</sup> [484 ft<sup>2</sup>]).




Figure 4-7 Location of Gold Meadows Spring



Photo 4-20 Dry pond at Gold Meadows Spring on December 10, 1990 (WS213-23.TIF)




Figure 4-9 Location of Oak Spring




Figure 4-8 Location of John's Spring




Photo 4-22 Wetland vegetation at John's Spring on February 24, 1997 (JOHNS2.TIF)

mormon tea, singleleaf pinyon, desert bitterbrush (*Furcraea glauca*), and Gambel's oak. The total area of the wetland habitat dominated by hydrophytic species was estimated to be about 50 m<sup>2</sup> (538 ft<sup>2</sup>) (Table 5-1, Section 5.0) and was restricted to seep areas along the rock face and the ledge pool below the rock face.

#### **4.2.8.3 Wetland Hydrology and Water Quality**

Areas observed to have field indicators of wetland hydrology included seeps located in the rock face and a surface pool at the base of the rock face. Water seeps out of the rock face in numerous locations across a distance of about 25 m (82 ft) and flows down a steep slope for about 5 m (16 ft). Flow rate was measured at 0.4 l/min (0.01 gal/min) in December 1996 (Table 5-1, Section 5.0). A 2-cm- (1-in)-deep surface pool exists on a relatively flat area at the base of the rock face. The surface area of this pool is about 5 m<sup>2</sup> (54 ft<sup>2</sup>) and is covered with a dense growth of seep monkeyflower (Photo 4-22). Water quality measurements were not taken at this site.

#### **4.2.8.4 Hydric Soils**

Hydric soils at John's Spring were confined to the seep areas at the base of the rocky cliff and the ledge pool, and comprised about 50 m<sup>2</sup> (538 ft<sup>2</sup>). Soils in these areas appeared to be saturated for at least seven days during the growing season, indicating the presence of hydric soil.

... and the large pool at the base of the rock face which covered an area about 50 m<sup>2</sup> (538 ft<sup>2</sup>).

#### **4.2.8.6 Wildlife Use**

Little is known about wildlife use of the spring because of limited study. Mule deer use the area as indicated by the presence of scat near the spring. Migratory passerine birds such as dark-eyed juncos (*Junco hyemalis*) were observed drinking from the spring on December 18, 1996. Many species of passerine birds probably benefit from drinking at this site during summer months.

#### **4.2.9 Oak Spring**

##### **4.2.9.1 Site Description and Historical Use**

Oak Spring is located southwest of Oak Spring Butte (Figure 4-9) in an oak grove on a hillside (Photo 4-23) near the northern NTS boundary. Native Americans were reported to

saltbush, cheatgrass, green rabbitbrush, mormon tea, singleleaf pinyon, desert bitterbrush, and Gambel's Oak.

From the survey observation point, located at the spring pool, nearly 100 percent of the dominant plant species were wetland species (sandbar willow) indicating that hydrophytic vegetation was present (Table 4-10). Beatley (1976) reported five additional wetland plant species from Oak Spring (Table 5-3, Section 5.0) that were not detected during the survey: sturdy sedge (*Carex alma*), beardless wildrye (*Leymus [Elymus] triticoides*), seep monkeyflower, and beardless rabbitsfoot grass (*Polypogon viridis*). All of these species are obligate or facultative wetland species. It is probable that some of these species reestablish during periods of greater spring flow and moist soil conditions. Seeds of some of these plants might readily disperse to the site from plants growing at John's Spring about 250 m (820 ft) away.




Figure 4-9 Location of Oak Spring



Photo 4-24 Closeup of wetland vegetation at Oak Spring on November 4, 1996 (WS343-16.TIF)

Table 4-10 Oak Spring wetland vegetation as surveyed on November 4, 1996

<b>Habitat: Spring Outflow</b>			
<b>Species</b>	<b>Common Name</b>	<b>Indicator Status<sup>a</sup></b>	<b>Absolute % Cover</b>
<b>Tree Layer:</b> no species			
<b><i>Salix exigua</i></b>	sandbar willow	FACW	<b>100</b>
<b>Herb Layer:</b> no species			
Percentage of dominant species that are OBL, FACW, or FAC indicator status: <u>100</u> %.			
Dominant plant species are indicated by bold Absolute % Cover values.			
<sup>a</sup> For Region 8 indicator status codes for plants, see Section 3.2.4.			
Hydrophytic vegetation: <u>Yes</u>			

An adjacent transitional area between the jurisdictional wetland and upland areas was comprised of basin wildrye and smooth sumac. Within this transitional area, basin wildrye constituted about 30 percent cover and smooth sumac about 5 percent.

#### **4.2.9.3 Wetland Hydrology and Water Quality**

Field indicators of wetland hydrology were restricted to one area during the wetland survey in November 1996. This area was a small pool of surface water less than 1 m<sup>2</sup> (11 ft<sup>2</sup>) in area and about 2.5 cm (1 in) deep. Flow out of this pool was measured at 0.4 l/min (0.1 gal/min) in December 1996 (Table 5-1, Section 5.0). No water quality measurements were taken at the site.

#### **4.2.9.4 Hydric Soils**

Field indicators for hydric soils were restricted to saturated soils at the surface pool at Oak Spring. Soils were fairly shallow and limited in areal extent. Soil mottling was not observed in the soil pit that was dug; however, soils had a moderate amount of dark organic matter and were saturated for what appeared to have been greater than seven days during the growing season, indicating the presence of hydric soils. The spring had apparently been excavated and soils may have represented an atypical situation.

#### **4.2.9.5 Determination of Jurisdictional Status**

Only the small area at Oak Spring occupied by sandbar willow met the criteria to be considered as a jurisdictional wetland because it had field indicators for all three required

located at the bottom of a hill and adjacent to a drainage channel. No water quality measurements were taken at this site.

#### **4.2.10.4 *Hydric Soils***

Field indicators for hydric soils were not observed at this site. Soil pits were not dug because of the potential to disturb the small pool site. Soils at this site are poorly developed, with little organic matter.




Figure 4-10 Location of Pavits Spring



Photo 4-26 Wash vegetation around Pavits Spring looking northeast on September 9, 1988  
(WS036-14.TIF)



Photo 4-28 Dry pool at Pavits Spring on September 4, 1996 (WS502-01.TIF)

Table 4-11 Pavits Spring wetland vegetation as surveyed on September 4, 1996

<b>Habitat: Dry Wash Pool</b>			
Species	Common Name	Indicator Status <sup>a</sup>	Absolute % Cover
<b>Tree Layer:</b> no species			
<b>Shrub Layer:</b>			
<i>Ericameria (Chrysothamnus) nauseosa</i>	rubber rabbitbrush	NL	40
<b>Herb Layer:</b>			
<i>Dactylus glomerata</i>	orchardgrass	FACU	10
<i>Sporobolus airoides</i>	alkali sacaton	FAC-	10
Percentage of dominant species that are OBL, FACW, or FAC indicator status: <u>33</u> %.			
Dominant plant species are indicated by bold Absolute % Cover values.			
<sup>a</sup> For Region 8 indicator status codes for plants, see Section 3.2.4.			
			Hydrophytic vegetation: <u>No</u>

#### 4.2.10.5 Determination of Jurisdictional Status

Because of the lack of hydrophytic vegetation and hydric soils, it is probable that Pavits Spring would not be considered a jurisdictional wetland.

#### 4.2.10.6 Wildlife Use

Wildlife which use Pavits Spring include desert cottontails (*Sylvilagus audobonii*), coyotes, Gambel's quail, hummingbirds, and mule deer. Invertebrate groups recorded from water samples from Pavits Spring in 1988 (Greger, unpublished notes) include chironomids (aquatic midges), cladocerans (water fleas), oligochaetes (aquatic earthworms), and nematodes (round worms).

#### 4.2.11 Rainier Spring

##### 4.2.11.1 Site Description and Historical Use

Rainier Spring is located in a wash adjacent to a dirt road approximately 300 m (984 ft) north of E Tunnel Portal (Figure 4-11, Photo 4-29). This site was apparently used in the




Figure 4-11 Location of Rainier Spring



Photo 4-30 Closeup of vegetation at Rainier Spring on February 20, 1997 (RAINIER2.TIF)

1900s to water livestock, as evidenced by the presence of metal water tanks. Several metal pipes exist in the wash. One outflow pipe was inserted into the rocks about 10 m (33 ft) above several water guzzlers which were made from 55-gallon (208-ℓ) drums and were filled partially with sediment (Photo 4-30). A wooden post was inserted into the wash sediment apparently to secure or stabilize one of the tanks. An overturned stock tank about 1.8 m (6 ft) long × 0.6 m (2 ft) wide was located about 10 m (33 ft) further down the wash from the drums.

#### 4.2.11.2 *Hydrophytic Vegetation*

The wetlands survey performed on December 18, 1996, showed that basin wildrye, a nonhydrophytic plant, was the only dominant plant species present in the wash at Rainier Spring and comprised about 30 percent of the cover in the wash (Table 4-12). The area of the wash where basin wildrye occurred was about 22 m<sup>2</sup> (237 ft<sup>2</sup>). Plant species in the surrounding upland included big sagebrush, mormon tea, rubber rabbitbrush, Utah juniper, and singleleaf pinyon. It was concluded that hydrophytic vegetation did not occur at this site at the time of the survey, although basin wildrye is indicative of mesic areas elsewhere on the NTS.

Table 4-12 Rainier Spring wetland vegetation as surveyed on December 18, 1996

<b>Habitat: Wash</b>			
<b>Species</b>	<b>Common Name</b>	<b>Indicator Status<sup>a</sup></b>	<b>Absolute % Cover</b>
<b>Tree Layer:</b> no species			
<b>Shrub Layer:</b> no species			
<b>Herb Layer:</b>			
<i>Leymus cinereus</i>	basin wildrye	FACU	<b>30</b>
Percentage of dominant species that are OBL, FACW, or FAC indicator status: <u>0</u> %.			
Dominant plant species are indicated by bold Absolute % Cover values.			
<sup>a</sup> For Region 8 indicator status codes for plants, see Section 3.2.4.			
Hydrophytic vegetation: <u>No</u>			

#### 4.2.11.3 *Hydrology*

Rainier Spring was reported to have water on September 18, October 4, and October 15, 1957 (Moore, 1961). This site was sampled for water quality and radioactivity at that time. Rainier Spring was reported to be dry on November 10, 1960 (Moore, 1961). No flow was observed from the outflow pipe in 1996. A dirt road leading to B Tunnel exists within 20 m (66 ft) of the spring site. The construction of this road may have influenced

drainage and recharge patterns of the area by intercepting rainfall and directing water flow down the road. This could have decreased spring discharge. Surface water was absent at Rainier Spring on December 18, 1996, and no water quality measurements were taken.

#### **4.2.11.4 Hydric Soils**

No field indicators of hydric soils (such as surface water or saturated soils) were observed at Rainier Spring.

#### **4.2.11.5 Jurisdictional Wetland Determination**

Rainier Spring would probably not be considered a jurisdictional wetland because it lacked all three characteristics of a jurisdictional wetland: hydrophytic vegetation, wetland hydrology, and hydric soils.

#### **4.2.11.6 Wildlife Use**

The Rainier Spring site has little value for wildlife use because it lacks surface hydrology. No wildlife or their sign were observed at the site during the wetland survey.

### **4.2.12 Reitmann Seep**

#### **4.2.12.1 Site Description and Historical Use**

Reitmann Seep was also known as Green Spring (Giles, 1976); however, this site is unnamed (i.e., referred to as “spring”) on the Paiute Ridge USGS 7.5-Minute Series quadrangle map (1986). The name “Reitmann Seep” has been commonly used in past DOE reports. It is about 3 km (2 mi) east of Yucca Flat and southwest of Slanted Buttes (Figure 4-12, Photo 4-31). The seep forms a very small pool (Photo 4-32) and contains about 23 ℓ (6 gal) of water throughout the year (Giles, 1976). The pool and surrounding vegetation have an area of about 1 m<sup>2</sup> (11 ft<sup>2</sup>). The slope above Reitmann Seep is moderately steep and heavy rainfall commonly fills the pool with sediment. Little information on historical use of Reitmann Seep by humans is known. The seep was improved by man during recent times, apparently for the purpose of supplying water to wildlife. A 55-gallon (208-ℓ) drum (now heavily rusted) is cut open on one side and buried flush with the soil surface. It was installed about 10 m (33 ft) downslope from the spring pool and is fed water through a pipe. The pipe from the pool to the drum is buried in a narrowly excavated channel.

#### **4.2.12.2 Hydrophytic Vegetation**

Wetlands vegetation at Reitmann Seep includes Parish’s spikerush (*Eleocharis parishii*) and annual rabbitsfoot grass. From the survey observation point at the spring pool, about 66 percent of the dominant plant species around the seep were wetland species (Table 4-13), indicating that hydrophytic vegetation was present at Reitmann Seep. This area of




Figure 4-12 Location of Reitmann Seep



Photo 4-32 Pool at Reitmann Seep on June 2, 1989 (WS110-23.TIF)

Table 4-13 Reitmann Seep wetland vegetation as surveyed on June 19, 1996

<b>Habitat: Spring Pool</b>			
Species	Common Name	Indicator Status <sup>a</sup>	Absolute % Cover
<b>Tree Layer:</b> no species			
<b>Shrub Layer:</b> no species			
<b>Herb Layer:</b>			
<i>Bromus rubens</i>	foxtail brome	FACU	10
<i>Eleocharis parishii</i>	Parish's spikerush	OBL	50
<i>Polypogon monspeliensis</i>	annual rabbitsfoot grass	FACW+	40
Percentage of dominant species that are OBL, FACW, or FAC indicator status: <u>66</u> %.			
Dominant plant species are indicated by bold Absolute % Cover values.			
<sup>a</sup> For Region 8 indicator status codes for plants, see Section 3.2.4.			
Hydrophytic vegetation:			<u>No</u>

wetland vegetation is very small (about 1 m<sup>2</sup> [11 ft<sup>2</sup>]). Vegetation of the upland plant community adjacent to the wetland included fourwing saltbush, foxtail brome, blackbrush, and Joshua tree (*Yucca brevifolia*).

#### 4.2.12.3 Wetland Hydrology and Water Quality

Surface water is present throughout the year at Reitmann Seep in a small pool and a guzzler whose combined area is approximately 1.5 m<sup>2</sup> (12 ft<sup>2</sup>). The pool is located at the base of a fractured, rocky hillside in an earthen basin that appears to have been formed by runoff water from the hillside above the pool. Flow rates measured from an existing pipe inserted into the ground at the spring were low and varied from 0.03 l/min (0.007 gal/min) in September 1996 to 0.2 l/min (0.05 gal/min) in November 1996 (Table 5-1, Section 5.0). Water quality measurements were taken in June, July, September, and November 1996. Data are presented in Table 5-2 (Section 5.0).

#### 4.2.12.4 Hydric Soils

Hydric soils were restricted to the saturated soils at the spring pool at Reitmann Seep. These soils appear to be saturated for more than seven days during the plant growing season, indicating the presence of hydric soil. Soils are very black and appear high in decaying organic matter, most of which appears to be plant litter that has blown into the pool from adjacent upland vegetation.

below the tank covered about 25 m<sup>2</sup> (269 ft<sup>2</sup>) surface area. Plant species in the surrounding upland area included white burrobush, Nevada jointfir, and creosotebush.

#### **4.2.13.3 Hydrology**

Surface water was present at the Rock Valley Tank on January 7, 1997, and was observed in the small rock opening measuring about 20 × 30 cm (8 × 12 in) at the base of a limestone ridge (Photo 4-34). No water flow from the rock cavity was observed. The depth of water inside the rock cavity near the surface was about 30 cm (1 ft). The water-filled cavity extended more than 1 m (3.3 ft) horizontally into the rock formation. It appears that the cavity may have been formed by a combination of standing water from runoff gradually dissolving the rock substrate, freezing and thawing of the water in the winter, and perhaps by water seeping through fractures in the rock formation. In the rock formation above the tank, several small 0.03-m<sup>3</sup> (1-ft<sup>3</sup>) depressions appear to have been formed by similar processes, and water in these depressions may seep through the rock to the tank below. Narrow erosion channels in the rock were also frequently observed in the area along fractures. No water quality measurements were taken.




Figure 4-13 Location of Rock Valley Tank







Photo 4-34 Tank opening (center) at Rock Valley Tank looking west on January 7, 1997 (WS340-20.TIF)

Worman, 1969). A gently sloping tunnel was excavated into the hillside, creating



-  Spring
-  Unimproved Road
-  Elevation Contour (50 m)
-  Primary Wash
-  Mine Tunnel or Cave Entrance



**Bechtel Nevada**

Figure 4-14 Location and sketch of Tippipah Spring




Photo 4-36 Habitat surrounding Tippipah Spring looking west on October 5, 1988 (WS501-01.TIF)

a cave approximately 9 m (30 ft) long (Giles, 1976). Spring water accumulates in the tunnel in a pool about 4 m (13 ft) from the cave entrance (Photo 4-35). The back of the tunnel is under water when the pool is full. The remaining 6 m (20 ft) of the adit are only visible when water levels in the pool are low. Water from the cave pool flows for about 40 m (130 ft) through subsurface strata and emerges in a long surface channel (Photo 4-36). The channel flows north for varying distances depending on seasonal rainfall (Figure 4-14). It appears that the upper portion of the spring channel has been excavated periodically by man judging from the mounds of dirt on either side of the channel. The upper spring channel was nearly dry during the Fall of 1990 resulting in a visible decline in cover of riparian plants (Photo 4-37). The same habitat showed recovery of the wetland vegetation during 1992 (Photo 4-38), a wet year which broke a three-year drought period.

#### **4.2.14.2 Hydrophytic Vegetation**

During June 1996, the wetland vegetation consisted of a narrow, linear corridor that extended for 170 m (558 ft). Three locations along its length were sampled: upper channel, mid-channel (Photo 4-39) and lower channel (Photo 4-40). Eleven species of wetland plants were recorded at these observation points. The dominant wetland species were Baltic rush, biennial cinquefoil, annual rabbitsfoot grass, and water speedwell (Tables 4-15 to 4-17). There were minor differences in the plant communities sampled at the three observation points. The number and composition of wetland plant species varies along the watercourse with distance from the spring source. Annual rabbitsfoot grass was present in the lower two sections and was not observed in the upper section. Biennial cinquefoil was present only in the lower channel. Other species within the jurisdictional wetland boundary were trefoil (*Lotus* sp.) and southern cattail. Louisiana sagewort and rubber rabbitbrush were encroaching on the edge of the wetland boundary. The overall wetland habitat was dominated by Baltic rush. The total surface area of this wetland, as defined by wetland vegetation, was approximately 500 m<sup>2</sup> (5,380 ft<sup>2</sup>) (Table 5-1, Section 5.0). Common upland plant species in the area included big sagebrush, blackbrush, rubber rabbitbrush, fourwing saltbush, Nevada jointfir, and desert bitterbrush.

Surveys from the observation points in the upper channel, mid-channel, and lower channel all determined that 100 percent of the dominant plant species were wetland species indicating that Tippipah Spring has hydrophytic vegetation (Tables 4-15 to 4-17). However, no vascular plants were observed at the spring pool, possibly because the pool lies within the cave where light is limited.

#### **4.2.14.3 Wetland Hydrology and Water Quality**

Field indicators of wetland hydrology were restricted to the cave pool and the three channel areas. Surface water was present at Tippipah Spring during the wetland survey in June 1996. The estimated area of surface inundation was approximately 190 m<sup>2</sup> (2,044 ft<sup>2</sup>) (Table 5-1, Section 5.0). Flow rates in the mid channel were estimated to be approximately 2.7 l/min (0.7 gal/min) on November 15, 1996. Water quality measurements were taken at the open channel pool in June, September, and November 1996. Measurements were taken at the cave pool in September and November 1996. Data are presented in Table 5-2 (Section 5.0).




Photo 4-38 Upper reach of Tippipah Spring channel showing recovery of vegetation following drought, looking north on June 4, 1992 (WS310-09.TIF)




Photo 4-39 Middle reach of Tippipah Spring channel looking southwest on June 18, 1996 (WSS41-25.TIF)




Photo 4-40 Lower reach of Tippipah Spring channel looking northwest on June 18, 1996 (WS341-24.TIF)

Table 4-15 Tippipah Spring wetland vegetation as surveyed on June 18, 1996

<b>Habitat: Upper Stretch of the Spring Channel</b>			
<b>Species</b>	<b>Common Name</b>	<b>Indicator Status<sup>a</sup></b>	<b>Absolute % Cover</b>
<b>Tree Layer:</b>			
no species			
<b>Shrub Layer:</b>			
no species			
<b>Herb Layer:</b>			
<i>Carex praegracilis</i>	clustered field sedge	FACW-	2
<i>Eleocharis palustris</i>	common spikerush	OBL	Tr
<i>Heliomeris multiflora</i> var. <i>nevadensis</i>	Nevada goldeneye	NL	Tr
<i>Juncus balticus</i>	Baltic rush	FACW	<b>49</b>
<i>Polypogon monspeliensis</i>	annual rabbitsfoot grass	FACW+	Tr
<i>Veronica anagallis-aquatica</i>	water speedwell	OBL	<b>49</b>
Percentage of dominant species that are OBL, FACW, or FAC indicator status: <u>100</u> %.			
Dominant plant species are indicated by bold Absolute % Cover values. Tr = trace, <1% absolute cover.			
<sup>a</sup> For Region 8 indicator status codes for plants, see Section 3.2.4.			
Hydrophytic vegetation: <u>No</u>			

Table 4-16 Tippipah Spring wetland vegetation as surveyed on June 18, 1996

**Habitat: Middle Stretch of Spring Channel**

Species	Common Name	Indicator Status <sup>a</sup>	Absolute % Cover
<b>Tree Layer:</b>			
no species			
<b>Shrub Layer:</b>			
no species			
<b>Herb Layer:</b>			
<i>Bromus rubens</i>	foxtail brome	FACU	Tr
<i>Bromus tectorum</i>	cheatgrass	NL	2
<i>Castilleja</i> sp.	unidentified Indian paintbrush	UNKN	Tr
<i>Deschampsia danthonioides</i>	annual hairgrass	FACW	Tr
<i>Eleocharis parishii</i>	Parish's spikerush	OBL	Tr
<i>Epilobium glaberrimum</i>	smooth willowweed	FACW	2
<i>Heliomeris multiflora</i> var. <i>nevadensis</i>	Nevada goldeneye	NL	2
<i>Juncus balticus</i>	Baltic rush	FACW	40
<i>Juncus longistylis</i>	longstyle rush	FACW+	Tr
<i>Lactuca serriola</i>	prickly lettuce	FACU	2
<i>Polypogon monspeliensis</i>	annual rabbitsfoot grass	FACW+	30
<i>Verbena bracteata</i>	bigbract verbena	FACU	2
<i>Veronica anagallis-aquatica</i>	water speedwell	OBL	20

Percentage of dominant species that are OBL, FACW, or FAC indicator status: 100 %.

Dominant plant species are indicated by bold Absolute % Cover values. Tr = trace, <1% absolute cover.

<sup>a</sup> For Region 8 indicator status codes for plants, see Section 3.2.4.

Hydrophytic vegetation: Yes

Table 4-17 Tippisah Spring wetland vegetation as surveyed on June 18, 1996

<b>Habitat: Lower Stretch of Spring Channel</b>			
<b>Species</b>	<b>Common Name</b>	<b>Indicator Status<sup>a</sup></b>	<b>Absolute % Cover</b>
<b>Tree Layer:</b>			
no species			
<b>Shrub Layer:</b>			
no species			
<b>Herb Layer:</b>			
<i>Bromus tectorum</i>	cheatgrass	NL	Tr
<i>Erodium cicutarium</i>	redstem stork's bill	NL	Tr
<i>Heliomeris multiflora</i> var. <i>nevadensis</i>	Nevada goldeneye	NL	2
<i>Juncus balticus</i>	Baltic rush	FACW	40
<i>Lactuca serriola</i>	prickly lettuce	FACU	2
<i>Polypogon monspeliensis</i>	annual rabbitsfoot grass	FACW+	10
<i>Potentilla biennis</i>	biennial cinquefoil	FAC	20
<i>Verbena bracteata</i>	bigbract verbena	FAC	1
<i>Veronica anagallis-aquatica</i>	water speedwell	OBL	25

Percentage of dominant plant species that are OBL, FACW, or FAC indicator status: 100 %.

Dominant plant species are indicated by bold values for Absolute % Cover. Tr = trace, <1% absolute cover.

<sup>a</sup>For Region 8 indicator status codes for plants, see Section 3.2.4).


**Hydrophytic vegetation: Yes**

#### 4.2.14.4 Hydric Soils

Field indicators of hydric soils were also restricted to areas at the cave pool and the three channel areas where soils appeared to be saturated for seven days or more during the plant growing season, indicating the presence of hydric soil. Several soil pits were dug to examine soils for other field indicators for hydric soils. No evidence of soil mottling was found. Because the upper spring outflow area appears to have been dug out or possibly blasted due to the presence of rock immediately adjacent to the channel, this site may represent an atypical situation for soils evaluation.

#### 4.2.14.5 Determination of Jurisdictional Status

The three channel areas at Tippisah Spring would probably qualify as jurisdictional wetlands because they had field indicators for all three required parameters: hydrophytic vegetation, wetland hydrology, and hydric soils. The cave pool lacks hydrophytic vegetation and would probably not be considered as a jurisdictional wetland. It may, however, be protected under the CWA as waters of the United States.






Figure 4-15 Location of Tongue Wash Tank





Photo 4-41 Cave opening at Tongue Wash Tank looking south on January 10, 1997 (WS340-24.TIF)



Photo 4-42 Tank inside cave at Tongue Wash Tank on January 10, 1997 (WS340-22.TIF)

Utah serviceberry (*Amelanchier utahensis*), big sagebrush, mormon tea, basin wildrye, desert almond (*Prunus fasciculata*), Stansbury cliffrose, Gambel's oak, and skunkbush sumac.

One hundred percent of the dominant plant species in the cave pool area (Table 4-18) were hydrophytic species, indicating that field indicators for hydrophytic vegetation are present at this area. At the hillside meadow sample area, only two of the four dominant plants (50 percent) were wetland species (Table 4-19). Normally, greater than 50 percent of the dominant plants must be hydrophytic to conclude that field indicators for hydrophytic vegetation are present. However, it is appropriate to conclude that such field indicators were present at the hillside meadow area because hydrophytic species at this site (the nine species categorized as FAC, FACW, or OBL [Table 4-19]) comprised 63 percent of the cover, whereas nonhydrophytic species comprised only 43 percent of the cover. It is therefore concluded that Topopah Spring has field indicators for hydrophytic vegetation



**Bechtel Nevada**




Photo 4-44 Vegetation of Topopah Spring cave pool on April 18, 1989 (WS109-15.TIF)




Photo 4-46 Hillside meadow at Topopah Spring looking north on June 20, 1996 (WS339-18.TIF)

Table 4-18 Topopah Spring wetland vegetation as surveyed on June 20, 1996

<b>Habitat: Cave Pool</b>			
<b>Species</b>	<b>Common Name</b>	<b>Indicator Status<sup>a</sup></b>	<b>Absolute % Cover</b>
<b>Tree Layer:</b> no species			
<b>Shrub Layer:</b> no species			
<b>Herb Layer:</b>			
<i>Artemisia ludoviciana</i>	Louisiana sagewort	FACU	5
<i>Bromus diandrus</i>	ripgut grass	NL	1
<i>Epilobium glaberrimum</i>	smooth willowweed	FACW	5
<i>Mimulus guttatus</i>	seep monkeyflower	OBL	<b>10</b>
<i>Polypogon monspeliensis</i>	annual rabbitsfoot grass	FACW+	1
<i>Potentilla biennis</i>	biennial cinquefoil	FAC	2
<i>Pseudognaphalium stramineum</i>	straw falsecudweed	NL	5
<i>Rumex salicifolius</i>	willow dock	FAC*	<b>15</b>
<i>Sisymbrium altissimum</i>	tall tumbled mustard	FACU	1
<i>Veronica anagallis-aquatica</i>	water speedwell	OBL	<b>20</b>
Percentage of dominant plant species that are OBL, FACW, or FAC indicator status: <u>100</u> %.			
Dominant plant species are indicated by bold Absolute % Cover values.			
<sup>a</sup> For Region 8 indicator status codes for plants, see Section 3.2.4.			
			Hydrophytic vegetation: <u>Yes</u>

Percentage of dominant plant species that are OBL, FACW, or FAC indicator status: 50 %.

Absolute % cover due to all hydrophytic species present: 63 %.

Dominant plant species are indicated by bold Absolute % Cover values. Tr = trace, <1% absolute cover.

<sup>a</sup> For Region 8 indicator status codes for plants, see Section 3.2.4.

Hydrophytic vegetation: Yes\*

\*Because the number of dominant hydrophytic species was equal to the number of dominant nonhydrophytic species, an alternative method for determining the presence of hydrophytic vegetation was used (Williams, 1992; see Section 3.2.4).

Topopah Spring comprised a much larger area of about 200 m<sup>2</sup> (2,152 ft<sup>2</sup>) (Table 5-1, Section 5.0).

Other area of standing water was estimated to be about 0.1 m (0.3 ft) deep. Flow rates at Topopah Spring, measured from the existing pipe, were very low (0.14 l/min [0.04 gal/min]) (Table 5-1, Section 5.0). Water flow was widely distributed on the hillside meadow and was not measured. Water quality measurements were taken in the spring pool in June and September 1996. Data are presented in Table 5-2 (Section 5.0).

#### **4.2.16.4 Hydric Soils**

Field indicators of hydric soils were observed at both the cave pool area and the hillside meadow and consisted of saturated or inundated soils which appeared to be inundated for seven days or longer during the plant growing season, indicating the presence of hydric soil. Two soil pits were dug to examine soils for field indicators for hydric soils. Mottling was not observed in the soil exposed from these soil pits.

#### **4.2.16.5 Determination of Jurisdictional Status**

Both the cave pool area and the hillside meadow at Topopah Spring would probably be considered jurisdictional wetlands because they had field indicators for all three required parameters: hydrophytic vegetation, wetland hydrology, and hydric soils.

#### **4.2.16.6 Wildlife Use**

Wildlife commonly using this spring include coyote, mountain lion, mule deer, raptors, and passerine birds. Upland game birds are common including chukar, Gambel's quail, and mourning dove. Extensive use of this spring is made by chukar in the summer. Biologists have observed hundreds of these birds around the cave pool during numerous visits.

### **4.2.17 Tub Spring**

#### **4.2.17.1 Site Description and Historical Use**

Tub Spring is located in a wash on a southeast-facing slope on the east side of Oak Spring Butte, north of Yucca Flat (Figure 4-17, Photo 4-47). Access to water at Tub Spring was developed by local mining operations located about 1.6 km (1 mi) southwest of the spring (Giles, 1976). After the spring was developed, it was probably used to water cattle that grazed in the region before 1950. A tunnel 0.6 to 0.9 m (2 to 3 ft) wide was excavated about 9 m (30 ft) into the hillside. Water collects in a pool within the tunnel throughout the year behind a small earthen dam at the tunnel entrance. The depth of the water is approximately 0.6 m (2 ft) (Giles, 1976). A 7.6-cm- (3-in)-diameter pipe had been installed from the tunnel down slope about 60 m (197 ft) to a large metal watering tank. In 1975, the spring and pipeline system was renovated after a rock slide plugged the pipeline (Smith *et al.*, 1978). Additionally, a wire screen was placed over the pipe entrance to prevent recurrence of the blockage. The original pipeline and watering tank




Figure 4-17 Location of Tub Spring



Photo 4-48 Vegetation around Tub Spring on August 24, 1996 (WS046-20.TIF)

were reused. In the 1980s, biologists conducting wildlife surveys at the site observed that the pipe was broken and that no water was flowing into the metal watering tank. Some time in the early 1990s, a 3.8-ℓ (1-gal) tin can was installed in the ground under the broken end of the pipe which was about 10 m (33 ft) upslope from the watering tank. Water currently runs through the pipe, fills the tin can, and overflows onto the ground forming a wetted area of about 1 m<sup>2</sup> (11 ft<sup>2</sup>) (Photo 4-48).

#### 4.2.17.2 Hydrophytic Vegetation

Only two plant species, skunkbush sumac, and an unidentified grass, were observed growing in a small moist area around the tin can (Table 4-20). Upland vegetation near this area includes desert needlegrass (*Achnatherum speciosum*), fourwing saltbush, big sagebrush, foxtail brome, cheatgrass, blackbrush, mormon tea, Nevada jointfir, Cooper's heathgoldenrod, rubber rabbitbrush, and Stansbury cliffrose. Hydrophytic vegetation was absent from this site and at the cave pool.

Table 4-20 Tub Spring wetland vegetation as surveyed on November 7, 1996

<b>Habitat: Tin Can Area</b>			
<b>Species</b>	<b>Common Name</b>	<b>Indicator Status<sup>a</sup></b>	<b>Absolute % Cover</b>
<b>Tree Layer:</b> no species			
<b>Shrub Layer:</b>			
<i>Rhus trilobata</i>	skunkbush sumac	NI	10
<b>Herb Layer:</b>			
unidentified grass		UNKN	1
Percentage of dominant plant species that are OBL, FACW, or FAC indicator status: <u>0</u> %.			
Dominant plant species are indicated by bold values for Absolute % Cover.			
<sup>a</sup> For Region 8 indicator status codes for plants, see Section 3.2.4.			
<b>Hydrophytic vegetation: <u>No</u></b>			

#### 4.2.17.3 Wetland Hydrology and Water Quality

Field indicators of wetland hydrology were limited to the cave pool and the saturated area around the tin can where surface water was present. It is likely that the area around the tin can would be dry if it had not been developed by the recent activities of man. Flow rate measured from the broken pipe on September 20, 1996 was 0.1 ℓ/min (0.03 gal/min) (Table 5-1, Section 5.0). Water quality measurements were taken in the guzzler can in June and September 1996 and are presented in Table 5-2 (Section 5.0).

#### **4.2.17.4 Hydric Soils**

Field indicators of hydric soils were observed at the cave pool and around the tin can. Soils in these areas appeared to have been saturated for more than seven days during the growing season, indicating the presence of hydric soil. However, because the soils around the can are dependent on water being delivered by the pipe, this area is considered artificially wet. No soil pits were dug due to the small size of the area and the evidence of prior disturbance. Soils appeared to be poorly developed in undisturbed areas around the site.

#### **4.2.17.5 Determination of Jurisdictional Status**

No area at Tub Spring would be considered a jurisdictional wetland because all areas lacked field indicators for hydrophytic vegetation. Although the cave pool would not be considered a jurisdictional wetland, it may be considered waters of the United States.

#### **4.2.17.6 Wildlife Use**

Chukar, coyotes, Gambel's quail, mourning doves, mountain lions, mule deer, and passerine birds drink water at the site.

#### **4.2.18 Tupapa Seep**

##### **4.2.18.1 Site Description and Historical Use**

This site was identified as "Tupapa Seep Spring" on the Camp Desert Rock USGS 7.5-Minute Series quadrangle map (1961). No spring could be found at the site and it was renamed for this report as "Tupapa Seep." It is located in a wash in a remote area about 1 km (0.6 mi) southeast of Hampel Hill and 3.2 km (2 mi) southwest of Frenchman Flat (Figure 4-18). The site appears to be unaltered by man (Photo 4-49), and no historical accounts of human activity at the spring were found.

##### **4.2.18.2 Hydrophytic Vegetation**

A vegetation survey from an observation point located within the wash was conducted at Tupapa Seep on November 7, 1996 (Photo 4-50). From the observation point only one dominant species was classified as a hydrophytic plant, foxtail barley (*Hordeum jubatum*). Cheatgrass, an upland grass, was the only other dominant species observed in the sample area (Table 4-21). Other plants in the sample area included: shadscale saltbush, rubber rabbitbrush, and Mexican bladdersage. Based on the weak showing of hydrophytic species at this site, it was concluded that the criteria for hydrophytic vegetation were lacking.

##### **4.2.18.3 Wetland Hydrology and Water Quality**

Field indicators of wetland hydrology were lacking at this site. No evidence of surface water or saturated soils was detected during visits to Tupapa Seep in September and November 1996. This seep is probably seasonally intermittent in flow and appears to




Figure 4-18 Location of Tupapa Seep



Photo 4-50 Closeup of vegetation at Tupapa Seep looking east on November 4, 1996 (WS505-03.TIF)

Table 4-21 Tupapa Seep wetland vegetation as surveyed on November 7, 1996

<b>Habitat: Wash</b>			
Species	Common Name	Indicator Status <sup>a</sup>	Absolute % Cover
<b>Tree Layer:</b> no species			
<b>Shrub Layer:</b>			
<i>Atriplex confertifolia</i>	shadscale saltbush	NL	2
<i>Ericameria nauseosa</i>	rubber rabbitbrush	NL	2
<i>Salazaria mexicana</i>	Mexican bladdersage	NL	2
<b>Herb Layer:</b>			
<i>Bromus tectorum</i>	cheatgrass	NL	20
<i>Hordeum jubatum</i>	foxtail barley	FAC*	60
Percentage of dominant plant species that are OBL, FACW, or FAC indicator status: <u>50</u> %.			
Dominant plant species are indicated by bold Absolute % Cover values.			
<sup>a</sup> For Region 8 indicator status codes for plants, see Section 3.2.4.			
			Hydrophytic vegetation: <u>No</u>

remain dry in low rainfall years. Knowledge of seasonal availability of water at this site is lacking because of limited prior study. No water quality measurements were taken at this site.

#### 4.2.18.4 Hydric Soils

No field indicators for hydric soils were observed at this site in 1996. Soils appeared dry at the time of the survey and no soil pits were dug. Soils in the area did appear to be dark with a low chroma value.

#### 4.2.18.5 Determination of Jurisdictional Status

This site would probably not qualify as a jurisdictional wetland based on a lack of field indicators for all three required parameters: hydrophytic vegetation, wetland hydrology, and hydric soils. Some seepage may occur in wet years and sustain plant species that are occasionally found in wetlands. However, during dry years few wetland species persist. The lack of archaeological and historical features at the site also suggests that this site did not provide a dependable supply of water.

#### 4.2.18.6 Wildlife Use

Because of limited study, little is known about wildlife use of the area; however, coyote scat and a common raven were observed near the seep area.

located in Fortymile Canyon about 31 m (100 ft) above the canyon floor on a steep (10 percent) west-facing slope (Figure 4-19, Photo 4-51). The area is believed to have been used as a trappers' camp by Native Americans until recent times (Stoffle *et al.*, 1990b). Numerous prehistoric artifacts exist near the spring and on the slope below the spring, including a petroglyph boulder, stone chips, grinding slabs, rock rings, rock coyote trap, and rock weights to hold nets and trap wildlife. Rock shelters also exist on the opposite side of the canyon (Stoffle *et al.*, 1990b).

Much Euroamerican activity has been reported in Fortymile Canyon including mining, prospecting, and travel. During the period of 1870 to 1900, there were freight and mail routes (i.e., Emigrant Trail) that passed through Fortymile Canyon, with relay stations at Whiterock Spring, Tippipah Spring, and Fortymile Canyon (Stoffle *et al.*, 1990a). The Twin Spring site may have been the location of the relay station, although direct evidence for this is lacking (Henton and Pippin, 1988). A lead-silver mine was also worked in the Fortymile Canyon area during the 1880s and was rediscovered and worked again in 1905 (Stoffle *et al.*, 1990a).

The most conspicuous human impact at Twin Spring is a man-made cave dug about 18 m (98 ft) into the hillside and located about 30 m (98 ft) north of the existing spring. This cave occurs at the same elevation as the existing spring and could be the original site of the second spring. The tailings from this cave were leveled off and a rock wall or foundation was built on it. A cave-in was noted about 3 m (10 ft) inside the entrance. Historic artifacts found at the cave include round nails, lumber, and a condensed milk can (Henton and Pippin, 1988). The purpose of the cave is unclear, but Henton and Pippin (1988) suggest it was not dug for mining purposes but as an improvement to the second spring.

Water was piped from the existing spring to the bottom of the wash, a distance of about 300 m (984 ft), as shown by the presence of metal pipes and a large cement water tank. The tank measured about 2 m (6 ft) wide by 3 m (10 ft) long and is inscribed with the date "1921" (Henton and Pippin, 1988).

#### **4.2.19.2 Hydrophytic Vegetation**

A wetlands vegetation survey was conducted on December 12, 1996, from an observation point located at the base of a rock ledge where water flows out and forms a small pool (referred to as the wash slope area). One hundred percent of the dominant plants in this area were hydrophytic species indicating that hydrophytic vegetation was present at Twin Spring (Table 4-22). *T. domingensis* was the only dominant wetland species growing in a small, inundated area within the sample area. Other wetland species which accounted for




Figure 4-19 Location of Twin Spring



Photo 4-52 Wetland vegetation at Twin Spring looking north on January 8, 1997 (WS347-22.TIF)

Table 4-22 Twin Spring wetland vegetation as surveyed on December 12, 1996

<b>Habitat: Wash Slope Area</b>			
<b>Species</b>	<b>Common Name</b>	<b>Indicator Status<sup>a</sup></b>	<b>Absolute % Cover</b>
<b>Tree Layer:</b> no species			
<b>Shrub Layer:</b> no species			
<b>Herb Layer:</b>			
<i>Polypogon monspeliensis</i>	annual rabbitsfoot grass	FACW+	Tr
<i>Rumex salicifolius</i>	willow dock	FACW*	5
<i>Typha domingensis</i>	southern cattail	OBL	90
Percentage of dominant plant species that are OBL, FACW, or FAC indicator status: <u>100</u> %.			
Dominant plant species are indicated by bold values for Absolute % Cover. Tr = trace, <1% absolute cover.			
<sup>a</sup> For Region 8 indicator status codes for plants, see Section 3.2.4.			
Hydrophytic vegetation: <u>Yes</u>			

less than 6 percent of the absolute cover at the site were willow dock and annual rabbitsfoot grass. The wetland area was limited to about 27 m<sup>2</sup> (291 ft<sup>2</sup>) (Table 5-1, Section 5.0). Plants growing in a transitional area on the edge of the inundated area included wormwood, Louisiana sagewort, and skunkbush sumac. The other wet area at Twin Spring was a man-made cave (see Section 4.2.19.3 below) which did not contain hydrophytic vegetation and represented an atypical situation because of prior disturbance. Vegetation in the upland area surrounding the spring included fourwing saltbush, cheatgrass, green rabbitbrush, Virgin River brittlebush (*Encelia virginensis*), Nevada jointfir, mormon tea, eastern Mojave buckwheat, spiny hopsage (*Grayia spinosa*), and Mexican bladdergrass.

#### 4.2.19.3 Wetland Hydrology and Water Quality

This site has been referred to as "Twin Springs" (Stoffle *et al.*, 1990b), which suggests that there were two spring sources at the site. However, only one spring with surface inundation is apparent at the site. Currently, water flows out from the base of a rock ledge and forms a small pool which measures about 2 m<sup>2</sup> (22 ft<sup>2</sup>) (Photo 4-52). This area is referred to as the wash slope area. The maximum depth of the pool within the wash slope area was about 10 cm (4 in) in December 1996. Minimal surface flow occurred down the slope from the pool, but this was not measured. Flow was visible below the wetland area for about 4 m (13 ft) in a steep rocky wash. Water quality measurements were taken in January 1997, and the data are presented in Table 5-2 (Section 5.0).

The man-made cave which occurs at the same elevation as the existing spring could be the original site of the second spring. The cave-in here could have stopped water flow from

wash about 30 m (98 ft) downstream from the observation point. Unidentified mosses were common in the wash channel. Sixty percent of the dominant species at the observation point were wetland species indicating the presence of hydrophytic vegetation at the site (Table 4-23). The total area of the wetland as defined by hydrophytic vegetation within the wash channel is approximately 250 m<sup>2</sup> (2,690 ft<sup>2</sup>) (Table 5-1, Section 5.0). Plant species in the adjacent upland included blackbrush, Cooper's heathgoldenrod, rubber rabbitbrush, basin wildrye, and Mexican bladdersage.




Figure 4-20 Location of Wahmonie Seep 1



Photo 4-54 Wahmonie Seep 2 looking south on June 20, 1996 (WS339-12.TIF)

(Section 5.0).

#### **4.2.20.4 *Hydric Soils***

Field indicators at Wahmonie Seep 1 consisted of saturated soils for what appeared to be more than seven days during the growing season, indicating the presence of hydric soil. A soil pit was dug to determine the presence of other field indicators of hydric soils. Soils lacked evidence of mottling and dark colors (low chroma values).

#### **4.2.21.3 Wetland Hydrology and Water Quality**

Field indicators of wetland hydrology were observed in the wash channel and consisted of surface water and saturated soils. On June 6, 1996, surface water less than 5 cm ( 2 in) deep was observed in isolated small pools less than 0.25 m<sup>2</sup> (2.7 ft<sup>2</sup>) in the wash bottom, but surface water was not observed on June 20, 1996. Based on the presence of wetland species, surface water, and saturated soils as late in the growing season as June 6, it was concluded that indicators of wetland hydrology were present. No water quality measurements were taken at this site.




Figure 4-21 Location of Wahmonie Seep 2



Photo 4-56 Wahmonie Seep 3 looking north on June 20, 1996 (WS339-13.TIF)

Table 4-24 Wahmonie Seep 2 wetland vegetation as surveyed on June 20, 1996

<b>Habitat: Wash Channel</b>			
<b>Species</b>	<b>Common Name</b>	<b>Indicator Status<sup>a</sup></b>	<b>Absolute % Cover</b>
<b>Tree Layer:</b> no species			
<b>Shrub Layer:</b>			
<i>Baccharis emoryi</i>	Emory's baccharis	FACW	<b>85</b>
<b>Herb Layer:</b>			
<i>Artemisia ludoviciana</i>	Louisiana sagewort	FACU	<b>2</b>
unidentified moss		UNKN	<b>1</b>
Percentage of dominant plant species that are OBL, FACW, or FAC indicator status: <u>100</u> %.			
Dominant plant species are indicated by bold Absolute % Cover values.			
<sup>a</sup> For Region 8 indicator status codes for plants, see Section 3.2.4.			
			Hydrophytic vegetation: <u>Yes</u>

#### 4.2.21.4 Hydric Soils

Biologists dug a soil pit and found saturated soils. No other hydric soil field indicators were observed. The soils were shallow, rocky, exhibited no mottling, and had little organic matter. Soils in the bottom of the wash channel at the seep appeared to have been saturated for more than seven days during the growing season, indicating the presence of hydric soils. The area of soil saturation appeared to correspond to the area dominated by Emory's baccharis.

#### 4.2.21.5 Determination of Jurisdictional Status

A portion of the wash channel at Wahmonie Seep 2 would probably be considered a jurisdictional wetland because it has field indicators for all three required parameters: hydrophytic vegetation, wetland hydrology, and hydric soils.

#### 4.2.21.6 Wildlife Use

Little is known about wildlife use of this seep. A bobcat skull was found near the seep suggesting use by this species. Desert cottontails, mule deer scat, and Gambel's quail were observed in the area on June 6, 1996.

saturated (Table 5-1, Section 5.0). Soils at this seep were shallow, rocky, and poorly developed.




Figure 4-22 Location of Wahmonie Seep 3

**Habitat: Wash Channel**

Species	Common Name	Indicator Status <sup>a</sup>	Absolute % Cover
<b>Tree Layer:</b> no species			
<b>Shrub Layer:</b> <i>Baccharis emoryi</i>			
	Emory's baccharis	FACW	60
<b>Herb Layer:</b> <i>Artemisia ludoviciana</i>			
	Louisiana sagewort	FACU	10
	<i>Bromus rubens</i>	UPL	30

Percentage of dominant plant species that are OBL, FACW, or FAC indicator status: 33 %.  
Absolute % cover due to all hydrophytic species present: 60 %. Dominant plant species are indicated by bold Absolute % Cover values.

<sup>a</sup>For Region 8 indicator status codes for plants, see Section 3.2.4.

Hydrophytic vegetation: Yes\*

\*Because the number of dominant hydrophytic species were equal to or less than the number of dominant nonhydrophytic species, an alternative method for determining the presence of hydrophytic vegetation was used (Williams, 1992; see Section 3.2.4).

#### 4.2.22.5 Determination of Jurisdictional Status

The narrow corridor in the wash channel at Wahmonie Seep 3 would probably be considered a jurisdictional wetland because it had field indicators for all three required parameters: hydrophytic vegetation, wetland hydrology, and hydric soils.

#### 4.2.22.6 Wildlife Use

Mule deer scat were located near the seep indicating use by this species. Other species use this area, but no observations were made during the limited sampling period.

#### 4.2.23 Whiterock Spring

##### 4.2.23.1 Site Description and Historical Use

Whiterock Spring is located in a wash at the northern end of Yucca Flat about 2.3 km (1.4 mi) east of the base of Rainier Mesa (Figure 4-23). This site was used as a winter camp by Native Americans from the Belted Range during the late 1800s to early 1900s (Stoffle *et al.*, 1990a). A stone cabin and corral occur near the site which were used during the 1920s (Worman, 1969). Sixty mining claims were recorded from the Whiterock Spring area in 1928 (Stoffle *et al.*, 1990a), suggesting that mining in the area occurred then. Ranching may have occurred in the area in the 1930s. Early ranchers or




Figure 4-23 Location and sketch of Whiterock Spring

Downstream in the lower stretch of the wash, several dry ephemeral pools exist that contained field indicators of wetland hydrology such as dried algae on rocks. Water quality measurements were taken in the flow box in June and September 1996 and in the west cave pool in September 1996. These water quality data are presented in Table 5-2 (Section 5.0).




Photo 4-58 Whiterock Spring without grazing looking north on June 18, 1996 (WS341-31.TIF)




Photo 4-60 Surface pool at Whiterock Spring looking north on December 7, 1990 (WS213-17.TIF)

1. If ROGER is indicator status codes for plants, see Section J.2.4.

Hydrophytic vegetation: Yes

---

Fortynine Canyon (Figure 4-24, Photo 4-61). There appears to be little evidence of human disturbance at the spring except for numerous Native American rock shelters in the area. These rock shelters are located about 23 m (75 ft) above the canyon floor (Stoffle *et al.*, 1990b). One rock shelter contained grinding slabs and a rock wall. It is believed that this area was used temporarily by Native Americans traveling through the area. The wash north of the site is believed to be the likely route used by Native Americans for collecting pinyon nuts on Shoshone Mountain (Stoffle *et al.*, 1990b).

#### **4.2.24.2 Hydrophytic Vegetation**

A wetland vegetation survey was conducted at the Yellow Rock Springs site on December 19, 1996. From the observation point in the rocky wash, no wetland plant species were observed, indicating that hydrophytic vegetation was absent from this site (Table 4-28). Skunkbush sumac was the most dominant plant species and, although it is not considered a hydrophytic species, it appears to be restricted to moist soil habitats on the NTS. Skunkbush sumac and basin wildrye have not been observed in upland habitat around any of the NTS springs visited, and therefore these plants may serve as indicators




Figure 4-24 Location of Yellow Rock Springs




Photo 4-62 Drainage channel at Yellow Rock Spring looking southeast on January 29, 1997  
(WS348-11.TIF)

survey, and water quality measurements were not taken. The surface area of inundation was estimated to be about 30 m<sup>2</sup> (323 ft<sup>2</sup>) (Table 5-1, Section 5.0).

#### **4.2.24.4 *Hydric Soils***

Field indicators for hydric soils were present and appeared to be confined to isolated pockets of soil within bedrock fissures located in the three drainage channels. The floor of all three channels appeared to have been saturated for more than seven days during the plant growing season, indicating the presence of hydric soils. These drainage channels were predominantly rocky with little accumulation of soil fines.

#### **4.2.24.5 *Determination of Jurisdictional Status***

Yellow Rock Springs would probably not be considered a jurisdictional wetland because it lacked field indicators for hydrophytic vegetation.

ephemeral pond defines the wetland plant community dominated by hydrophytic vegetation. Aerial photographs taken in 1964 did not reveal the same pattern of trees around the pond as are seen in aerial photographs taken in 1994, suggesting that saltcedar has colonized the site since 1964. Photo interpretation of the 1964 photos suggest that there may have been about six large shrubs around the pond at that time, and it is unknown if these shrubs were saltcedar or other plant species. Only one other species was observed around the pond edge, southern cattail, which was limited to one small clump of about 1 m<sup>2</sup> (11 ft<sup>2</sup>) in the north end of the pond. There was no evidence of other herbaceous vegetation within the pond during the site survey in January 1997. The hydrophytic vegetation surrounding Yucca Playa Pond covered an estimated area of 3,400 m<sup>2</sup> (36,597 ft<sup>2</sup>) (Table 5-1, Section 5.0).




Figure 4-25 Location and aerial photo of Yucca Playa Pond



Photo 4-64 Drainage channel entering Yucca Playa Pond looking northwest on January 7, 1997  
(WS346-06.TIF)

Table 4-29 Yucca Playa Pond wetland vegetation as surveyed on January 7, 1997

<b>Habitat: Playa Pond</b>			
Species	Common Name	Indicator Status <sup>a</sup>	Absolute % Cover
<b>Tree Layer:</b>			
<i>Tamarix ramosissima</i>	saltcedar	FACW	20
<b>Shrub Layer:</b>			
no species			
<b>Herb Layer:</b>			
<i>Typha domingensis</i>	southern cattail	OBL	Tr
Percentage of dominant plant species that are OBL, FACW, or FAC indicator status: <u>100</u> %.			
Dominant plant species are indicated by bold Absolute % Cover values. Tr = trace, <1% absolute cover.			
<sup>a</sup> For Region 8 indicator status codes for plants see Section 3.2.4.			
			Hydrophytic vegetation: <u>Yes</u>

Upland vegetation surrounding Yucca Playa Pond was quite diverse and somewhat weedy. Key species of plants included fourwing saltbush, cheatgrass, bottlebrush squirreltail, halogeton (*Halogeton glomerata*), prickly Russian thistle (*Salsola paulsenii*), tall tumbled mustard (*Sisymbrium altissimum*), Joshua tree, and several herbs and grasses that were not identified because of a lack of distinguishing taxonomic characteristics at the time of the site survey in January of 1997.

#### 4.2.25.3 Wetland Hydrology and Water Quality

Field indicators of wetland hydrology were present at Yucca Playa Pond. Field indicators included surface water at the pond (also observed in previous years) and a well-eroded shoreline, apparently from wave action, which was barren of vegetation. Several drainage channels direct water off the playa into the pond (Photo 4-64). The surface area of the pond measured approximately 22,930 m<sup>2</sup> (246,840 ft<sup>2</sup>) or 2.3 ha (5.7 ac) (Table 5-1, Section 5.0). Depth of water in January 1997 appeared to be about 150 cm (59 in), although the exact depth could not be determined because of ice. No springs or seeps are known to contribute water to the pond, and water supply appears to be primarily from surface runoff from precipitation events during winter months. It is likely that water persists within the ephemeral pond during the spring but dries during the summer. Water quality measurements were taken on January 7, 1996, and data are presented in Table 5-2 (Section 5.0).

#### 4.2.25.4 Hydric Soils

Field indicators of hydric soils consisted of saturated soil within the pond. The soil appeared to remain saturated for more than seven days during the plant growing season

or decrease the wetness of the area. Indicators of hydric soils are usually poorly represented or absent, while indicators of hydrophytic vegetation and wetland hydrology are more apparent. As mentioned in Section 3.1, some man-induced wetlands are not subject to Section 404 of the CWA. For example, if hydrophytic vegetation is maintained only by wetland hydrology that would no longer exist if an activity (e.g., irrigation) were to be terminated, the area is not considered a jurisdictional wetland. Two areas on the NTS known to support man-induced wetlands include Frenchman Lake and Yucca Flat.

depth of from 1 m (3 ft) to 10 m (33 ft). The excavations intercept and collect surface runoff after storms and periodically fill with water. One of the three field indicators needed for a site to be considered a jurisdictional wetland is the presence of hydrophytic vegetation, which around these excavations consists almost exclusively of saltcedar trees. The presence of the other two required field indicators, wetland hydrology and hydric soils, is questionable and will require further study to determine if ponded water and saturated soils persist long enough into the growing season to be considered wetland hydrology and hydric soils. The presence of hydric soils at these sites is questionable because saturation within the upper 46 cm (18 in) of the soil does not appear to occur. At the majority of the 24 known excavations, the soil supporting the roots of these trees does not appear to be saturated near the soil surface, and it appears that the only time the upper 46 cm (18 in) of soil would be saturated is immediately following a precipitation event. The saturated soil zone appears to be located 1 to 2 m (3 to 6 ft) deep and, therefore, would not meet the criteria needed to be considered hydric soils. If these excavations do not meet the criteria for jurisdictional wetlands they would, however, still be considered waters of the United States by virtue of the fact that they are located on Frenchman Lake (see Section 4.2.27).

Within Yucca Flat, numerous subsidence craters created by historical belowground nuclear tests retain surface water and support hydrophytic vegetation. These craters are another category of man-induced wetlands. These circular depressions (typically about 150 m [500 ft] in diameter and 20 m [70 ft] in depth) also intercept and collect surface runoff after storms, and many of them support hydrophytic vegetation (again, mostly saltcedar trees).

Vegetation within the craters of Yucca Flat and the excavations on Frenchman Lake may be mapped during 1997 as part of the continuing vegetation mapping of the NTS. Further assessments will be made of these areas only when a proposed NTS project may affect them. At that time, the presence of field indicators positive for jurisdictional wetlands would be determined.

#### **4.2.27 Waters of the United States**

There are other natural bodies of water on the NTS that are unvegetated but which attract wildlife and are therefore considered important biological resources. These water bodies include Yucca and Frenchman lakes which periodically flood during heavy precipitation and are known to offer seasonal habitat to migratory waterfowl. Although they are not wetlands by definition, they qualify as waters of the United States. Waters of the United States is a broad category of waters under the jurisdiction of the USACE as authorized under the CWA. They include “. . . waters such as intrastate lakes, rivers, streams (including intermittent streams), mudflats, sandflats, wetlands, sloughs, prairie potholes, wet meadows, playa lakes or natural ponds . . . .”

(msl) (3,925 ft msl) with an area of approximately 2,016 ha (4,982 ac) (Raytheon Services Nevada, 1994). This water level is located approximately at the edge of the unvegetated playa. At Frenchman Lake, the 100-year, 6-hour model elevation is estimated at 939 m msl (3,080 ft msl), and no estimate of area has been made (Julianne Miller, Hydrologist [BN], personal communication, March 3, 1997). This water level is also located approximately at the edge of the unvegetated playa.

## 5.0 DISCUSSION AND SUMMARY OF NTS WETLANDS

### 5.1 Jurisdictional Wetlands Determination

Sixteen of the 25 NTS study sites surveyed in 1996 and 1997 met the three required criteria (hydrophytic vegetation, wetland hydrology, and hydric soils) to be considered jurisdictional wetlands (Figure 5-1, Table 5-1). Copies of wetland field data sheets used to document site conditions and delineate jurisdictional wetland boundaries at the sites are located in Appendix B. Nine sites lacked one or more of the three field indicators needed to be considered jurisdictional wetlands.

All 16 sites which may be considered jurisdictional wetlands had field indicators of hydrophytic vegetation (Table 5-1), suggesting that the sizes of the jurisdictional wetlands increase slightly with increased seepage and water flow. During years of abundant seepage, wetland vegetation increases in area of coverage, and during years of reduced seepage, the total wetland area is reduced. Four sites have cave pools which contain water throughout most of the year but lack vegetation (Cane, Tippipah, Tub, and Whiterock springs), apparently because of reduced light reaching the caves. These pools that lacked hydrophytic vegetation did not meet the criteria to be considered jurisdictional wetlands but may be considered waters of the United States which still fall under the jurisdiction of the USACE. Yucca Playa Pond is an ephemeral pond, the margins of which support wetland vegetation. However, the pond periodically inundates a large area of about 16,246 m<sup>2</sup> (174,807 ft<sup>2</sup>) which appears to support no vegetation. Like the unvegetated cave pools, this area may also be considered waters of the United States.

Twenty-three of the 25 study sites had field indicators of wetland hydrology (Table 5-1). No such field indicators were found at Rainier Spring and Tupapa Seep when these sites were visited (Table 5-1). The source of water at most of the study sites is groundwater discharge from seeps and springs. Surface runoff from precipitation was found at Yucca Playa Pond and at four sites with natural rock catchment basins (tanks). The length of time soils are saturated at these sites which retain surface runoff varies depending on the time of year when precipitation events occur. Winter storms provide water that stays longer in the tanks and pond than water from summer storms. Two sites have tanks (Rock Valley Tank and Yellow Rock Springs), but they also have some water from seepage. Little is known about the persistence of water in all of the known NTS tanks through the year, but it is possible that water in these natural rock depressions, as well as in Yucca Playa Pond, would be considered waters of the United States by the USACE.

Twenty sites were observed to have field indicators for hydric soils (Table 5-1). Field indicators were limited at these sites and were often inferred from site hydrology and past observations of surface water or saturated soils. The five sites lacking field indicators for hydric soils either had no water or no saturated soils at the time the survey was conducted (Pavits Spring, Rainier Spring, Rock Valley Tank, and Tupapa Seep) or were located in bedrock and soils were absent (Tongue Wash Tank). The soils at the remaining 20 sites often lacked field indicators such as mottling and low chroma values (i.e., dark colored soils due to high organic matter content) that are common in wetter climates such as the Great Basin. Desert wetland soils on the NTS are often subject to severe erosion during




Figure 5-1 Water sources identified as jurisdictional wetlands at the NTS

Hippair Spring	500	190	2.7	58	yes	yes	yes	yes
Tongue Wash Tank	0	4.5	0	25	no	yes	no	no
Topopah Spring	200	8	0.12	25	yes	yes	yes	yes
Tub Spring	0	0.1	0.1	5	no	yes	yes	no
Tupapa Seep	0	0	0	0	no	no	no	no
Twin Spring	27	2	NM	10	yes	yes	yes	yes
Wahmonie Seep 1	250	5	NM	8	yes	yes	yes	yes
Wahmonie Seep 2	150	0.25	0	0	yes	yes	yes	yes
Wahmonie Seep 3	180	0	0	0	yes	yes	yes	yes
Whiterock Spring	1,800	0.1	1.9	3	yes	yes	yes	yes
Yellow rock Spring	0	30	NM	40	yes	yes	yes	yes
Yucca Playa Pond	3,400 <sup>f</sup>	22,930	0	150	yes	yes	yes	yes
<b>Total Area (m<sup>2</sup>)</b>	<b>7,193</b>	<b>23,316</b>						

<sup>a</sup>Total surface area over which wetland plans were located. <sup>b</sup>Maximum inundated area recorded at the time of the survey. <sup>c</sup>Maximum flow rate recorded during the year 1996. NM = maximum depth of natural surface water pools. <sup>d</sup>Whether site qualifies as a jurisdictional wetland based on presence of wetland indicators. <sup>e</sup>Area includes that defined by tree canopy cover and an equal area of extended roots.

however, which current flow rates are compared with those from the 1960s. There is one historical record which suggests that flow rates have decreased markedly over the last 25 years at two of the largest NTS springs. In November 1960, Moore (1961) reported discharge rates at Cane Spring of 7.6 to 11.4 *l*/min (2 to 3 gal/min) and at Whiterock Spring of 3.8 to 7.6 *l*/min (1 to 2 gal/min). About 25 years later (1981-1988), measurements ranged from 1.1 to 6.3 *l*/min (0.29 to 1.66 gal/min) for Cane Springs and 0.5 to 4.6 *l*/min (0.37 to 1.22 gal/min) for Whiterock Spring (Ingraham *et al.*, 1991; Lyles *et al.*, 1990). Other evidence which suggests declining flow rates at Cane Spring is the area of inundation which has decreased dramatically from 1963 to the present (see *Photos 4-5 and 4-6*). This reduction probably occurred in the late 1960s or early 1970s when a reduction in flow from NTS springs was noted by Giles (1976).

and surface hydrology. For example, natural activities around springs have caused vegetation changes such as plant introductions, selected grazing of some species, and physical disturbance.

Twenty-three of the 25 study sites discussed in this report demonstrate field indicators for wetland hydrology (Table 5-1). The presence of standing water (inundated soils), however, was limited to 21 of the 25 sites.

### **5.2.2 Chemical Features**

Biologically important water quality parameters (water temperature, dissolved oxygen, pH, total dissolved solids, and electrical conductivity) were measured at 11 wetland sites which had surface water exceeding 5 cm (2 in) in depth (Table 5-2). At three sites (Cane, Tippisah, and Whiterock springs), water quality was sampled at two locations within the

Reitmann	11/22/96	spring pool	12.4	2.7	7.4	287	557
Tippipah	6/18/96	open channel pool	18.6*	1.2	6.8	114	—
Tippipah	9/03/96	open channel pool	18.5	1.0	6.7	135	267
Tippipah	11/15/96	open channel pool	13.7	4.6	7.2	119	243
Tippipah	9/03/96	cave pool	15.3	6.7	7.0	114	227
Tippipah	11/22/96	cave pool	14.3	7.8	7.1	106	212
Topopah	6/20/96	spring pool	14.9*	3.8	7.5	66	—
Topopah	9/09/96	spring pool	20.0	2.7	6.7	69	139
Tub	6/24/96	guzzler can	26.0*	—	7.6	147	—
Tub	9/10/96	guzzler can	26.5	6.0	7.5	146	294
Twin	1/08/97	spring pool	16.8	1.0	7.0	137	271
Wahmonie Seep 1	6/20/96	wash pool	17.8*	1.8	7.5*	259	—
Whiterock	6/18/96	flow box	16.8	8.1*	7.0	124	—
Whiterock	9/03/96	flow box	18.7	6.6	7.2	139	277
Whiterock	9/03/96	west cave pool	15.6	5.8	7.4	142	276
Yucca Playa	1/07/97	pond	1.7	13.6	8.1	162	328

\*Values represent single readings. All other values are an average of three readings. "—" indicates no data collected.

from 7.09 to 7.65. In addition, pH values from selected springs measured by Moore (1961) are similar in most cases to measurements taken in 1996, although readings from Cane and Tippipah springs were more alkaline. These values differ from those obtained by Taylor and Giles (1979) who reported that seven of eight NTS springs monitored had slightly acidic water.

Total dissolved solids and electrical conductivity measured at NTS springs in 1996 (Table 5-2) appear within similar ranges of previously reported values (Moore, 1961; Lyles *et al.*, 1990). Total dissolved solids were fairly constant across season and were generally low (66 to 379 parts per million [ppm]) when compared to irrigation water standards 600 ppm (U.S. Salinity Laboratory, 1954). Topopah Spring had the lowest values, while Reitmann Seep had the highest values. Data collected by others on major cations, anions, and mineral and trace elements in NTS spring water suggest that the water chemistry of NTS springs has not changed greatly over the past 35 years (Moore, 1961; Taylor and Giles, 1979; Romney and Greger, 1994; Lyles *et al.*, 1990; Stetzenbach, 1994).

up approximately 74 percent of the total number of species. Grasses, rushes, and sedges make up only 12 percent, while trees/shrubs make up 14 percent (Beatley, 1976).

The total number of plant species recorded at each wetland site during 1996 varied between 0 and 25. Cane, Topopah, Tippipah, and Whiterock springs had the most species recorded (25, 24, 23, and 22 respectively) (Table 5-3). Tongue Wash Tank had no plant species recorded in the observation area. Three sites, Gold Meadows Spring, Rainier Spring, and Rock Valley Tank had only one species. Low species numbers at sites are due primarily to the rocky nature and lack of soil moisture at these sites. In general, the smaller, ephemeral wetlands had fewer species, while the larger more permanent wetlands had more species (Tables 5-1 and 5-3). Thirty-six species of plants occurred at only a single wetland site, while others (e.g., basin wildrye, Louisiana sagewort, Baltic rush, annual rabbitsfoot grass, and seep monkeyflower) had much wider distributions (Table 5-3).

Overall, there are 24 species of plants at NTS wetlands that had 10 percent or greater absolute cover (classified as dominants) (Table 5-3). Half of these dominants (12 species) are grasses, rushes, or sedges. Forbs, which include more species than the other plant groups, have only five species that are dominants, while trees and shrubs comprised seven dominant species. These dominant species are very important in determining if hydrophytic vegetation is present at a site. Seventeen of the 24 dominant species are listed as obligate or facultative wetland species. They typify wetland areas because they generally do not occur outside very wet habitats. Three dominant species (basin wildrye, Sandberg bluegrass, and Louisiana sagewort) found in NTS wetlands are listed as facultative upland species. Species in this category can tolerate moist conditions but are more typically found in drier upland habitats. On the NTS, however, these three species are found primarily in spring areas, washes, and other areas where moisture is abundant and should probably be listed as facultative wetland species for southern Nevada. The brome grasses, foxtail brome and cheatgrass, were dominants at one or more study sites. They are ubiquitous on disturbed sites. Foxtail brome is listed as an upland species, cheatgrass, is not listed on the Region 8 species list, and they and are not indicators of mesic habitats on the NTS. There are two other dominants which were recorded at several study sites:





Wetlands are important habitats for many species of animals. A total of 138 species of animals have been documented at NTS wetland sites (Appendix D). These species include various classes of animals including mammals, birds, reptiles, and terrestrial insects. The largest group of vertebrates using wetlands are birds (Table 5-4). Throughout the arid west, an extraordinary diversity of bird species depend on wetland habitats (Carothers *et al.*, 1974; Knopf *et al.*, 1988a,b; Dobkin, 1994). Among the 134 species of migratory landbirds that breed regularly in the Great Basin, more than half are associated primarily with riparian habitats (Dobkin, 1996). Destruction or degradation of wetland habitats is widely viewed as the most important factor in the decline of landbird populations in western North America (Bock *et al.*, 1993; Desante and George, 1994; Ohmart, 1994). For this reason, land management agencies are aware of the importance of protecting and restoring riparian habitats for birds and other wildlife (Warner and Hendrix, 1984).

### 5.3.3 Wildlife Use

Wetlands are important habitats for many species of animals. A total of 138 species of animals have been documented at NTS wetland sites (Appendix D). These species include various classes of animals including mammals, birds, reptiles, and terrestrial insects. The largest group of vertebrates using wetlands are birds (Table 5-4). Throughout the arid west, an extraordinary diversity of bird species depend on wetland habitats (Carothers *et al.*, 1974; Knopf *et al.*, 1988a,b; Dobkin, 1994). Among the 134 species of migratory landbirds that breed regularly in the Great Basin, more than half are associated primarily with riparian habitats (Dobkin, 1996). Destruction or degradation of wetland habitats is widely viewed as the most important factor in the decline of landbird populations in western North America (Bock *et al.*, 1993; Desante and George, 1994; Ohmart, 1994). For this reason, land management agencies are aware of the importance of protecting and restoring riparian habitats for birds and other wildlife (Warner and Hendrix, 1984).

**Table 5-4** Number of vertebrate species recorded at each wetland study site on the NTS

148

Groups	Ammonia Tanks	Cane Spring	Captain Jack Spring	Cottonwood Spring	Coyote Spring	Fortymile Canyon Tanks	Gold Meadows Spring	John's Spring	Oak Spring	Pavits Spring	Rainier Spring	Reitmann Seep	Rock Valley Tank	Tippipah Spring	Tongue Wash Tank	Topopah Spring	Tub Spring	Tupapa Seep	Twin Spring	Wahmonie Seep 1	Wahmonie Seep 2	Wahmonie Seep 3	Whiterock Spring	Yellow Rock Springs	Yucca Playa Pond	Total (unique species)
Mammals	3	11	5	2	2	2	9	2	2	2	0	3	3	6	1	5	5	1	4	2	3	1	4	3	3	15
Upland game birds	1	3	3	0	0	0	3	0	1	1	0	2	0	3	1	3	3	0	1	1	0	0	2	0	1	3
Migratory waterfowl	0	6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	6
Raptors	0	6	5	0	0	1	6	0	0	0	0	1	0	3	0	4	0	0	0	0	0	0	3	0	1	11
Passerine birds	0	69	16	0	0	0	25	2	2	0	0	3	0	16	2	7	6	1	0	1	0	0	13	0	0	80
T & E species	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1
<b>Totals</b>	<b>4</b>	<b>95</b>	<b>29</b>	<b>2</b>	<b>2</b>	<b>3</b>	<b>43</b>	<b>4</b>	<b>5</b>	<b>3</b>	<b>0</b>	<b>9</b>	<b>4</b>	<b>28</b>	<b>4</b>	<b>19</b>	<b>14</b>	<b>2</b>	<b>5</b>	<b>4</b>	<b>3</b>	<b>1</b>	<b>22</b>	<b>3</b>	<b>7</b>	<b>116</b>

~~Yucca Playa Pond is known to support three species of shrimp (fairy, tadpole, and clam shrimp) (Starkweather, 1995). The one species of hydrobiid snail (*Pyrgulopsis* sp.) is now~~  
pond, which existed at the site when discharge rates were higher, probably attracted more bird species, particularly migratory waterfowl and passerine birds. Gold Meadows Spring, Captain Jack Spring, and Tippipah Spring also support fair numbers of species (44, 29, 26, respectively). There are several study sites that show very low numbers of species (Cottonwood Spring, Coyote Spring, Fortymile Canyon Tanks, Rainier Spring, Tupapa Seep, and Wahmonie Seep 3). All of these sites (except Cottonwood Spring) have only ephemeral water that is unavailable for wildlife use much of the year. These low numbers of species recorded may be a result of very limited study (particularly for Cottonwood Spring) and may underestimate their importance to wildlife.

Many aquatic invertebrate organisms occur in NTS springs despite their small surface area. Observed groups of aquatic organisms in springs include oligochaetes (segmented worms), nematodes (roundworms), copepods, ostracods (seed shrimps), cladocerans (water fleas), chironomids (midge larvae), and hydrobiid gastropods (springsnails); and Yucca Playa Pond is known to support three species of shrimp (fairy, tadpole, and clam shrimp) (Starkweather, 1995). The one species of hydrobiid snail (*Pyrgulopsis* sp.) is now

restricted to a small area of the spring cave pool at Cane Spring. This organism also occurs in four springs located in the northern region of the Spring Mountains some 64 km (40 mi) south of Cane Spring. Relatively little is known about aquatic invertebrates from riparian habitats in the Great Basin (Hersher and Pratt, 1990). They are also poorly known on the NTS.

will be considered and may be incorporated into new goals. For example, DOE/NV's goal for wetlands preservation may be linked with the Nevada Division of Wildlife's goal to maintain viable populations of chukar that are known to use selected NTS springs. The management goals and practices described below are currently implemented on the NTS and ensure compliance with wetlands legislation and DOE policy to manage the NTS under the guiding principles of ecosystem management.

### **6.1 Protection as Important Biological Resources**

Many natural water sources on the NTS provide islands of unique habitat within the Mojave and Great Basin Desert communities of south-central Nevada. Wetlands, by definition, support plants that do not normally grow in unsaturated desert soils, and therefore increase the biodiversity of plant communities on the NTS. Similarly, some NTS wetlands provide rare environments within the region for isolated populations of aquatic organisms. They also attract migratory waterfowl, passerine birds, and game species of wildlife, as documented in this report. For these reasons, and apart from their jurisdictional status under the CWA, the NTS wetlands are regarded as rare habitats which are important biological resources. The management goals for all important biological resources on the NTS include (1) avoiding impacts to the resource whenever possible, and (2) minimizing all unavoidable impacts. Additional goals germane to NTS wetlands specifically include restoring the biological integrity of wetlands if degradation occurs and preserving and enhancing the natural and beneficial values of NTS wetlands. These goals combined meet the intent of CWA, NEPA, Executive Order 11990, and the DOE Land- and Facility-Use Management Policy.

### **6.2 Continued Monitoring**

DOE/NV has conducted periodic surveys of nine NTS natural water sources (Cane, Captain Jack, Gold Meadows, Oak, Tippihah, Topopah, Tub, and Whiterock springs, and Reitmann Seep) since 1989. These surveys have focused on wildlife observations and have served to identify wetland function and values, such as seasonal habitat for migrating water fowl. They are integrated with other ecological monitoring tasks such as surveys to census wild horses and chukar. These surveys will continue, and opportunities will be taken to sample for aquatic organisms and to collect basic water quality data at these sites. Results of continued monitoring of these selected water sources may prove valuable in better understanding the function and value of the NTS wetlands.

### **6.3 Identification and Evaluation of Other NTS Wetlands**

Surveys will continue to be conducted on the NTS to locate other springs, seeps, and playa wetlands. These surveys will be conducted primarily in concert with other ecological monitoring field tasks. The three seeps called Wahmonie Seeps 1, 2, and 3 were discovered in the summer of 1996 during field surveys to map tortoise habitat in the southern one-third of the NTS. Habitat mapping of the northern two-thirds of the NTS is planned for Fiscal Years 1997 and 1998, and it is likely that more wetlands will be found during that effort. To identify all important biological resources on the NTS, biologists will take advantage of all opportunities in the field to visit, describe, sample, and photograph new wetland sites according to the methods described in Section 3.0 of this report. Each new wetland discovered will also be evaluated for its jurisdictional status under the CWA.

### **6.4 Maintenance of Wetlands Geospatial Database**

DOE has developed several tools to manage important biological resources, including wetlands, on the NTS. These tools include the EGIS which contains a wetlands geospatial database containing the point locations of all known NTS wetlands linked to field data on vegetation, hydrology, soils, and wildlife usage. Selected digitized photographs of each wetland site are also linked to the database. This database will be updated annually to incorporate all new data collected at known sites or to add data on newly discovered wetland sites. The wetlands geospatial database will eventually be linked with information about other natural and man-made resources for use during implementation of the NTS comprehensive land- and facility-use management plan. These databases will also be used for the preparation of future project-specific EAs and EISs.

### **6.5 Inclusion in Project Siting and Permitting Procedures**

Siting procedures for proposed NTS projects include an evaluation of impacts on important biological resources, including wetlands. Projects are redesigned or relocated whenever possible to avoid adverse impacts. It is possible that a proposed NTS project may unavoidably impact a jurisdictional wetland and require a permit under Section 404 of the CWA. The types of activities that require a permit would be those that destroy wetland habitat or alter the discharge, flow, or movement of water through a wetland, which may include

- clearing wetlands if it involves excavating, leveling, filling, or using heavy equipment in the wetlands;
- constructing or improving ditches or berms in wetlands, especially if the ditches or berms alter the flow of water through the wetlands;
- destroying vegetation either physically, mechanically, or chemically (e.g., burning, mowing, or herbicides); and
- dewatering seeps or springs by intercepting ground water.

A limited number of activities are exempted from regulation under Section 404. Examples of these activities include

- maintenance of currently serviceable structures such as dikes, dams, bridge abutments, and transportation structures; and
- construction of temporary sediment basins where fill material is not placed into navigable waters.

Much of the site-specific information provided in this report is sufficient to initiate a preapplication consultation with the USACE if an NTS project required a Section 404 permit.

**This Page Intentionally Left Blank**

1975, 1981. Desert wildlife water developments. Questioning use in the Southwest. Wildlife Society Bulletin, Vol 23, No. 4, pp. 663-675.

Carothers, S. W., R. R. Johnson, and S. W. Aitchison, 1974. Population structure and social organization of southwestern riparian birds. *American Zoologist* 14:97-108.

Castetter, R. C., and H. O. Hill, 1979. Additions to the birds of the Nevada Test Site. *Western Birds* 10:221-223.

Clebsch, A., Jr., 1960. Ground water in the Oak Spring formation and hydrologic effects of underground nuclear explosions at the Nevada Test Site. U.S. Geological Survey open-file report TEI-759, p. 29.

Cowardin, L., V. Carter, F. Golet, and E. LaRoe, 1979. Classification of wetlands and deep water habitats of the United States. U.S. Fish and Wildlife Service, Report No. F'S/OBS-79/31, p. 103. U.S. Department of Interior, Washington, DC.

Davis, M., B. Dicey, C. Fontana, K. Giles, R. Grossman, P. Huff, B. Hooda, A. Mullen, M. Sells, and C. Wills, 1996. Nevada Test Site Annual Site Environmental Report - Calendar Year 1995. DOE/NV/11718. S. C. Black, and Y. E. Townsend (eds.) U.S. Department of Energy/Nevada Operations Office, Las Vegas, Nevada.

Desante, D. F., and T. L. George, 1994. Population trends in the landbirds of western North America. *In*: J. R. Jehl, Jr., and N. K. Johnson (eds.), A century of avifaunal change in western North America. *Studies in Avian Biology*, Volume 15, pp. 173-190.

- , 1982. Title 33: Navigation and Navigable Waters, Chapter 1, Regulatory Programs of the Corps of Engineers. Vol 47, No. 138, p. 31810. U.S. Government Printing Office, Washington, DC.
- , 1980. 40CFR Part 230: Section 404(b)(1), Guidelines for Specification of Disposal Sites for Dredged or Fill Material. Vol 45, No. 249, p. 85352-85353. U.S. Government Printing Office, Washington, DC.
- Fernald, A. T., G. S. Corchary, and W. P. Williams, 1968. Surficial Geologic Map of Yucca Flat, Nye and Lincoln Counties, Nevada. U.S. Department of the Interior, U.S. Geological Survey. Miscellaneous Geologic Investigations Map 1-550.
- Frizzell, V. A. Jr., and J. Shulters, 1990. A geologic map of the Nevada Test Site, Southern Nevada. U.S. Geological Survey, U.S. Department of the Interior.
- Giles, K. R., 1976. Springs on the Nevada Test Site and their use by wildlife. NERC-LV-539-26, p. 14. U.S. Environmental Protection Agency, Las Vegas, Nevada, April, 1976.
- Greger, P. D., 1995. Status of horses, deer, and birds on the Nevada Test Site. *In*: Status of the flora and fauna of Nevada Test Site 1994. DOE/NV/11432-195. Compiled by R. B. Hunter.
- Greger, P. D., 1994. Status of large mammals and birds on the Nevada Test Site in 1993. *In*: Status of the flora and fauna on the Nevada Test Site, 1993. DOE/NV/11432-162, pp. 124-149. Compiled by R. B. Hunter.
- Greger, P. D., and E. M. Romney, 1994a. Trends in wildlife utilization of water sources and adjacent habitats at the Nevada Test Site from 1989-1991. *In*: Status of the flora and fauna on the Nevada Test Site, 1989- 1991. DOE/NV/11432-57, pp. 170-235. Compiled by R. B. Hunter.

- Greger, P. D., and E. M. Romney, 1994b. Status of large mammals and birds on the Nevada Test Site, 1992. *In: Status of flora and fauna on the Nevada Test Site, 1992.* DOE/NV/11432-58, pp. 144-175. Compiled by R. B. Hunter.
- Hayward, C. L., M. L. Killpack, and G. L. Richards, 1963. Birds of the Nevada Test Site. *Brigham Young University Science Bulletin, Biological Series, Vol 3. No.1,* p. 27.
- Henton G. H., and L. C. Pippin, 1988. Prehistoric and historic archaeology of Fortymile Canyon, Yucca Wash, and Midway Valley, near Yucca Mountain, Nye County, Southern Nevada. Technical Report 60, p. 163. Desert Research Institute, University of Nevada System.
- Hersher, R., and W. L. Pratt, 1990. A new *Pyrgulopsis* (Gastropoda: Hydrobiidae) from southwestern California, with a model for historical development of the Death Valley hydrographic system. *Proceedings of the Biological Society of Washington.* 103(2):279-299.
- Ingraham, N. L., B. F. Lyles, R. L. Jacobson, and J. W. Hess, 1991. Stable Isotopic Study of Precipitation and Spring Discharge in Southern Nevada. *Journal of Hydrology* 125: 243-258.
- Jorgensen, C. D., and C. L. Hayward, 1965. Mammals of the Nevada Test Site. *Brigham Young University Science Bulletin, Biological Series* 6(3):1-81.
- Knopf, F. L., R. R. Johnson, T. Rich, F. B. Samson, and R. C. Szaro, 1988a. Conservation of riparian ecosystems in the United States. *Wilson Bulletin* 100:272-284.
- Knopf, F. L., J. A. Sedgewick, and R. W. Cannon, 1988b. Guild Structure of a riparian avifauna relative to seasonal cattle grazing. *Journal of Wildlife Management* 52:280-290.
- Lewis G. B., M. A. Glora, and I. Aoki, 1965. Environmental Radioactivity at the Nevada Test Site. NVO-162-18. Radiological Services Department, U.S. Department of Energy/Nevada Operations Office, Las Vegas, Nevada.
- Lyles, B. F., J. Edkins, R. L. Jacobson, and J. W. Hess, 1990. Time-series analysis of ion and isotope geochemistry of selected springs of the Nevada Test Site, Nye County, Nevada. DOE/NV/10384-27, p. 121. Water Resources Institute, Desert Research Institute, University of Nevada System.
- McKeown, F. A., D. L. Healey, and C. H. Miller, 1976. Geologic Map of the Yucca Lake Quadrangle, Nye County, Nevada. Map GQ-1327, 1:24,000. U.S. Geological Survey, U.S. Department of the Interior.
- Moore, J. E., 1961. Records of wells, test holes, and springs in the Nevada Test Site and surrounding area. U.S. Geological Survey open-file report TEI-781, p. 22.

Cooperation with National Technical Committee for Hazardous Waste, Washington, DC.

NRC (National Research Council), 1995. Wetlands: Characteristics and Boundaries, Committee on Characterization of Wetlands. National Academy Press, Washington, DC.

O'Farrell, T. P., and L. A. Emery, 1976. Ecology of the Nevada Test Site: A Narrative Summary and Annotated Bibliography. Report NVO-167, U.S. Energy Research and Development Administration (ERDA).

Ohmart, R. D., 1994. The effects of human induced changes on the avifauna of western riparian habitats. *In*: J. R. Jehl Jr., and N. K. Johnson (eds.), A century of avifaunal change in western North America. Studies in Avian Biology. Vol. 15, pp. 273-285.

Raytheon Services Nevada, 1994. Flood Assessment at the Proposed Area 6 Liquid Waste Treatment System. DOE/Nevada Test Site, Nye County, Nevada. Draft, October 1994.

Reed, P. B., Jr., 1996. National list of plant species that occur in wetlands: Intermountain (Region 8) revised draft. National Ecology Research Center, U.S. Fish and Wildlife Service, St. Petersburg, Florida.

Rhoads, W. A., and M. P. Williams, 1977. Status of endangered and threatened plant species on the Nevada Test Site - A survey. EG&G Report 1183-2356. Santa Barbara, California.

IIIUCM/Guion Program Environmental Report Nevada Test Site and Vicinity. Report No. EMSL-LV-0539-14, p. 45. U.S. Environmental Protection Agency, Las Vegas, Nevada.

Starkweather, P., 1995. Modern and paleolimnology of the endorheic playas of the Nevada Test Site. University of Nevada, Las Vegas. Final report to DOE/NV DE-FC08-93NV11399, p.19 plus figures. Supported under the cooperative agreement with the U.S. Department of Energy.

Stetzenbach, K. J., 1995. Fingerprinting of ground water by ICP-MS. Unpublished progress report DOE/NV/10872-T203. Harry Reid Center for Environmental Studies, University of Nevada, Las Vegas, Nevada.

Stoffle, R.W., J. E. Olmsted, and M. J. Evans, 1990a. Literature review and ethnohistory of Native American occupancy and use of the Yucca Mountain area. DOE/NV-10576-21, p. 197. Institute for Social Research, University of Michigan, Ann Arbor, Michigan. Report to Science Applications International Corporation, Las Vegas, Nevada.

Stoffle, R.W., D. B. Halmo, J. E. Olmsted, and M. J. Evans, 1990b. Native American cultural resource studies at Yucca Mountain Nevada. p. 232. Institute for Social Research. The University of Michigan, Ann Arbor, Michigan. Report to Science Applications International Corporation, Las Vegas, Nevada.

Impact Statement for the Nevada Test Site and Off Site Locations in the State of Nevada. DOE/EIS 0234. Volume 2. U.S. Department of Energy/Nevada Operations Office, Las Vegas, Nevada.

———, 1988. Site Characterization Plan, Yucca Mountain Site, Nevada Research and Development Area, Nevada. DOE/R.W.-0199, Volume 11, Part A.

U.S. Department of Interior. 1994. An Ecosystem Approach to Fish and Wildlife Conservation, U.S. Fish and Wildlife Service, Washington, DC.

U.S. Salinity Laboratory, 1954. Diagnosis and improvement of saline and alkali soils. Agriculture Handbook 60, USDA.

Warner, R. E., and K. M. Hendrix, 1984. California riparian systems: ecology, conservation and productive management. p. 1035. University of California Press, Berkeley, California.

Williams, A. E., 1992. Clarification and Interpretation of the 1987 Manual, USACE Memorandum from Arthur E. Williams, Major General, USA, Directorate of Civil Works, Correspondence CECW-OR.

Winograd, I. J., and W. Thordarson, 1975. Hydrogeologic and Hydrochemical Framework, South-Central Great Basin, Nevada-California, with special reference to the Nevada Test Site. Hydrology of Nuclear test Sites. U.S. Geological Survey Professional Paper 712-C, p. 126. U.S. Department of Interior.

Worman, F. C. V., 1969. Archeological investigations at the U.S. Atomic Energy Commission's Nevada Test Site and Nuclear Rocket Development Station. LA-4125, p. 201. Los Alamos Scientific Laboratory of the University of California, Los Alamos, New Mexico.

**This Page Intentionally Left Blank**

**APPENDIX A**

**LIST OF SCIENTIFIC AND  
COMMON PLANT NAMES**

Appendix A. List of scientific and common plant names\*

SCIENTIFIC NAME	COMMON NAME	SCIENTIFIC NAME	COMMON NAME
<b>Trees/Shrubs</b>		<b>Grasses/Rushes/Sedges (Cont.)</b>	
<i>Ambrosia dumosa</i>	white burrobush	<i>Bromus rubens</i>	foxtail brome
<i>Amelanchier utahensis</i>	Utah serviceberry	<i>Bromus tectorum</i>	cheatgrass
<i>Artemisia tridentata</i>	big sagebrush	<i>Carex</i> sp.	sedge
<i>Atriplex canescens</i>	fourwing saltbush	<i>Carex alma</i>	sturdy sedge
<i>Atriplex confertifolia</i>	shadscale saltbush	<i>Carex praegracilis</i>	clustered field sedge
<i>Baccharis emoryi</i>	Emory's baccharis	<i>Carex alma</i>	sturdy sedge
<i>Cercocarpus intricatus</i>	littleleaf mountain mahogany	<i>Carex praegracilis</i>	clustered field sedge
<i>Chrysothamnus</i> sp.	rabbitbrush	<i>Dactylis glomerata</i>	orchardgrass
<i>Chrysothamnus viscidiflorus</i>	green rabbitbrush	<i>Deschampsia danthonioides</i>	annual hairgrass
<i>Coleogyne ramossissima</i>	blackbrush	<i>Distichlis spicata</i>	inland saltgrass
<i>Encelia virginensis</i>	Virgin River brittlebush	<i>Eleocharis palustris</i>	common spikerush
<i>Ephedra nevadensis</i>	Nevada jointfir	<i>Eleocharis parishii</i>	Parish's spikerush
<i>Ephedra viridis</i>	mormon tea	<i>Elymus elymoides</i>	bottlebrush squirreltail
<i>Ericameria cooperi</i>	Cooper's heathgoldenrod	<i>Elymus</i> sp.	wildrye
<i>Ericameria nauseosa</i>	rubber rabbitbrush	<i>Hordeum jubatum</i>	foxtail barley
<i>Ericameria teretifolia</i>	needle leaf rabbitbrush	<i>Hordeum murinum</i> ssp. <i>glaucum</i>	smooth barley
<i>Eriogonum fasciculatum</i>	eastern Mojave buckwheat	<i>Juncus balticus</i>	baltic rush
<i>Grayia spinosa</i>	spiny hopsage	<i>Juncus longistylis</i>	longstyle rush
<i>Juniperus osteosperma</i>	Utah juniper	<i>Juncus saximontanus</i>	Rocky Mountain rush
<i>Larrea tridentata</i>	creosote+B3bush	<i>Leymus cinereus</i>	basin wildrye
<i>Pinus monophylla</i>	singleleaf pinyon	<i>Leymus triticoides</i>	beardless wildrye
<i>Populus fremontii</i>	Fremont's cottonwood	<i>Phragmites australis</i>	common reed
<i>Prosopis glandulosa</i> var. <i>torreyana</i>	western honey mesquite	<i>Poa secunda</i>	Sandberg bluegrass
<i>Prunus fasciculata</i>	desert almond	<i>Poa</i> sp.	bluegrass
<i>Purshia glandulosa</i>	desert bitterbrush	<i>Polypogon interruptus</i>	ditch polypogon
<i>Purshia stansburiana</i>	Stansbury cliffrose	<i>Polypogon monspeliensis</i>	annual rabbitsfoot grass
<i>Purshia tridentata</i>	antelope bitterbrush	<i>Polypogon viridis</i>	beardless rabbitsfoot grass
<i>Quercus gambelii</i>	Gambel's oak	<i>Potamogeton pectinatus</i>	sago pondweed
<i>Rhus trilobata</i>	skunkbush sumac	<i>Sporobolus airoides</i>	alkali sacaton
<i>Salazaria mexicana</i>	Mexican bladdersage	<i>Typha domingensis</i>	southern cattail
<i>Salix exigua</i>	sandbar willow	<i>Typha latifolia</i>	broadleaf cattail
<i>Salix gooddingii</i>	Goodding's willow		
<i>Symphoricarpos longiflorus</i>	desert snowberry	<b>Forbs</b>	
<i>Tamarix chinensis</i>	fivestamen tamarisk	<i>Amaranthus albus</i>	prostrate pigweed
<i>Tamarix ramosissima</i>	saltcedar	<i>Artemisia dracunculus</i>	wormwood
<i>Yucca brevifolia</i>	Joshua tree	<i>Artemisia ludoviciana</i>	Louisiana sagewort
<i>Yucca schidigera</i>	Mojave yucca	<i>Berula erecta</i>	cutleaf waterparsnip
		<i>Camissonia megalantha</i>	largeflower suncup
<b>Grasses/Rushes/Sedges</b>		<i>Camissonia</i> sp.	suncup
<i>Agrostis exarata</i> var. <i>monolepis</i>	monolepis bentgrass	<i>Castilleja applegatei</i> ssp. <i>martinii</i>	Martin's wavyleaf Indian paintbrush
<i>Bromus diandrus</i>	ripgut brome		

\* Species nomenclature is according to NRCS, 1996a

Appendix A. List of scientific and common plant names\*

SCIENTIFIC NAME	COMMON NAME
<b>Forbs (Cont.)</b>	
<i>Cirsium neomexicanum</i>	New Mexico thistle
<i>Cleome lutea</i>	yellow spiderflower
<i>Conyza canadensis</i>	Canadian horseweed
<i>Encelia</i> sp.	brittlebush
<i>Epilobium ciliatum</i>	hairy willowherb
<i>Epilobium glaberrimum</i>	smooth willowweed
<i>Erigeron divergens</i>	spreading fleabane
<i>Erodium cicutarium</i>	redstem stork's bill
<i>Galium aparine</i>	stickywilly
<i>Halogeton glomerata</i>	halogeton
<i>Heliomeris multiflora</i> var. <i>nevadensis</i>	Nevada goldeneye
<i>Kochia scoparia</i>	common kochia
<i>Lactuca serriola</i>	prickly lettuce
<i>Linum lewisii</i>	prairie flax
<i>Linum</i> sp.	flax
<i>Lotus</i> sp.	trefoil
<i>Melilotus indicus</i>	annual yellow sweetclover
<i>Mimulus guttatus</i>	seep monkeyflower
<i>Oenothera cespitosa</i> ssp. <i>marginata</i>	tufted eveningprimrose
<i>Penstemon rostriflorus</i>	Bridge penstemon
<i>Penstemon</i> sp.	penstemon
<i>Pentagramma triangularis</i>	western goldfern
<i>Potentilla biennis</i>	biennial cinquefoil
<i>Pseudognaphalium stramineum</i>	straw falsecudweed
<i>Rumex crispus</i>	curly dock
<i>Rumex salicifolius</i>	willow dock
<i>Salsola pausensii</i>	prickly Russian thistle
<i>Sisymbrium altissimum</i>	tall tumbledustard
<i>Suaeda moquinii</i>	Mojave seablite
<i>Verbena bracteata</i>	bigbract verbena
<i>Veronica anagallis-aquatica</i>	water speedwell

\*Species nomenclature is according to NRCS, 1996a

**APPENDIX B**

**DATA SHEETS FOR ROUTINE  
JURISDICTIONAL WETLAND DETERMINATION**

# Routine Jurisdictional Wetland Determination

Name: PDG, DJH Wetland Unit: Wash pool  
 Location: Ammonia Tanks UTM Coordinates Easting: 5-62-850 Northing: 41-10-240  
 Date: 1-7-97

## Hydrology

Type: Seep  Spring  Pond  Detention basin  Stream  Natural Tanks   
 Source: Natural  Man-enhanced  Man-made  Ephemeral  Permanent  Temporary   
 Date of construction/Period of flow: On 1-7-97 there was no water flow out of the tanks.  
 Disturbance type (if any) and date: None: A nearby rock shelter indicates human historical use  
 Inundated: Yes  No  Depth of standing water >50cm; Saturated: Yes  No  Depth to saturation 0  
 Other field indicators: \_\_\_\_\_  
 Atypical situation: Yes  No ; Wetland hydrology: Yes  No   
 Basis: Surface water exists at the observation point.

## Vegetation *List 3 dominant species, percent cover in bold, in each vegetation layer (5 if only 1 or 2 layers are present)*

	<i>Species</i>	<i>Indicator Status<sup>a</sup></i>	<i>% Cover</i>
<b>Trees</b>			
1.	<u>none</u>	<u>_____</u>	<u>_____</u>
2.	<u>_____</u>	<u>_____</u>	<u>_____</u>
3.	<u>_____</u>	<u>_____</u>	<u>_____</u>
4.	<u>_____</u>	<u>_____</u>	<u>_____</u>
5.	<u>_____</u>	<u>_____</u>	<u>_____</u>
<b>Shrubs</b>			
1.	<u>none</u>	<u>_____</u>	<u>_____</u>
2.	<u>_____</u>	<u>_____</u>	<u>_____</u>
3.	<u>_____</u>	<u>_____</u>	<u>_____</u>
4.	<u>_____</u>	<u>_____</u>	<u>_____</u>
5.	<u>_____</u>	<u>_____</u>	<u>_____</u>
<b>Herbs</b>			
1.	<u>Artemisia ludoviciana</u>	<u>FACU</u>	<u>30</u>
2.	<u>Encelia sp.</u>	<u>UNKN</u>	<u>2</u>
3.	<u>Juncus balticus</u>	<u>FACW</u>	<u>8</u>
4.	<u>Leymus cinereus</u>	<u>FACU</u>	<u>5</u>
5.	<u>_____</u>	<u>_____</u>	<u>_____</u>
6.	<u>_____</u>	<u>_____</u>	<u>_____</u>
7.	<u>_____</u>	<u>_____</u>	<u>_____</u>
8.	<u>_____</u>	<u>_____</u>	<u>_____</u>
9.	<u>_____</u>	<u>_____</u>	<u>_____</u>
10.	<u>_____</u>	<u>_____</u>	<u>_____</u>

Other field indicators: \_\_\_\_\_  
 Percentage of species that are OBL, FACW, and/or FAC: 0%; Hydrophytic vegetation: Yes  No   
 Basis: Lack of dominance of hydrophytic species at the observation point.

## Hydric Soils

Field indicators: Hydric soils exist at the observation point due to presence of saturated soils for longer than 7 days duration.

Hydric Soils: Yes  No

Jurisdictional Wetland Determination : Wetland  Nonwetland

## Notes:

<sup>a</sup> Wetland indicator status for plants in region 8. FACW = facultative wetland species. FACU = Facultative upland species.  
 UNKN = Unknown wetland status of this species.

# Routine Jurisdictional Wetland Determination

Name: PDG, DJH, JAA Wetland Unit: Drainage channel below cave pool  
 Location: Cane Spring UTM Coordinates Easting: 5-80-775 Northing: 40-72-730  
 Date: 6-19-96

## Hydrology

Type: Seep  Spring  Pond  Detention basin  Stream  Mechanically contained   
 Source: Natural  Man-enhanced  Man-made  Ephemeral  Permanent  Temporary   
 Date of construction/Period of flow: Unknown date of construction. PVC pipe directs water from cave to a flow gauge box.  
 Disturbance type (if any) and date: Channels dug out leading to a dry pond. An old road forms a berm-Cave adit pool dug out.  
 Inundated: Yes  No  Depth of standing water 0; Saturated: Yes  No  Depth to saturation 12"-18"  
 Other field indicators: Hydrobiid snails, *Pyrgulopsis sp* restricted to cave pool living on filamentous algae present near opening of the cave pool  
 Atypical situation: Yes  No ; Wetland hydrology: Yes  No   
 Basis: At the observation point, the outflow channel is dry at the surface, but is saturated at 1ft depth. No surface outflow from the cave pool occurs on this date. Flow rate measured at the cave pool in November 22, 1996 was about 3 liters/min.

**Vegetation** List 3 dominant species, % cover in bold, in each vegetation layer (5 if only 1 or 2 layers are present)

	Species	Indicator Status <sup>a</sup>	% Cover
<b>Trees</b>			
1.	<u>none</u>	<u></u>	<u></u>
2.	<u></u>	<u></u>	<u></u>
3.	<u></u>	<u></u>	<u></u>
4.	<u></u>	<u></u>	<u></u>
<b>Shrubs</b>			
1.	<u><i>Atriplex canescens</i></u>	<u>UPL</u>	<u>15</u>
2.	<u></u>	<u></u>	<u></u>
3.	<u></u>	<u></u>	<u></u>
4.	<u></u>	<u></u>	<u></u>
5.	<u></u>	<u></u>	<u></u>
<b>Herbs</b>			
1.	<u><i>Leymus cinereus</i></u>	<u>FACU</u>	<u>40</u>
2.	<u><i>Rumex salicifolius</i></u>	<u>FACW*</u>	<u>15</u>
3.	<u></u>	<u></u>	<u></u>
4.	<u></u>	<u></u>	<u></u>
5.	<u></u>	<u></u>	<u></u>
6.	<u></u>	<u></u>	<u></u>
7.	<u></u>	<u></u>	<u></u>
8.	<u></u>	<u></u>	<u></u>
9.	<u></u>	<u></u>	<u></u>

Percentage of species that are OBL, FACW, and/or FAC: 33%; Hydrophytic vegetation: Yes  No   
 Basis: Observation point was dry during the survey - other areas nearby have hydrophytic vegetation.

## Hydric Soils

Field indicators: Hydric soils-undisturbed: Soils have mottling at depths of about 5-6 ft. under the surface, shown on a cut away bank. Overall depths of soils approximately 6-8 ft.

Hydric Soils: Yes  No

Jurisdictional Wetland Determination: Wetland  Nonwetland

## Notes:

Meadow above cave pool has *Juncus balticus* and *leymus cinereus* (7m x 10m =70 m<sup>2</sup> area); *Typha domingensis*, *Tamarix ramossissima*, *Salix goodingii*, *Polypogon monspeliensis* are other species in the area. Cattails occur under large willows. *Tamarix sp* occurs on the old pond berm (i.e. dirt road). <sup>a</sup> Wetland indicator status for plants in region 8. UPL = Upland plant species. FACW\* = tentative assignment as a facultative wetland plant species based on limited information available. FACU = Facultative upland plant species.

Name: PDG Wetland Unit: Seep under the willow trees  
 Location: Cane Spring UTM Coordinates Easting: 5-80-775 Northing: 40-72-730  
 Date: 9-9-96

**Hydrology**

Type: Seep  Spring  Pond  Detention basin  Stream  Mechanically contained   
 Source: Natural  Man-enhanced  Man-made  Ephemeral  Permanent  Temporary   
 Date of construction/Period of flow: Unknown period of flow  
 Disturbance type (if any) and date: Channel dug out by man leads to a dry pond  
 Inundated: Yes  No  Depth of standing water 2-3 cm; Saturated: Yes  No  Depth to saturation 0  
 Other field indicators: Water flow is present under willow \ fenced area  
 Atypical situation: Yes  No ; Wetland hydrology: Yes  No   
 Basis: Surface water exists at the observation point.

**Vegetation** List 3 dominant species, % cover in bold, in each vegetation layer (5 if only 1 or 2 layers are present)

Species	Indicator Status <sup>a</sup>	% Cover
<b>Trees</b>		
1. <u>Salix goodingii</u>	<u>FACW</u>	<u>90</u>
2. _____	_____	_____
3. _____	_____	_____
4. _____	_____	_____
5. _____	_____	_____
<b>Shrubs</b>		
1. _____	_____	_____
2. _____	_____	_____
3. _____	_____	_____
4. _____	_____	_____
5. _____	_____	_____
<b>Herbs</b>		
1. <u>Juncus balticus</u>	<u>FACW</u>	<u>30</u>
2. <u>Leymus cinereus</u>	<u>FACU</u>	<u>40</u>
3. <u>Typha domingensis</u>	<u>OBL</u>	<u>15</u>
4. _____	_____	_____
5. _____	_____	_____
6. _____	_____	_____
7. _____	_____	_____
8. _____	_____	_____
9. _____	_____	_____
10. _____	_____	_____

Percentage of species that are OBL, FACW, and/or FAC: 75%; Hydrophytic vegetation: Yes  No   
 Basis: The observation point has a dominance of hydrophytic vegetation. Total wetland area estimated to be about 230m<sup>2</sup>.

**Hydric Soils**

Field indicators Soil mottling, dark organic soils (low chroma), and soil saturation for greater than 7 days duration are present indicating hydric soils exist at the observation points.

Hydric Soils: Yes  No

Jurisdictional Wetland Determination : Wetland  Nonwetland

**Notes :**

Drainage area appears excavated by man, i.e. man made channel. Polypogon monspeliensis occurs in the spring area.  
Hydrobid snails are absent from east channel (where they occurred in 1988) but are now present only in the cave pool.  
<sup>a</sup> Wetland indicator status for plants in region 8. FACU = Facultative upland species. FACW = Facultative wetland species  
 OBL = Obligate wetland species.

# Routine Jurisdictional Wetland Determination

Name: PDG, DJH, JAA Wetland Unit: Drainage channel below pool  
 Location: Captain Jack Spring UTM Coordinates Easting: 5-73-834 Northing: 41-13-668  
 Date: 6-19-96

## Hydrology

Type: Seep  Spring  Pond  Detention basin  Stream  Mechanically contained   
 Source: Natural  Man-enhanced  Man-made  Ephemeral  Permanent  Temporary   
 Date of construction/Period of flow: In 1977, pipes were installed which lead to watering tanks. Pipes no longer exist. A metal tank (now dry) was bolted to rocks adjacent to the wash channel. A flood damaged tank exists in the wash.  
 Disturbance type and date: Heavy feral horse use has apparently impacted upland vegetation near the spring entrance.  
 Inundated: Yes  No  Depth of standing water 20cm; Saturated: Yes  No  Depth to saturation 0  
 Other field indicators: The spring consists of a small pool (24" x 30") and about 8" deep below some rock ledges with a surface outflow varies from 20-50cm wide by 2 cm deep. Surface flow out of the pool is < 1L/min on 6-19-96.  
 Atypical situation: Yes  No ; Wetland hydrology: Yes  No   
 Basis: Surface water exists at the observation point.

**Vegetation** List 3 dominant species, % Cover in bold, in each vegetation layer (5 if only 1 or 2 layers are present)

	Species	Indicator Status	% Cover
--	---------	------------------	---------

### Trees

1.	<u>none</u>	<u></u>	<u></u>
2.	<u></u>	<u></u>	<u></u>
3.	<u></u>	<u></u>	<u></u>
4.	<u></u>	<u></u>	<u></u>
5.	<u></u>	<u></u>	<u></u>

### Shrubs

1.	<u>none</u>	<u></u>	<u></u>
2.	<u></u>	<u></u>	<u></u>
3.	<u></u>	<u></u>	<u></u>
4.	<u></u>	<u></u>	<u></u>
5.	<u></u>	<u></u>	<u></u>

### Herbs

1.	<u>Mimulus guttatus</u>	<u>OBL</u>	<u>10</u>
2.	<u>Potentilla biennis</u>	<u>FAC</u>	<u>5</u>
3.	<u>Rumex salicifolius</u>	<u>FACW*</u>	<u>50</u>
4.	<u>Veronica anagallis-aquatica</u>	<u>OBL</u>	<u>10</u>
5.	<u></u>	<u></u>	<u></u>
6.	<u></u>	<u></u>	<u></u>
7.	<u></u>	<u></u>	<u></u>
8.	<u></u>	<u></u>	<u></u>
9.	<u></u>	<u></u>	<u></u>

Other field indicators:   
 Percentage of species that are OBL, FACW, and/or FAC: 100%; Hydrophytic vegetation: Yes  No   
 Basis: The observation point has a dominance of hydrophytic vegetation. Total wetland area is about 30m<sup>2</sup>.

### Hydric Soils

Field indicators: Hydric soils exist based on the presence of saturated soils for 7 or more days duration.

Hydric Soils: Yes  No

Jurisdictional Wetland Determination : Wetland  Nonwetland

### Notes:

Drainage from the spring pool continues for ≈ (30m length by 1m width) = 30m<sup>2</sup> surface area of saturated soils. Soils are very rocky, but with moderate accumulation of fines in the lower end of drainage. <sup>a</sup> Wetland indicator status for plants in region 8.  
\* Indicates a tentative assignment to the facultative wetland category based on limited information for this species in region 8.  
FACW = Facultative wetland species. FAC = Facultative wetland species. OBL = Obligate wetland species.



# Routine Jurisdictional Wetland Determination

Name: PDG, WKO Wetland Unit: Wash slope  
 Location: Coyote Spring UTM Coordinates Easting: 5-83-561 Northing: 40-66-755  
 Date: 9-4-96

## Hydrology

Type: Seep  Spring  Pond  Detention basin  Stream  Mechanically contained   
 Source: Natural  Man-enhanced  Man-made  Ephemeral  Permanent  Temporary   
 Date of construction/Period of flow: N/A  
 Disturbance type (if any) and date: None  
 Inundated: Yes  No  Depth of standing water 0; Saturated: Yes  No  Depth to saturation 0  
 Other field indicators: Dark colored, wet surface soil found at two locations in the wash.  
 Atypical situation: Yes  No ; Wetland hydrology: Yes  No   
 Basis: Surface water occurs at two locations at the spring area.

## Vegetation

*List 3 dominant species, % cover in bold, in each vegetation layer (5 if only 1 or 2 layers are present)*

### Trees

1.	Species	Indicator Status <sup>a</sup>	% Cover
1.	<u>none</u>	<u></u>	<u></u>
2.	<u></u>	<u></u>	<u></u>
3.	<u></u>	<u></u>	<u></u>
4.	<u></u>	<u></u>	<u></u>
5.	<u></u>	<u></u>	<u></u>

### Shrubs

1.	<u>none</u>	<u></u>	<u></u>
2.	<u></u>	<u></u>	<u></u>
3.	<u></u>	<u></u>	<u></u>
4.	<u></u>	<u></u>	<u></u>
5.	<u></u>	<u></u>	<u></u>

### Herbs

1.	<u><i>Distichlis spicata</i></u>	<u>FAC+*</u>	<u>60</u>
2.	<u></u>	<u></u>	<u></u>
3.	<u></u>	<u></u>	<u></u>
4.	<u></u>	<u></u>	<u></u>
5.	<u></u>	<u></u>	<u></u>
6.	<u></u>	<u></u>	<u></u>
7.	<u></u>	<u></u>	<u></u>
8.	<u></u>	<u></u>	<u></u>
9.	<u></u>	<u></u>	<u></u>
10.	<u></u>	<u></u>	<u></u>

Other field indicators: Dark soil  
 Percentage of species that are OBL, FACW, and/or FAC: 100%; Hydrophytic vegetation: Yes  No   
 Basis: The observation point has hydrophytic vegetation. Area of wetland was estimated to be about 200m<sup>2</sup>.

## Hydric Soils:

Field indicators: Hydric soils exist based on the presence of saturated soils for a period of 7 days or greater duration.

Hydric Soils: Yes  No

Jurisdictional Wetland Determination : Wetland  Nonwetland

## Notes:

*Kochia scoparius* (FACU), a facultative upland species occurs nearby. *Leymus cinereus* (FACU) occurs in a rocky wash 100m upstream from the observation point. <sup>a</sup> Wetland indicator status for plants in region 8. OBL = Obligate wetland species. FAC+\* = Tentative assignment to the facultative wetland category based on limited information available for this species in region 8.

# Routine Jurisdictional Wetland Determination

Name: PDG Wetland Unit: Ephemeral Pond  
 Location: Gold Meadows Spring UTM Coordinates Easting: 5-70-450 Northing: 41-20-440  
 Date: 7-22-96

## Hydrology

Type: Seep  Spring  Pond  Detention basin  Stream  Mechanically contained   
 Source: Natural  Man-enhanced  Man-made  Ephemeral  Permanent  Temporary   
 Date of construction/Period of flow: There is no surface outflow from the pond.  
 Disturbance type (if any) and date: Southwest side of the pond has a man-made berm. This was possibly constructed by man to deepen the pond for livestock use. Date of excavation is unknown.  
 Inundated: Yes  No  Depth of standing water 0; Saturated: Yes  No  Depth to saturation 0  
 Other field indicators: Dark organic matter, damp mud present - water mark on rocks  
 Atypical situation: Yes  No ; Wetland hydrology: Yes  No   
 Basis: Water marks exist on large rocks at the observation point. Water was present earlier in the year. Survey was conducted during a very dry year.

## Vegetation List 3 dominant species, % cover in bold, in each vegetation layer (5 if only 1 or 2 layers are present)

	<i>Species</i>	<i>Indicator Status<sup>a</sup></i>	<i>% Cover</i>
<b>Trees</b>			
1.	<u>none</u>	<u></u>	<u></u>
2.	<u></u>	<u></u>	<u></u>
3.	<u></u>	<u></u>	<u></u>
4.	<u></u>	<u></u>	<u></u>
5.	<u></u>	<u></u>	<u></u>
<b>Shrubs</b>			
1.	<u>none</u>	<u></u>	<u></u>
2.	<u></u>	<u></u>	<u></u>
3.	<u></u>	<u></u>	<u></u>
4.	<u></u>	<u></u>	<u></u>
5.	<u></u>	<u></u>	<u></u>
<b>Herbs</b>			
1.	<u><i>Juncus balticus</i></u>	<u>FACW</u>	<u>15</u>
2.	<u></u>	<u></u>	<u></u>
3.	<u></u>	<u></u>	<u></u>
4.	<u></u>	<u></u>	<u></u>
5.	<u></u>	<u></u>	<u></u>
6.	<u></u>	<u></u>	<u></u>
7.	<u></u>	<u></u>	<u></u>
8.	<u></u>	<u></u>	<u></u>
9.	<u></u>	<u></u>	<u></u>

Other field indicators:   
 Percentage of species that are OBL, FACW, and/or FAC: 100%; Hydrophytic vegetation: Yes  No   
 Basis: A dominance of hydrophytic plants occurs at the observation point. Wetland area estimated at about 45m<sup>2</sup>.

## Hydric Soils:

Field indicators Hydric soils exist based on presence of saturated soils for a period of 7 or greater days in duration.

Hydric Soils: Yes  No

Jurisdictional Wetland Determination : Wetland  Nonwetland

## Notes:

Sagebrush, *Artemisia tridentata* is the major upland species in the area. Basin wild rye *Leymus cinereus* occurs on the edges of the dried pond. Horse, deer and antelope use of the spring is significant. <sup>a</sup> Wetland indicator status for plants in region 8. FACW = Facultative wetland species.

CHVI, EPNE, ARTR. <sup>a</sup> Wetland indicator status for plants in region 8. FACU = Facultative upland species. UPL = Upland species. OBL = Obligate wetland species.

---

# Routine Jurisdictional Wetland Determination

Name: PDG, DJH Wetland Unit: Ledge pool  
 Location: John's Spring UTM Coordinates Easting: 5-82-100 Northing: 41-22-490  
 Date: 12-18-96

## Hydrology

Type: Seep  Spring  Pond  Detention basin  Stream  Mechanically contained   
 Source: Natural  Man-enhanced  Man-made  Ephemeral  Permanent  Temporary   
 Date of construction/Period of flow: None  
 Disturbance type (if any) and date: None  
 Inundated: Yes  No  Depth of standing water 2-5cm; Saturated: Yes  No  Depth to saturation 0  
 Other field indicators: \_\_\_\_\_  
 Atypical situation: Yes  No ; Wetland hydrology: Yes  No   
 Basis: Surface water exists at the observation point.

**Vegetation** List 3 dominant species, % cover in bold, in each vegetation layer (5 if only 1 or 2 layers are present)

	Species	Indicator Status <sup>a</sup>	% Cover
<b>Trees</b>			
1.	<u>none</u>	<u>_____</u>	<u>_____</u>
2.	<u>_____</u>	<u>_____</u>	<u>_____</u>
3.	<u>_____</u>	<u>_____</u>	<u>_____</u>
4.	<u>_____</u>	<u>_____</u>	<u>_____</u>
5.	<u>_____</u>	<u>_____</u>	<u>_____</u>
<b>Shrubs</b>			
1.	<u>Rhus trilobata</u>	<u>NI</u>	<u>20*</u>
2.	<u>_____</u>	<u>_____</u>	<u>_____</u>
3.	<u>_____</u>	<u>_____</u>	<u>_____</u>
4.	<u>_____</u>	<u>_____</u>	<u>_____</u>
5.	<u>_____</u>	<u>_____</u>	<u>_____</u>
<b>Herbs</b>			
1.	<u>Carex praegracilis</u>	<u>FACW-</u>	<u>20</u>
2.	<u>Gallium aparine</u>	<u>FACU</u>	<u>Tr</u>
3.	<u>Leymus cinereus</u>	<u>FACU</u>	<u>60*</u>
4.	<u>Mimulus guttatus</u>	<u>OBL</u>	<u>80</u>
5.	<u>Penstemon sp.</u>	<u>UNKN</u>	<u>Tr.</u>
6.	<u>Oenothera cespitosa ssp. marginata</u>	<u>NL</u>	<u>Tr.</u>
7.	<u>_____</u>	<u>_____</u>	<u>_____</u>
8.	<u>_____</u>	<u>_____</u>	<u>_____</u>
9.	<u>_____</u>	<u>_____</u>	<u>_____</u>

Percentage of species that are OBL, FACW, and/or FAC: 100%; Hydrophytic vegetation: Yes  No   
 Basis: The observation point has a dominance of hydrophytic vegetation. Size of wetland area estimated at about 50m<sup>2</sup>.

## Hydric Soils

Field indicators: Hydric soils exist based on the presence of saturated soils for 7 days or greater duration.  
 Hydric Soils: Yes  No

Jurisdictional Wetland Determination : Wetland  Nonwetland

## Notes:

<sup>a</sup> Wetland indicator status for region 8. \**R. trilobata* and *L. cinereus* occur in the transition zone between the jurisdictional wetland and the upland habitat. OBL = Obligate wetland species. FACW- = Facultative wetland species. FACU = Facultative upland plant species. NI = Insufficient information to determine wetland status for this species in region 8. NL = not listed in National List of Plants that occur in Wetlands for Region 8. UNKN = Unknown wetland status.

5.	_____	_____	_____
<b>Herbs</b>			
1.	<i>Leymus cinereus</i>	FACU	30*
2.	_____	_____	_____
3.	_____	_____	_____
4.	_____	_____	_____
5.	_____	_____	_____
6.	_____	_____	_____
7.	_____	_____	_____
8.	_____	_____	_____
9.	_____	_____	_____
10.	_____	_____	_____

Other field indicators: \_\_\_\_\_  
 Percentage of species that are OBL, FACW, and/or FAC: 100 %; Hydrophytic vegetation: Yes X No \_\_\_\_\_  
 Basis: A dominance of hydrophytic vegetation occurs at the observation point.

**Hydric Soils**

Field indicators: Hydric soils exist based on the presence saturated soils for 7 or greater days duration .

Hydric Soils: Yes X No \_\_\_\_\_

Jurisdictional Wetland Determination : Wetland X Nonwetland \_\_\_\_\_

**Notes:**

\* *Leymus cinereus* and *Rhus trilobata* are present on the edge of the delineated wetland and comprised about 30 % cover of this area. <sup>a</sup> Wetland indicator status for plants in region 8. FACW = Facultative wetland species. NI = Insufficient information to determine wetlands status of this species in region 8. FACU = Facultative upland species.

**Vegetation** *List 3 dominant species (% cover in bold) in each vegetation layer (5 if only 1 or 2 layers are present)*

	<i>Species</i>	<i>Indicator Status<sup>a</sup></i>	<i>% Cover</i>
<b>Trees</b>			
1.	none		
2.			
3.			
4.			
5.			
<b>Shrubs</b>			
1.	<i>Ericameria nauseosa</i>	NL	40
2.			
3.			
4.			
5.			
<b>Herbs</b>			
1.	<i>Dactylus glomerata</i>	FACU	10
2.	<i>Sporobolus airoides</i>	FAC-	10
3.			
4.			
5.			
6.			
7.			
8.			
9.			
10.			

Other field indicators: Mesic grasses including *Poa sp.* and *Camissonia sp.*  
 Percentage of species that are OBL, FACW, and/or FAC: 33 %; **Hydrophytic vegetation:** Yes      No X  
 Basis: A dominance of hydrophytic vegetation is not present at the observation site.

**Hydic soils:**  
 Field indicators: None - no inundation (saturated soils) during surveys in September 1996.

Hydic soils: Yes      No X

Jurisdictional Wetland Determination : Wetland      Nonwetland X

**Notes:**  
Wetlands survey was performed during a very dry year. <sup>a</sup> Wetland indicator status for plants in region 8. FAC- = Facultative wetland species. FACU = Facultative upland species. NL = Not listed on the National List of Plants that occur in Wetlands for Region 8.

1. A. C. H. G. or species that are O. B. S., F. H. C. W., and/or F. H. C. \_\_\_\_\_ No, hydrophytic vegetation: Yes \_\_\_\_\_ No X  
Basis: Lack of dominance of hydrophytic vegetation at the observation point.

**Hydric Soils**

Field indicators: Hydric soils are absent because saturated soils are not present for 7 days or greater duration.

Hydric Soils: Yes \_\_\_\_\_ No X

Jurisdictional Wetland Determination : Wetland \_\_\_\_\_ Nonwetland X

Notes: Records from Moore (1961), (USGS report TEI-781) indicated water (flow) was present on several dates in the fall of 1957, but the spring was dry in October 1960. <sup>a</sup> Wetland indicator status for plants in region 8. FACU = Facultative upland species.



# Routine Jurisdictional Wetland Determination

Name: PDG, DJH Wetland Unit: Solution crevice in limestone outcrop  
 Location: Rock Valley Tank UTM Coordinates Easting: 5-68-070 Northing: 40-61-000  
 Date: 1-7-97

## Hydrology

Type: Seep  Spring  Pond  Detention basin  Stream  Natural Tank   
 Source: Natural  Man-enhanced  Man-made  Ephemeral  Permanent  Temporary   
 Date of construction/Period of flow: Unknown  
 Disturbance type (if any) and date: None  
 Inundated: Yes  No  Depth of standing water 25-30cm; Saturated: Yes  No  Depth to saturation 0  
 Other field indicators: \_\_\_\_\_  
 Atypical situation: Yes  No ; Wetland hydrology: Yes  No   
 Basis: Surface water exists at the observation point but is confined to an opening (20 by 40 cm) in rock. Water exists in a solution cavern of unknown dimensions.

## Vegetation *List 3 dominant species, percent cover in bold, in each vegetation layer (5 if only 1 or 2 layers are present)*

Species	Indicator Status <sup>a</sup>	% Cover
<b>Trees</b>		
1. <u>none</u>	_____	_____
2. _____	_____	_____
3. _____	_____	_____
4. _____	_____	_____
5. _____	_____	_____
<b>Shrubs</b>		
1. <u>none</u>	_____	_____
2. _____	_____	_____
3. _____	_____	_____
4. _____	_____	_____
5. _____	_____	_____
<b>Herbs</b>		
1. <u><i>Bromus rubens</i></u>	<u>UPL</u>	<u>20</u>
2. _____	_____	_____
3. _____	_____	_____
4. _____	_____	_____
5. _____	_____	_____
6. _____	_____	_____
7. _____	_____	_____
8. _____	_____	_____
9. _____	_____	_____

Other field indicators: \_\_\_\_\_  
 Percentage of species that are OBL, FACW, and/or FAC: 0%; Hydrophytic vegetation: Yes  No   
 Basis: Hydrophytic vegetation is absent from the observation site.

## Hydric Soils

Field indicators: Hydric soils appear to be absent because surface water is confined to a limestone rock opening. An area of dark mesic soil accumulation (5m by 10m) occurs below the limestone outcrop. Soil depths in this area appear shallow although soil pits were not dug in this area.

Hydric Soils: Yes  No

Jurisdictional Wetland Determination : Wetland  Nonwetland

## Notes:

Heavy coyote use of the water source is indicated by numerous scats in the area. <sup>a</sup> Wetland indicator status for plants in region 8. UPL = Upland plant species.



# Routine Jurisdictional Wetland Determination

Name: PDG, JAA Wetland Unit: Wash channel  
 Location: Wahmonie Seep 2 UTM Coordinates Easting: 5-77-597 Northing: 40-73-418  
 Date: 6-20-96

## Hydrology

Type: Seep  Spring  Pond  Detention basin  Stream  Mechanically contained   
 Source: Natural  Man-enhanced  Man-made  Ephemeral  Permanent  Temporary   
 Date of construction/Period of flow: None/Unknown availability of water  
 Disturbance type (if any) and date: None  
 Inundated: Yes  No  Depth of standing water 0; Saturated: Yes  No  Depth to saturation 12 inches  
 Other field indicators: Surface water was present at this seep on June 6, 1996 but not on June 20, 1996.  
 Atypical situation: Yes  No ; Wetland hydrology: Yes  No   
 Basis: Surface water was present at the site during 1996.

## Vegetation *List 3 dominant species, percent cover in bold, in each vegetation layer (5 if only 1 or 2 layers are present)*

	<i>Species</i>	<i>Indicator Status<sup>a</sup></i>	<i>% Cover</i>
<b>Trees</b>			
1.	<u>none</u>	<u></u>	<u></u>
2.	<u></u>	<u></u>	<u></u>
3.	<u></u>	<u></u>	<u></u>
4.	<u></u>	<u></u>	<u></u>
5.	<u></u>	<u></u>	<u></u>
<b>Shrubs</b>			
1.	<u>Baccharis emoryi</u>	<u>FACW</u>	<u>85</u>
2.	<u></u>	<u></u>	<u></u>
3.	<u></u>	<u></u>	<u></u>
4.	<u></u>	<u></u>	<u></u>
5.	<u></u>	<u></u>	<u></u>
<b>Herbs</b>			
1.	<u>Artemisia ludoviciana</u>	<u>FACU</u>	<u>2</u>
2.	<u>Moss</u>	<u></u>	<u>1</u>
3.	<u></u>	<u></u>	<u></u>
4.	<u></u>	<u></u>	<u></u>
5.	<u></u>	<u></u>	<u></u>
6.	<u></u>	<u></u>	<u></u>
7.	<u></u>	<u></u>	<u></u>
8.	<u></u>	<u></u>	<u></u>
9.	<u></u>	<u></u>	<u></u>
10.	<u></u>	<u></u>	<u></u>

Other field indicators: \_\_\_\_\_  
 Percentage of species that are OBL, FACW, and/or FAC: 100 %; Hydrophytic vegetation: Yes  No   
 Basis: A dominance of hydrophytic plants occurs at the observation point. Wetland area was estimated to be about 150m<sup>2</sup>.

## Hydric soils

Field indicators: Hydric soils exist based on the presence of saturated soils for a period of 7 days or greater duration.

Hydric Soils: Yes  No

Jurisdictional Wetland Determination : Wetland  Nonwetland

## Notes:

Soils are very rocky. Wetland dimensions are approximately 30m x 5 m. Damp soil exists under plants in wash. <sup>a</sup> Wetland indicator status for plants in region 8. FACW = Facultative wetland species. FACU = Facultative upland species.

# Routine Jurisdictional Wetland Determination

Name: JAA, PDG Wetland Unit: Seep channel  
 Location: Wahmonie Seep 3 UTM Coordinates Easting: 5-77-044 Northing: 40-73-438  
 Date: 6-20-96

## Hydrology

Type: Seep  Spring  Pond  Detention basin  Stream  Mechanically contained   
 Source: Natural  Man-enhanced  Man-made  Ephemeral  Permanent  Temporary   
 Date of construction/Period of flow: None/Unknown seasonal availability of water  
 Disturbance type (if any) and date: None  
 Inundated: Yes  No  Depth of standing water 0; Saturated: Yes  No  Depth to saturation undetermined  
 Other field indicators: Water marks on rocks in wash. Surface water was present on June 6, 1996 but not on June 20, 1996  
 Atypical situation: Yes  No ; Wetland hydrology: Yes  No   
 Basis: Field indicators for surface water were recorded at the observation point.

## Vegetation *List 3 dominant species, percent cover in bold, in each vegetation layer (5 if only 1 or 2 layers are present)*

Species	Indicator Status <sup>a</sup>	% Cover
<b>Trees</b>		
1. <u>none</u>	<u></u>	<u></u>
2. <u></u>	<u></u>	<u></u>
3. <u></u>	<u></u>	<u></u>
4. <u></u>	<u></u>	<u></u>
5. <u></u>	<u></u>	<u></u>
<b>Shrubs</b>		
1. <u>Baccharis emoryi</u>	<u>FACW</u>	<u>60</u>
2. <u></u>	<u></u>	<u></u>
3. <u></u>	<u></u>	<u></u>
4. <u></u>	<u></u>	<u></u>
5. <u></u>	<u></u>	<u></u>
<b>Herbs</b>		
1. <u>Artemisia ludoviciana</u>	<u>FACU</u>	<u>10</u>
2. <u>Bromus rubens</u>	<u>UPL</u>	<u>30</u>
3. <u></u>	<u></u>	<u></u>
4. <u></u>	<u></u>	<u></u>
5. <u></u>	<u></u>	<u></u>
6. <u></u>	<u></u>	<u></u>
7. <u></u>	<u></u>	<u></u>
8. <u></u>	<u></u>	<u></u>
9. <u></u>	<u></u>	<u></u>
10. <u></u>	<u></u>	<u></u>

Other field indicators:   
 Percentage of species that are OBL, FACW, and/or FAC: 33 %; Hydrophytic vegetation: Yes  No   
 Basis: A dominance of hydrophytic vegetation occurs at the observation site. Area of wetland estimated to be about 180m<sup>2</sup>.

## Hydric Soils:

Field indicators: Hydric soils exist based on the presence of saturated soils for 7 days or greater duration during 1996.

Hydric Soils: Yes  No

Jurisdictional Wetland Determination : Wetland  Nonwetland

## Notes:

Soils are very thin next to bedrock in the wash bottom. Salt encrusted water marks along the wash occur for 50-60m distance. (Water marks show a 3 meter width). Artemisia ludoviciana 10 % cover, and Bromus rubens 30% cover, occur on the edges of the wash. <sup>a</sup> = Wetland indicator status for plants in region 8. FACW = Facultative wetland species. FACU = Facultative upland species. UPL = Upland plant species.



# Routine Jurisdictional Wetland Determination

Name: DJH, PDG, JAA Wetland Unit: 2- Middle stretch of the spring channel  
 Location: Tippipah Spring UTM Coordinates Easting: 5-70-810 Northing: 40-99-723  
 Date: 6-18-96

## Hydrology

Type: Seep  Spring  Pond  Detention basin  Stream  Mechanically contained   
 Source: Natural  Man-enhanced  Man-made  Ephemeral  Permanent  Temporary   
 Date of construction/Period of flow: None/ unknown  
 Disturbance type (if any) and date: None  
 Inundated: Yes  No  Depth of standing water 10-15cm; Saturated: Yes  No  Depth to saturation 0  
 Other field indicators: \_\_\_\_\_  
 Atypical situation: Yes  No ; Wetland hydrology: Yes  No   
 Basis: Surface water occurs at the observation point. Flow rate measured about 80m downstream from the spring source on November 15, 1996 was about 2.7 L/per min.

## Vegetation List 3 dominant species, percent cover in bold, in each vegetation layer (5 if only 1 or 2 layers are present)

	Species	Indicator Status <sup>a</sup>	% Cover
<b>Trees</b>			
1.	<u>none</u>	<u>_____</u>	<u>_____</u>
2.	<u>_____</u>	<u>_____</u>	<u>_____</u>
3.	<u>_____</u>	<u>_____</u>	<u>_____</u>
4.	<u>_____</u>	<u>_____</u>	<u>_____</u>
5.	<u>_____</u>	<u>_____</u>	<u>_____</u>
<b>Shrubs</b>			
1.	<u>none</u>	<u>_____</u>	<u>_____</u>
2.	<u>_____</u>	<u>_____</u>	<u>_____</u>
3.	<u>_____</u>	<u>_____</u>	<u>_____</u>
4.	<u>_____</u>	<u>_____</u>	<u>_____</u>
<b>Herbs</b>			
1.	<u>Bromus rubens</u>	<u>UPL</u>	<u>Tr.</u>
2.	<u>Bromus tectorum</u>	<u>NL</u>	<u>2</u>
3.	<u>Castilleja sp.</u>	<u>UNKN</u>	<u>Tr.</u>
4.	<u>Deschampsia danthonioides</u>	<u>FACW</u>	<u>Tr.</u>
5.	<u>Eleocharis parishii</u>	<u>OBL</u>	<u>Tr.</u>
6.	<u>Epilobium glaberrimum</u>	<u>FACW</u>	<u>2</u>
7.	<u>Heliomeris multiflora var nevadensis</u>	<u>NL</u>	<u>2</u>
8.	<u>Juncus balticus</u>	<u>FACW</u>	<u>40</u>
9.	<u>Juncus longistylis</u>	<u>FACW+</u>	<u>Tr.</u>
10.	<u>Lactuca serriola</u>	<u>FACU</u>	<u>2</u>
11.	<u>Polypogon monspeliensis</u>	<u>FACW+</u>	<u>30</u>
12.	<u>Verbena bracteata</u>	<u>FACU</u>	<u>2</u>
13.	<u>Veronica anagallis-aquatica</u>	<u>OBL</u>	<u>20</u>

Other field indicators: \_\_\_\_\_  
 Percentage of species that are OBL, FACW, and/or FAC: 100 %; Hydrophytic vegetation: Yes  No   
 Basis: A dominance of hydrophytic vegetation exists at the observation point.

## Hydric Soils

Field indicators: Hydric soils exist based on the presence of saturated soils for 7 days or greater duration.

Hydric Soils: Yes  No

Jurisdictional Wetland Determination : Wetland  Nonwetland

## Notes:

Obsidian flakes found at the spring suggest Native American use. Tr = Trace amounts (<1 % cover). <sup>a</sup> = Wetland indicator status for plants in region 8. OBL = Obligate wetland species. FACW, FACW+ are both facultative wetland species. FACU = Facultative upland species. NL = not listed in National List of Plants that occur in Wetlands for Region 8. UPL = Upland species. UNKN = Unknown status in region 8.

Percentage of species that are OBL, FACW, and/or FAC: 100 %; Hydrophytic vegetation: Yes X No \_\_\_\_\_  
Basis: A dominance of hydrophytic plants occurs at the observation point. Size of wetland was estimated at about 500m<sup>2</sup>.

**Hydric Soils**

Field indicators: Hydric soils exist based on the presence of saturated soils for a period of 7 days or greater duration.

Hydric Soils: Yes X No \_\_\_\_\_

Jurisdictional Wetland Determination : Wetland X Nonwetland \_\_\_\_\_

**Notes:**

Artemisia tridentata, Ericameria nauseosa encroaching on edge of wetland. A large water tank exists near lower end of the spring channel. On 6-18-96, water flowed 170 m down the wash to the old water tank. <sup>a</sup> = Wetlands indicator status for plants in region 8. Tr = Trace amounts (<1% cover). OBL = Obligate wetland species. FAC, FACW, FACW+ are all types of facultative wetland species. FACU = Facultative upland species. NL = not listed in National List of Plants that occur in Wetlands for Region 8.

# Routine Jurisdictional Wetland Determination

Name: PDG, AA Wetland Unit: Rock water tank in cave  
 Location: Tongue Wash Tank UTM Coordinates Easting: 5-71-360 Northing: 41-13-050  
 Date: 9-10-96

## Hydrology

Type: Seep  Spring  Pond  Detention basin  Stream  Natural tank   
 Source: Natural  Man-enhanced  Man-made  Ephemeral  Permanent  Temporary   
 Date of construction/Period of flow: It is unknown if water flows out of the tank during any time.  
 Disturbance type (if any) and date: None Inundated: Yes  No   
 Depth of standing water 20-25cm; Saturated: Yes  No  Depth to saturation 0  
 Other field indicators: A dark water mark on the rocks indicate a previous water level that is higher than the present water level.  
 Atypical situation: Yes  No ; Wetland hydrology: Yes  No   
 Basis: Surface water exists at the observation point.

## Vegetation *List 3 dominant species, percent cover in bold, in each vegetation layer (5 if only 1 or 2 layers are present)*

	<i>Species</i>	<i>Indicator Status<sup>a</sup></i>	<i>% Cover</i>
<b>Trees</b>			
1.	<u>none</u>	<u></u>	<u></u>
2.	<u></u>	<u></u>	<u></u>
3.	<u></u>	<u></u>	<u></u>
4.	<u></u>	<u></u>	<u></u>
5.	<u></u>	<u></u>	<u></u>

<b>Shrubs</b>			
1.	<u>none</u>	<u></u>	<u></u>
2.	<u></u>	<u></u>	<u></u>
3.	<u></u>	<u></u>	<u></u>
4.	<u></u>	<u></u>	<u></u>
5.	<u></u>	<u></u>	<u></u>

<b>Herbs</b>			
1.	<u>none</u>	<u></u>	<u></u>
2.	<u></u>	<u></u>	<u></u>
3.	<u></u>	<u></u>	<u></u>
4.	<u></u>	<u></u>	<u></u>
5.	<u></u>	<u></u>	<u></u>
6.	<u></u>	<u></u>	<u></u>
7.	<u></u>	<u></u>	<u></u>
8.	<u></u>	<u></u>	<u></u>
9.	<u></u>	<u></u>	<u></u>
10.	<u></u>	<u></u>	<u></u>

Other field indicators: Some grasses and annuals exist in a small unsaturated area (2-3 m<sup>2</sup>) below the cave opening where water may seep out or overflow the tank during very wet years.  
 Percentage of species that are OBL, FACW, and/or FAC: 0%; Hydrophytic vegetation: Yes  No

Basis: Hydrophytic vegetation is absent at the observation point.

## Hydric Soils

Field indicators: Hydric soils appear to be absent at the observation point because water is confined to a bedrock pool. No soil pits were dug at this site.

Hydric Soils: Yes  No

Jurisdictional Wetland Determination: Wetland  Nonwetland

## Notes:

The water tank measures about 1-2m wide by 3-4m long and occurs in a small natural cave in tuff rock formation. Numerous petroglyphs at the site indicate use by Native Americans. Large numbers of birds were observed entering the cave drinking on 9-10-96. <sup>a</sup> = Wetlands indicator status for plants in region 8.

# Routine Jurisdictional Wetland Determination

Name: PDG, JAA Wetland Unit: Cave Pool  
 Location: Topopah Spring UTM Coordinates Easting: 5-65-024 Northing: 40-88-369  
 Date: 6-20-96

## Hydrology

Type: Seep  Spring  Pond  Detention basin  Stream  Mechanically contained   
 Source: Natural  Man-enhanced  Man-made  Ephemeral  Permanent  Temporary   
 Date of construction/Period of flow: Unknown  
 Disturbance type (if any) and date: Pipe was installed in the ground below the cave pool and has flow. Fire burned the area around the spring - date unknown: Cave pool was dug out to increase access to water. Dates of disturbance unknown.  
 Inundated: Yes  No  Depth of standing water 15-20cm; Saturated: Yes  No  Depth to saturation 0  
 Other field indicators: \_\_\_\_\_  
 Atypical situation: Yes  No ; Wetland hydrology: Yes  No   
 Basis: Surface water exists at the observation point. Total area of wetland (all habitats combined) estimated at about 200m<sup>2</sup>. Flow rate measured from an existing pipe was estimated at 0.140 L/min on September 9 1996.

## Vegetation *List 3 dominant species, percent cover in bold, in each vegetation layer (5 if only 1 or 2 layers are present)*

Species	Indicator Status <sup>a</sup>	% Cover
<b>Trees</b>		
1. <u>none</u>	_____	_____
2. _____	_____	_____
3. _____	_____	_____
4. _____	_____	_____
5. _____	_____	_____
<b>Shrubs</b>		
1. <u>none</u>	_____	_____
2. _____	_____	_____
3. _____	_____	_____
4. _____	_____	_____
5. _____	_____	_____
<b>Herbs</b>		
1. <u>Artemisia ludoviciana</u>	<u>FACU</u>	<u>5</u>
2. <u>Bromus diandrus</u>	<u>NL</u>	<u>1</u>
3. <u>Epilobium glaberrimum</u>	<u>FACW</u>	<u>5</u>
4. <u>Mimulus guttatus</u>	<u>OBL</u>	<u>10</u>
5. <u>Polypogon monspeliensis</u>	<u>FACW+</u>	<u>1</u>
6. <u>Potentilla biennis</u>	<u>FAC</u>	<u>2</u>
7. <u>Rumex salicifolius</u>	<u>FACW*</u>	<u>15</u>
8. <u>Sisymbrium altissimum</u>	<u>FACU-</u>	<u>1</u>
9. <u>Veronica anagallis-aquatica</u>	<u>OBL</u>	<u>20</u>
10. <u>Pseudognaphalium stramineum</u>	<u>FAC</u>	<u>5</u>

Other field indicators: \_\_\_\_\_  
 Percentage of species that are OBL, FACW, and/or FAC: 100%; Hydrophytic vegetation: Yes  No   
 Basis: Hydrophytic vegetation is present at the observation point.

## Hydric Soils

Field indicators: Hydric soils exist based on the presence of saturated soils for a period of 7 days or greater duration.

Hydric Soils: Yes  No

Jurisdictional Wetland Determination : Wetland  Nonwetland

## Notes:

A pipe was installed into the ground which forms a second shallow pool with vegetation 10 meters downslope from the cave pool. <sup>a</sup> = Wetland indicator status for plants in region 8. OBL = Obligate wetland species. FAC, FACW, FACW+ are all types of facultative wetland species. FACU, FACU- are both Facultative upland species. \* = a tentative assignment to this category based on limited information for this species. NL = not listed in the National List of Plants that occur in Wetlands for Region 8.

Hydric Soils: Yes  No

Jurisdictional Wetland Determination : Wetland  Nonwetland

**Notes:**

An area of surface inundation near the bottom of the slope was about 6 ft. long by 3 ft. wide. Another inundated spot near top of the slope was 8 ft. wide x 3 ft. long. Saturated soils are about 20 m long by 3-4 m wide. Soils were dark grey, with little organic matter and no mottling was detected. Wetland vegetation on the sloped meadow is about 20x 6m in dimensions.

<sup>a</sup> = Wetlands plant indicator status for region 8. OBL = Obligate wetland species, FAC, FACW, FACW+ - are all types of facultative wetland species. FACU = Facultative upland plants. NL = not listed in the National List of Plants that occur in Wetlands for Region 8.

Date of construction/Period of flow: Unknown period of flow.

Disturbance type (if any) and date: None

Inundated: Yes  No  Depth of standing water 0; Saturated: Yes  No  Depth to saturation Unknown

Other field indicators: \_\_\_\_\_

Atypical situation: Yes  No ; Wetland hydrology: Yes  No

Basis: No field indicators of surface hydrology were detected but the survey was performed in a very dry year.

**Vegetation** List 3 dominant species, percent cover in bold, in each vegetation layer (5 if only 1 or 2 layers are present)

Species	Indicator Status <sup>a</sup>	% Cover
<b>Trees</b>		
1. <u>none</u>	_____	_____
2. _____	_____	_____
3. _____	_____	_____
4. _____	_____	_____
5. _____	_____	_____
<b>Shrubs</b>		
1. <u>Atriplex confertifolia</u>	<u>NL</u>	<u>2</u>
2. <u>Salazaria mexicana</u>	<u>NL</u>	<u>2</u>
3. <u>Ericameria nauseosa</u>	<u>NL</u>	<u>2</u>
4. _____	_____	_____
5. _____	_____	_____
<b>Herbs</b>		
1. <u>Hordeum jubatum</u>	<u>FAC+</u>	<u>60</u>
2. <u>Bromus tectorum</u>	<u>NL</u>	<u>20</u>
3. _____	_____	_____
4. _____	_____	_____
5. _____	_____	_____
6. _____	_____	_____
7. _____	_____	_____
8. _____	_____	_____
9. _____	_____	_____
10. _____	_____	_____

Other field indicators: \_\_\_\_\_

Percentage of species that are OBL, FACW, and/or FAC: 50%; Hydrophytic vegetation: Yes  No

Basis: Only a weak indication of dominance of hydrophytic vegetation was shown at the observation point.

**Hydric Soils**

Field indicators: No indicators of hydric soils were detected at the observation point.

Hydric Soils: Yes  No

Jurisdictional Wetland Determination : Wetland  Nonwetland

**Notes:**

Large unidentified composite - toothed leaves 4-5 ft. high. Animal use - coyote scats - raven flew over on survey date.

<sup>a</sup> Wetland indicator status for plants in region 8. FAC+ = Facultative wetland species. NL= not listed in National List of Plants that occur in Wetlands for Region 8.

Percentage of species that are OBL, FACW, and/or FAC \_\_\_\_\_  
Basis: No hydrophytic vegetation was present at the observation point.

**Hydric Soils:**

Field indicators: Hydric soils are present in the cave based on presence of saturated soils for greater than 7 days duration.

Hydric Soils: Yes X No \_\_\_\_\_

Jurisdictional Wetland Determination : Wetland \_\_\_\_\_ Nonwetland X

Notes: <sup>a</sup> Wetland indicator status for plants in region 8. NI = Insufficient information to determine wetland status in region 8.

2.	<i>Salazaria mexicana</i>	NL	2
3.	<i>Ericameria nauseosa</i>	NL	2
4.			
5.			

**Herbs**

1.	<i>Hordeum jubatum</i>	FAC+	60
2.	<i>Bromus tectorum</i>	NL	20
3.			
4.			
5.			
6.			
7.			
8.			
9.			
10.			

Other field indicators: \_\_\_\_\_  
 Percentage of species that are OBL, FACW, and/or FAC: 50 %; Hydrophytic vegetation: Yes \_\_\_\_\_ No X  
 Basis: Only a weak indication of dominance of hydrophytic vegetation was shown at the observation point.

**Hydric Soils**

Field indicators: No indicators of hydric soils were detected at the observation point.

Hydric Soils: Yes \_\_\_\_\_ No X

Jurisdictional Wetland Determination : Wetland \_\_\_\_\_ Nonwetland X

**Notes:**

Large unidentified composite - toothed leaves 4-5 ft. high. Animal use - coyote scats - raven flew over on survey date.  
<sup>a</sup> Wetland indicator status for plants in region 8. FAC+ = Facultative wetland species. NL = not listed in national list of plants that occur in wetlands for region 8.

# Routine Jurisdictional Wetland Determination

Name: PDG, WKO, Wetland Unit: Wash slope  
 Location: Twin Springs UTM Coordinates Easting: 5-55-484 Northing: 40-89-984  
 Date: 12-19-96

## Hydrology

Type: Seep  Spring  Pond  Detention basin  Stream  Mechanically contained   
 Source: Natural  Man-enhanced  Man-made  Ephemeral  Permanent  Temporary   
 Date of construction/Period of flow: Unknown  
 Disturbance type (if any) and date: Tunnel was dug into hillside 15-18m by man. A cave-in has occurred. No water visible in cave, but soils are saturated. Tailings excavated from cave form a terrace on hillside with stone structure/foundation on it.  
 Inundated: Yes  No  Depth of standing water 10 cm; Saturated: Yes  No  Depth to saturation 0  
 Other field indicators: Outside pool of water exists below a rock ledge.  
 Atypical situation: Yes  No ; Wetland hydrology: Yes  No   
 Basis: Surface water occurs at the observation point.

## Vegetation List 3 dominant species, % cover in bold, in each vegetation layer (5 if only 1 or 2 layers are present)

	Species	Indicator Status <sup>a</sup>	% Cover
<b>Trees</b>			
1.	<u>none</u>	<u></u>	<u></u>
2.	<u></u>	<u></u>	<u></u>
3.	<u></u>	<u></u>	<u></u>
4.	<u></u>	<u></u>	<u></u>
5.	<u></u>	<u></u>	<u></u>
<b>Shrubs</b>			
1.	<u></u>	<u></u>	<u></u>
2.	<u></u>	<u></u>	<u></u>
3.	<u></u>	<u></u>	<u></u>
4.	<u></u>	<u></u>	<u></u>
5.	<u></u>	<u></u>	<u></u>
<b>Herbs</b>			
1.	<u><i>Polypogon monspeliensis</i></u>	<u>FACW+</u>	<u>Tr.</u>
2.	<u><i>Rumex salicifolius</i></u>	<u>FACW*</u>	<u>5</u>
3.	<u><i>Typha domingensis</i></u>	<u>OBL</u>	<u>90</u>
4.	<u></u>	<u></u>	<u></u>
5.	<u></u>	<u></u>	<u></u>
6.	<u></u>	<u></u>	<u></u>
7.	<u></u>	<u></u>	<u></u>
8.	<u></u>	<u></u>	<u></u>
9.	<u></u>	<u></u>	<u></u>

Percentage of species that are OBL, FACW, and/or FAC: 100 %; Hydrophytic vegetation: Yes  No   
 Basis: The observation point has a dominance of hydrophytic vegetation. Size of the wetland estimated at about 27m<sup>2</sup>.

## Hydric Soils

Field indicators: Hydric soils exist based on the presence of saturated soils for a period of 7 or more days duration.

Hydric Soils: Yes  No

Jurisdictional Wetland Determination : Wetland  Nonwetland

## Notes:

*Artemesia dracuncululus* (NL), *A ludoviciana* (Facu), and *Rhus trilobata* (NI), occur on the edges of the wetland habitat. *R. trilobata* occupied about 50 % of the wetland edge area. <sup>a</sup> Wetland indicator status for plants in region 8. OBL = Obligate wetland species. FACW+ = a facultative wetland species. FACW\* refers to a tentative assignment of this species to a facultative wetland category based on limited information in region 8. Tr = trace amounts (< 1 % cover).

Name: PDG, DJH, JAA Wetland Unit: 1 - Upper stretch of spring channel  
 Location: Whiterock Spring UTM Coordinates Easting: 5-77-015 Northing: 41-17-396  
 Date: 6-18-96

**Hydrology**

Type: Seep  Spring  Pond  Detention basin  Stream  Mechanically contained   
 Source: Natural  Man-enhanced  Man-made  Ephemeral  Permanent  Temporary   
 Date of construction/Period of flow: Flow gauge box has water directed to it through PVC pipes from two tunnels (adits) excavated by man. Unknown date of tunnel excavations.  
 Disturbance type (if any) and date: A cement retaining wall (3-4 "high) was constructed at the entrance to the east cave with a pipe for drainage to the outside. Date of construction unknown. Inundated: Yes  No   
 Depth of standing water 8-10cm in Caves; Saturated: Yes  No  Depth to saturation 18 inches  
 Other field indicators: A broken pipe from the east cave allows water to form a small shallow (1" deep) pool outside the cave entrance. Flow rate from both caves combined was estimated to be about 1.9 L/min on September 3, 1996.  
 Atypical situation: Yes  No ; Wetland hydrology: Yes  No   
 Basis: Surface water occurs at the observation point. Gravelly soils allow quick drainage of spring flow to subsurface levels of the soil stratum.

**Vegetation** List 3 dominant species, percent cover in bold, in each vegetation layer (5 if only 1 or 2 layers are present)

	Species	Indicator Status <sup>a</sup>	% Cover
<b>Trees</b>			
1.	<u>none</u>	<u></u>	<u></u>
2.	<u></u>	<u></u>	<u></u>
3.	<u></u>	<u></u>	<u></u>
4.	<u></u>	<u></u>	<u></u>
<b>Shrubs</b>			
1.	<u>Salix exigua</u>	<u>FACW</u>	<u>80</u>
2.	<u></u>	<u></u>	<u></u>
3.	<u></u>	<u></u>	<u></u>
4.	<u></u>	<u></u>	<u></u>
<b>Herbs</b>			
1.	<u>Potentilla biennis</u>	<u>FAC</u>	<u>Tr.</u>
2.	<u>Rumex salicifolius</u>	<u>FACW*</u>	<u>5</u>
3.	<u></u>	<u></u>	<u></u>
4.	<u></u>	<u></u>	<u></u>
5.	<u></u>	<u></u>	<u></u>
6.	<u></u>	<u></u>	<u></u>
7.	<u></u>	<u></u>	<u></u>
8.	<u></u>	<u></u>	<u></u>

Other field indicators:   
 Percentage of species that are OBL, FACW, and/or FAC: 100 %; Hydrophytic vegetation: Yes  No   
 Basis: A dominance of hydrophytic vegetation occurs at the observation point.

**Hydric Soils**

Field indicators: Hydric soils exist based on the presence of saturated soils for 7 days or longer duration at the observation point.

Hydric Soils: Yes  No

Jurisdictional Wetland Determination : Wetland  Nonwetland

**Notes:**

Salix exigua stand measures about 70m by 20-30 m across. Artemisia ludoviciana and Ericameria nauseosa are encroaching on the edge of the delineated area. Limited surface water: Salix shrubs ≈ 10 ft. high (3m) - Rhus trilobata is possibly spread by birds. Willow trees provide considerable cover for birds and wildlife. <sup>a</sup> Wetland indicator status for plants in region 8. FACW= Facultative wetland species. FAC = Facultative wetland species. \* refers to a tentative assignment to a category based on limited information in region 8. Tr = Trace amounts (<1% cover).



6.

10.

Percentage of species that are OBL, FACW, and/or FAC: 0%; Hydrophytic vegetation: Yes  No   
Basis: Hydrophytic vegetation was absent from the observation point.

**Hydric Soils**

Field indicators: Hydric soils appear to exist based on the presence of saturated soils for a period of 7 days or greater duration.

Hydric Soils: Yes  No

Jurisdictional Wetland Determination : Wetland  Nonwetland

**Notes:**

Water flows in three washes for about 40-60m length. Inundated pools exist with some shallow gravelly soils. <sup>a</sup> Wetlands indicator status for plants in region 8. NL = not listed in the National List of Plants that occur in Wetlands for Region 8. NI = insufficient information to classify plants according to wetlands status.

# Routine Jurisdictional Wetland Determination

Name: PDG, DJH Wetland Unit: Playa Pond  
 Location: Yucca Playa Pond UTM Coordinates Easting: 584-805 Northing: 40-90-584  
 Date: 1-7-97

## Hydrology

Type: Seep  Spring  Pond  Detention basin  Stream  Mechanically contained   
 Source: Natural  Man-enhanced  Man-made  Ephemeral  Permanent  Temporary   
 Date of construction/Period of flow: None\ unknown  
 Disturbance type (if any) and date: None Inundated: Yes  No   
 Depth of standing water >100cm; Saturated: Yes  No  Depth to saturation surface  
 Other field indicators: \_\_\_\_\_  
 Atypical situation: Yes  No ; Wetland hydrology: Yes  No   
 Basis: Surface water is present at the observation point.

## Vegetation *List 3 dominant species, percent cover in bold, in each vegetation layer (5 if only 1 or 2 layers are present)*

<i>Species</i>	<i>Indicator Status<sup>a</sup></i>	<i>% Cover</i>
<b>Trees</b>		
1. <u><i>Tamarix ramosissima</i></u>	<u>FACW</u>	<u>5</u>
2. _____	_____	_____
3. _____	_____	_____
4. _____	_____	_____
5. _____	_____	_____
<b>Shrubs</b>		
1. _____	_____	_____
2. _____	_____	_____
3. _____	_____	_____
4. _____	_____	_____
5. _____	_____	_____
<b>Herbs</b>		
1. <u><i>Typha domingensis</i></u>	<u>OBL</u>	<u>Tr.</u>
2. _____	_____	_____
3. _____	_____	_____
4. _____	_____	_____
5. _____	_____	_____
6. _____	_____	_____
7. _____	_____	_____
8. _____	_____	_____
9. _____	_____	_____
10. _____	_____	_____

Other field indicators: About 50 Tamarix trees exist at the water source.  
 Percentage of species that are OBL, FACW, and/or FAC: 100 %; Hydrophytic vegetation: Yes  No   
 Basis: A dominance of hydrophytic plants occurs at the observation point. The wetland area was estimated to be about 3400m<sup>2</sup>, including the area of plants and an equal area of rooting zone.

## Hydric Soils

Field indicators: Hydric soils exist based on the presence of saturated soils for a period of 7 days or greater duration.

Hydric Soils: Yes  No

Jurisdictional Wetland Determination : Wetland  Nonwetland

## Notes:

Soils at the observation site have high silt content. The pond was frozen over at the time of the survey. <sup>a</sup> Wetland indicator status for plants in region 8. Tr. = trace amounts (<1% absolute cover). FACW = Faculative wetland species. OBL = Obligate wetland species.

**This Page Intentionally Left Blank**

**APPENDIX C**

**LIST OF ALGAE AND ALGAE-LIKE SPECIES  
IDENTIFIED FROM SPRINGS ON THE NEVADA TEST SITE**

Appendix C. List of algae and algae-like species identified from springs on the NTS.

Scientific Name	Cane Spring	Captain Jack Spring	Reitmann Seep	Oak Spring	Tippah Spring	Topopah Spring	Tub Spring	Whiterock Spring
<b>Chrysophyta (Golden Algae, Diatoms)</b>								
<i>Achnanthes exigua</i>				X		X		X
<i>Achnanthes lanceolata</i>	X	X		X	X	X	X	X
<i>Achnanthes minutissima</i>	X							
<i>Achnanthes saxonica</i>		X		X				X
<i>Amphora submontana</i>								X
<i>Asterionella formosa</i>			X					
<i>Denticula elegans</i>								X
<i>Epithemia adnata</i> var. <i>proboscidea</i>					X			
<i>Epithemia sorex</i>							X	
<i>Fragilaria</i> sp.				X		X		
<i>Fragilaria construens</i>								X
<i>Gomphonema parvulum</i>	X	X		X	X	X		X
<i>Hantzschia</i> sp.								X
<i>Melosira granulata</i>			X					
<i>Meridian circulare</i>		X						
<i>Navicula cryptocephala</i>		X						
<i>Navicula cuspidata</i> var. <i>ambigua</i>	X							
<i>Navicula laevissima</i>				X				
<i>Navicula minima</i>	X			X		X		
<i>Navicula rhynchocephala</i> var. <i>amphiceras</i>							X	
<i>Nitzschia</i> sp.	X	X	X	X		X		X
<i>Nitzschia amphibia</i>								X
<i>Nitzschia gracilis</i>				X				
<i>Nitzschia linearis</i>	X			X				X
<i>Nitzschia palea</i>	X			X				
<i>Nitzschia tryblionella</i>								X
<i>Pinnularia</i> sp.								X
<i>Pinnularia abaujensis</i> var. <i>subundulata</i>						X		
<i>Pinnularia viridis</i> var. <i>minor</i>	X							
<i>Stauroneis anceps</i>		X						
<i>Stephanodiscus niagarae</i>			X					
<i>Surirella ovalis</i>	X					X		X
<i>Vaucheria</i> sp.	X							

Sources: Shields and Drouet, 1962; Taylor and Giles, 1979

Appendix C. List of algae and algae-like species identified from springs on the NTS.

Scientific Name	Cane Spring	Captain Jack Spring	Reitmann Seep	Oak Spring	Tippipah Spring	Topopah Spring	Tub Spring	Whiterock Spring
<b>Chlorophyta (Green Algae)</b>								
<i>Ankistrodesmus falcatus</i>			X					
<i>Bulbochaete</i> sp.	X							
<i>Chara</i> sp.	X							
<i>Chlamydomonas</i> sp.								X
<i>Chlorella vulgaris</i>		X	X					
<i>Closterium turgidum</i>						X		
<i>Cosmarium</i> sp.		X					X	
<i>Franceia droescheri</i>	X							
<i>Haematococcus lacustris</i>				X				
<i>Microthamnion kuetzingianum</i>	X			X				
<i>Oedogonium</i> sp.	X	X						X
<i>Oocystis borgei</i>				X				
<i>Oocystis crassa</i>	X							
<i>Pandorina morum</i>	X							
<i>Protoderma viride</i>				X				
<i>Scenedesmus acutus</i>				X				
<i>Scenedesmus bijuga</i>	X							
<i>Spirogyra juergensii</i>		X						
<i>Stigeoclonium</i> sp.	X	X		X			X	
<i>Ulothrix</i> sp.				X				
<b>Cyanobacteria (formerly known as blue-green algae)</b>								
<i>Amphithrix janthina</i>	X							
<i>Calothrix</i> sp.		X		X				
<i>Lyngbya</i> sp.								X
<i>Nodularia sphaerocarps</i>	X							
<i>Nostoc entophyllum</i>	X							
<i>Oscillatoria brevis</i>	X							
<i>Oscillatoria</i> sp.		X	X			X		
<i>Phormidium autumnale</i>	X							
<i>Phormidium</i> sp.								X
<i>Phormidium tenue</i>	X						X	
<i>Plectonema boryanum</i>	X							
<b>Total Number of Species</b>	<b>27</b>	<b>14</b>	<b>7</b>	<b>19</b>	<b>3</b>	<b>10</b>	<b>6</b>	<b>19</b>

Sources: Shields and Drouet, 1962; Taylor and Giles, 1979

**APPENDIX D**

**LIST OF ANIMALS DOCUMENTED TO OCCUR AT  
NTS WETLANDS AND THEIR SEASON OF USE**



Appendix D. List of animals documented to occur at NTS wetlands.

The season in which animals were observed are shown: F=fall, Sp=spring, S=summer, W=winter, Y=year round, X=time of year unspecified

Common Name	Scientific Name	Ammonia Tanks	Cane Spring	Captain Jack Spring	Cottonwood Spring	Coyote Spring	Fortymile Canyon Tanks	Gold Meadows Spring	John's Spring	Oak Spring	Pavits Spring	Rainier Spring	Reitmann Seep	Rock Valley Tank	Tipipah Spring	Tongue Wash Tank	Topopah Spring	Tub Spring	Tupapa Seep	Twin Spring	Wahmonie Seep 1	Wahmonie Seep 2	Wahmonie Seep 3	Whiterock Spring	Yellow Rock Springs	Yuca Playa Pond	Total Sites
Brewer's blackbird	<i>Euphagus cyanocephalus</i>		X					S							S												1
Brewer's sparrow	<i>Spizella breweri</i>		X					S							S									X			4
Broad-tailed hummingbird	<i>Selasphorus platycercus</i>		X					S Sp							S												1
Brown-headed cowbird	<i>Molothrus ater</i>		X					S							S												4
Bushiti	<i>Psaltriparus minimus</i>			S																							1
Cassin's finch	<i>Carpodacus cassinii</i>							S																			1
Cassin's kingbird	<i>Tyrannus vociferans</i>		X					S																			1
Cedar waxwing	<i>Bombicilla cedrorum</i>		Sp					S							X								Sp				2
Chipping sparrow	<i>Spizella passerina</i>		Sp					S																			3
Chukar	<i>Alcedo alaxandri</i>	X	W	SF	Sp			X						SF	X	Y	SF			X						X	10
Common raven	<i>Corvus corax</i>		SF	Y				SF		Sp				SF	S	W Sp			F				S Sp				9
Common snipe	<i>Gallinago gallinago</i>		Sp																								1
Common yellowthroat	<i>Geothlypis trichas</i>		X																				X				2
Cooper's hawk	<i>Accipiter cooperii</i>		X	F				X									W						F				5
Costa's hummingbird	<i>Calypte costae</i>		Sp						X					X													2
Dark-eyed junco	<i>Junco hyemalis</i>		X	FW									F														3
European starling	<i>Sturnus vulgaris</i>		SF	SF																							1
Gambel's quail	<i>Callipepla gambelli</i>		SF	SF				X			F				SF	S	S	F					SF				10
Golden eagle	<i>Aquila chrysaetos</i>		Sp	Sp											Sp								Sp				3
Gray catbird	<i>Dumetella carolinensis</i>		X				W																				1
Great blue heron	<i>Ardea herodias</i>																								X		1
Great egret	<i>Casmerodius albus</i>		X																								1
Great-horned owl	<i>Bubo virginianus</i>																								X		1
Greater roadrunner	<i>Geococcyx californianus</i>		Sp																						X		1
Green-tailed towhee	<i>Pipilo chlorurus</i>		Sp					S																			2
Green-winged teal	<i>Anas crecca</i>		X																						X		1
Hermit thrush	<i>Cathartes guttatus</i>		Sp																								1
House finch	<i>Carpodacus mexicanus</i>		Y					S				SW	Sp			F	S						S				7
Indigo bunting	<i>Passerina cyanea</i>							S																			1
Lark sparrow	<i>Chondestes grammacus</i>		X					S																			2
Lazuli bunting	<i>Passerina amoena</i>		S Sp																								2
Le Conte's thrasher	<i>Toxostoma lecontei</i>		Sp																								1
Lesser goldfinch	<i>Carduelis psaltria</i>		F					S																			2
Lesser yellowlegs	<i>Tringa flavipes</i>		X																								1
Lewis' woodpecker	<i>Melanerpes lewis</i>		X																								1
Lincoln's sparrow	<i>Melospiza lincolni</i>		X																								1

Sources: Allred et al., 1963; Hayward et al., 1963; Jorgensen and Haywood, 1965; Casstetter, 1979; Romney and Greger, 1992; Greger and Romney, 1994a,b; BN, 1996; and Bechtel Nevada unpublished field records, 1974-1997.

Appendix D. List of animals documented to occur at NTS wetlands.  
 The season in which animals were observed are shown: F=fall, Sp=spring, S=summer, W=winter, Y=year round, X=time of year unspecified

Common Name	Scientific Name	Amonia Tanks	Cane Spring	Captain Jack Spring	Cottonwood Spring	Coyote Spring	Forty-mile Canyon Tanks	Gold Meadows Spring	John's Spring	Oak Spring	Pavits Spring	Rainier Spring	Reitmann Seep	Rock Valley Tank	Tipipah Spring	Tongue Wash Tank	Topah Spring	Tub Spring	Tupapa Seep	Twin Spring	Wahmonie Seep 1	Wahmonie Seep 2	Wahmonie Seep 3	Whitrock Spring	Yellow Rock Springs	Yucca Playa Pond	Total Sites
Loggerhead shrike	<i>Lanius ludovicianus</i>		F W												X												2
Long-eared owl	<i>Asio otus</i>		Sp																					S W			2
MacGillivray's warbler	<i>Oporornis tolmiei</i>		Y												X												2
Mountain bluebird	<i>Sialia currucoides</i>		Sp					S Sp							X												2
Mourning doves	<i>Zenaidura macroura</i>		S Sp	S Sp				S Sp		S					S Sp			F	S					S Sp			9
Northern flicker	<i>Sialia sialis</i>		X					Sp F																			1
Northern goshawk	<i>Accipiter gentilis</i>		W F					S																			2
Northern harrier	<i>Circus cyaneus</i>							S					X														1
Northern mockingbird	<i>Mimus polyglottos</i>							S									W										3
Northern oriole	<i>Icterus galbula</i>		Sp	F				S							S		S	S						X			7
Olive-sided flycatcher	<i>Contopus borealis</i>		Sp																								1
Orange-crowned warbler	<i>Vermivora celata</i>		Sp	X																							1
Phainopepla	<i>Phainopepla nitens</i>		X	F																							2
Pine siskin	<i>Carolinensis pinus</i>		X	S W				S								X		Sp F									2
Pinyon jay	<i>Gymnorhinus cyanocephala</i>		X	F																							5
Prairie falcon	<i>Falco mexicanus</i>		Sp																								1
Red-naped sapsucker	<i>Sphyrapicus varius</i>		F																								1
Red-necked phalarope	<i>Phalaropus lobatus</i>		X																								1
Red-tailed hawk	<i>Buteo jamaicensis</i>		W	S Sp				S F							S F		X							X			6
Red-winged blackbird	<i>Agelaius phoeniceus</i>		X												S												1
Rock wren	<i>Salpinctes obsoletus</i>		X	S Sp											S												4
Rose-breasted grosbeak	<i>Phoenicurus phoeniceus</i>			Sp				X																			2
Ruby-crowned kinglet	<i>Regulus calendula</i>		F Sp	F																							2
Rufous-sided towhee	<i>Pipilo erythrophthalmus</i>		X																								1
Sage sparrow	<i>Amphispiza belli</i>		W												F												3
Say's phoebe	<i>Sayornis saya</i>		S																								4
Scott's oriole	<i>Icterus parisorum</i>		X	S					X	X														S			2
Scrub jay	<i>Aphelocoma coerulescens</i>			F											S F		F	S									6
Sharp-shinned hawk	<i>Accipiter striatus</i>		Sp	W				S																			3
Song sparrow	<i>Melospiza melodia</i>		X																								1
Swinson's thrush	<i>Cathartes ustulatus</i>		Sp																								2
Townsend's warbler	<i>Dendroica townsendi</i>		X																								1
Tree swallow	<i>Tachycineta bicolor</i>		X																								1
Turkey vulture	<i>Cathartes aura</i>		X	S Sp											S Sp									S			4
Vesper sparrow	<i>Pooecetes gramineus</i>		X					S																			2
Violet-green swallow	<i>Tachycineta thalassina</i>							S																			1
Warbling vireo	<i>Vireo gilvus</i>		Sp																								1

Sources: Allred et al., 1963; Hayward et al., 1963; Jorgensen and Hayward, 1965; Castetter, 1979; Romney and Greger, 1992; Greger and Romney, 1994a,b; BN, 1996; and Dechtel Nevada unpublished field records, 1974-1997.

Appendix D. List of animals documented to occur at NTS wetlands.

The season in which animals were observed are shown: F=fall, Sp=spring, S=summer, W=winter, Y=year round, X=time of year unspecified

Common Name	Scientific Name	Ammonia Tanks	Cane Spring	Captain Jack Spring	Cottonwood Spring	Coyote Spring	Fortymile Canyon Tanks	Gold Meadows Spring	John's Spring	Oak Spring	Favits Spring	Rainier Spring	Reitmann Seep	Rock Valley Tank	Tippipah Spring	Tongue Wash Tank	Topopah Spring	Tub Spring	Tupapa Seep	Twin Spring	Wahmonie Seep 1	Wahmonie Seep 2	Wahmonie Seep 3	Whiterock Spring	Yellow Rock Springs	Yucca Playa Pond	Total Sites	
Western bluebird	<i>Sialia mexicana</i>							S																				1
Western meadowlark	<i>Sturnella neglecta</i>		X																									1
Western pipstrelle	<i>Pipistrellus hesperus</i>		X																									1
Western tanager	<i>Piranga ludoviciana</i>		Sp																					Sp				2
Western wood-pewee	<i>Contopus sordidulus</i>		X																									1
White-crowned sparrow	<i>Zonotrichia leucophrys</i>		F W	F											F													3
White-throated swift	<i>Aeronautes saxatalis</i>			S																								1
Wilson's warbler	<i>Wilsonia pusilla</i>		Sp					S																Sp				3
Wood duck	<i>Aix sponsa</i>		X																									1
Yellow warbler	<i>Dendroica petechia</i>		Sp																					X				2
Yellow-headed blackbird	<i>Xanthocephalus xanthocephalus</i>		X																									1
Yellow-rumped warbler	<i>Dendroica coronata</i>		Sp																									1
<b>Reptiles</b>																												
Desert horned lizard	<i>Phrynosoma platyrhinos</i>		X																									1
Desert spiny lizard	<i>Sceloporus magister</i>		X																									1
Desert tortoise	<i>Gopherus agassizii</i>													X														1
Long-nosed snake	<i>Rhinocheilus lecontei</i>		X																									1
Side-blotched lizard	<i>Uta stansburiana</i>		X																									1
Sidewinder	<i>Crotalus cerastes</i>		X																									1
Western fence lizard	<i>Sceloporus occidentalis</i>			X																								1
Western whiptail	<i>Cnemidophorus tigris</i>		X																									1
Zebra-tailed lizard	<i>Callisaurus draconoides</i>		X																									1
<b>Invertebrates</b>																												
Ant	<i>Neivamyrmex minor</i>		X																									1
Black-horned tree cricket	<i>Oecanthus nigricornis</i>		X																									1
California tree cricket	<i>Oecanthus californicus</i>		X																									1
Desert long-horned grasshopper	<i>Tmaecerus koebelei</i>		X																									1
Field cricket	<i>Acheta assimilis</i>		X																									1
Ground mantid	<i>Litanentria minor</i>		X																									1
long-horned grasshopper	<i>Capnobotes fuliginosus</i>		X																									1
Mantid	<i>Stagnomantis californicus</i>		X																									1
Pallid-winged grasshopper	<i>Trimerotropis pallidipennis</i>		X																									1
Scarab beetle	<i>Paracotalpa granicollis</i>		X																									1
short-horned grasshopper	<i>Aeoloplides minor</i>		X																									1
short-horned grasshopper	<i>Aeoloplides tenuipennis</i>		X																									1
short-horned grasshopper	<i>Amphitornius coloradus</i>		X																									1
short-horned grasshopper	<i>Eremiacris pallida</i>		X																									1

Sources: Allred et al., 1963; Hayward et al., 1963; Jorgensen and Haywood, 1965; Castetter, 1979; Romney and Greger, 1992; Greger and Romney, 1994a,b; BN, 1996; and Bechtel Nevada unpublished field records, 1974-1997.

D-4



**This Page Intentionally Left Blank**

Kenneth Vogel

Death Valley National Park  
Death Valley, California  
Richard Martin

Nellis Air Force Base  
Las Vegas, Nevada  
Susan Barrow (2)

Nevada Department of Wildlife, Region III  
Las Vegas, Nevada  
Mike Wickersham

Ash Meadows National Wildlife Refuge  
Ash Meadows, Nevada  
David Ledig

~~Sandia National Laboratory~~  
Las Vegas, Nevada  
Larry Posey

The Nature Conservancy  
Las Vegas, Nevada  
Teri Knight

DOE Office of Scientific and Technical  
Information  
Alexandria, Virginia (2)

Technical Information Resource Center  
Las Vegas, Nevada

Bechtel Nevada Public Reading Room  
Las Vegas, Nevada