

Incidence Rates[†] for New Mexico, 2003 - 2007

**All Cancer Sites
All Races (includes Hispanic), Both Sexes, All Ages**

County 	Annual Incidence Rate over rate period cases per 100,000 (95% Confidence Interval) 	Average Cases per Year over rate period 	Rate Period	Interval Range	Interval Color
US (SEER + NPCR) ¹	464.5 (464.1 - 464.8)	§	2003 - 2007	N/A	N/A
New Mexico ³	403.6 (399.6 - 407.6)	7,866	2003 - 2007	N/A	N/A
Quay County ⁷	488.5 (434.9 - 547.5)	65	2003 - 2007	457.4 - 488.5	
Roosevelt County ⁷	465.6 (420.4 - 514.3)	79	2003 - 2007	457.4 - 488.5	
Sandoval County ⁷	444.7 (426.4 - 463.5)	457	2003 - 2007	426.2 - 457.3	
Chaves County ⁷	435.9 (413.7 - 459.0)	295	2003 - 2007	426.2 - 457.3	
Los Alamos County ⁷	433.4 (393.5 - 476.4)	95	2003 - 2007	426.2 - 457.3	
Sierra County ⁷	433.4 (392.7 - 477.9)	98	2003 - 2007	426.2 - 457.3	
Eddy County ⁷	431.9 (407.8 - 457.2)	243	2003 - 2007	426.2 - 457.3	
Grant County ⁷	422.6 (393.7 - 453.1)	169	2003 - 2007	394.9 - 426.1	
Union County ⁷	422.4 (347.0 - 511.1)	23	2003 - 2007	394.9 - 426.1	
Santa Fe County ⁷	417.2 (402.5 - 432.3)	634	2003 - 2007	394.9 - 426.1	
Lea County ⁷	415.9 (392.0 - 441.0)	226	2003 - 2007	394.9 - 426.1	
Valencia County ⁷	415.8 (393.9 - 438.6)	275	2003 - 2007	394.9 - 426.1	
Bernalillo County ⁷	415.5 (408.3 - 422.9)	2,512	2003 - 2007	394.9 - 426.1	
Curry County ⁷	413.5 (386.2 - 442.2)	172	2003 - 2007	394.9 - 426.1	
Guadalupe County ⁷	406.0 (331.1 - 493.4)	21	2003 - 2007	394.9 - 426.1	
Torrance County ⁷	401.9 (359.0 - 448.6)	66	2003 - 2007	394.9 - 426.1	
Taos County ⁷	399.5 (370.9 - 429.7)	151	2003 - 2007	394.9 - 426.1	
Dona Ana County ⁷	396.2 (383.0 - 409.7)	687	2003 - 2007	394.9 - 426.1	
Catron County ⁷	388.1 (317.9 - 473.7)	24	2003 - 2007	363.6 - 394.8	
Otero County ⁷	385.9 (364.8 - 407.8)	254	2003 - 2007	363.6 - 394.8	
Socorro County ⁷	382.6 (342.7 - 425.8)	69	2003 - 2007	363.6 - 394.8	
Luna County ⁷	379.4 (349.9 - 411.0)	131	2003 - 2007	363.6 - 394.8	
San Miguel County ⁷	377.0 (346.9 - 408.9)	119	2003 - 2007	363.6 - 394.8	
Lincoln County ⁷	363.6 (332.5 - 397.1)	107	2003 - 2007	363.6 - 394.8	
San Juan County ⁷	355.3 (339.2 - 372.0)	373	2003 - 2007	332.4 - 363.5	
Cibola County ⁷	350.6 (319.4 - 384.1)	94	2003 - 2007	332.4 - 363.5	
Rio Arriba County ⁷	336.9 (312.2 - 363.1)	140	2003 - 2007	332.4 - 363.5	
Colfax County ⁷	316.7 (280.8 - 356.3)	60	2003 - 2007	301.2 - 332.3	
McKinley County ⁷	316.2 (295.1 - 338.3)	175	2003 - 2007	301.2 - 332.3	
Hidalgo County ⁷	313.2 (253.4 - 383.9)	19	2003 - 2007	301.2 - 332.3	
De Baca County ⁷	312.6 (228.9 - 422.9)	11	2003 - 2007	301.2 - 332.3	
Harding County ⁷	311.3 (191.1 - 513.2)	4	2003 - 2007	301.2 - 332.3	
Mora County ⁷	301.2 (242.7 - 370.5)	19	2003 - 2007	301.2 - 332.3	

Notes:

Created by statecancerprofiles.cancer.gov on 02/04/2011 9:13 pm.

[State Cancer Registries](#) may provide more current or more local data.

Data presented on the State Cancer Profiles Web Site may differ from statistics reported by the State Cancer Registries ([for more information](#)).

¹ Incidence rates (cases per 100,000 population per year) are age-adjusted to the [2000 US standard population](#) (19 age groups: <1, 1-4, 5-9, ... , 80-84, 85+). Rates are for invasive cancer only (except for bladder which is invasive and in situ) or unless otherwise specified. Rates calculated using SEER*Stat. Population counts for denominators are based on Census populations as modified by NCI. The US populations included with the data release have been adjusted for the population shifts due to [hurricanes Katrina and Rita](#) for 62 counties and parishes in Alabama, Mississippi, Louisiana, and Texas The 1969-2007 US Population Data File is used with SEER November 2009 data. The 1969-2006 US Population Data File is used with NPCR data November 2008/January 2009 data.

² Source: CDC's National Program of Cancer Registries Cancer Surveillance System (NPCR-CSS) November 2009/January 2010 data submission and [SEER November 2009 submission](#).

³ Source: [SEER November 2009 submission](#). State Cancer Registry also receives funding from CDC's National Program of Cancer Registries.

⁷ Source: SEER November 2009 submission.

⁸ Data not provided because it did not meet USCS data quality standards for one or more years during the rate period of data collection. While 96% of the US population resided in geographic areas with population-based cancer registries meeting the registry eligibility criteria for 2004, 4% of the US population was not yet represented in the [United States Cancer Statistics](#). American Cancer Society's [Facts & Figures](#) provides estimates of numbers of new cancer cases and deaths.

[▲ return to top](#)

Age-Adjusted Death Rates for New Mexico, 2003 - 2007

All Cancer Sites

All Races (includes Hispanic), Both Sexes, All Ages

County ▲	Annual Death Rate over rate period deaths per 100,000 (95% Confidence Interval) ▼	Average Deaths per Year over rate period ▼	Rate Period
United States	183.8 (183.6 - 184.0)	558,564	2003 - 2007
New Mexico	162.2 (159.6 - 164.8)	3,132	2003 - 2007
Guadalupe County	223.3 (169.4 - 289.7)	12	2003 - 2007
Sierra County	206.0 (179.8 - 235.9)	51	2003 - 2007
Roosevelt County	202.3 (172.9 - 235.2)	34	2003 - 2007
De Baca County	184.3 (116.1 - 282.3)	6	2003 - 2007
Valencia County	183.7 (168.9 - 199.4)	116	2003 - 2007
Quay County	183.4 (152.8 - 219.2)	26	2003 - 2007
Torrance County	181.7 (152.5 - 214.8)	29	2003 - 2007
Chaves County	180.2 (166.3 - 195.0)	127	2003 - 2007
Eddy County	180.1 (164.8 - 196.5)	103	2003 - 2007
Curry County	176.9 (159.3 - 195.9)	75	2003 - 2007
Lea County	176.4 (160.9 - 192.9)	96	2003 - 2007
Otero County	174.5 (160.3 - 189.6)	113	2003 - 2007
Union County	165.7 (120.7 - 224.3)	9	2003 - 2007
Sandoval County	165.3 (154.2 - 177.1)	166	2003 - 2007

Catron County	164.6 (118.9 - 227.1)	10	2003 - 2007
Rio Arriba County	163.1 (145.8 - 181.8)	66	2003 - 2007
Socorro County	162.3 (136.4 - 191.6)	28	2003 - 2007
Bernalillo County	162.1 (157.5 - 166.8)	969	2003 - 2007
San Miguel County	161.8 (142.2 - 183.4)	50	2003 - 2007
Grant County	160.8 (143.7 - 179.6)	67	2003 - 2007
Hidalgo County	158.7 (117.4 - 210.9)	10	2003 - 2007
San Juan County	158.7 (147.7 - 170.2)	159	2003 - 2007
Luna County	157.0 (138.7 - 177.2)	57	2003 - 2007
Dona Ana County	156.3 (148.0 - 164.9)	269	2003 - 2007
Taos County	153.8 (136.1 - 173.2)	56	2003 - 2007
Cibola County	152.9 (132.5 - 175.5)	41	2003 - 2007
Santa Fe County	148.3 (139.4 - 157.6)	213	2003 - 2007
McKinley County	145.7 (131.0 - 161.5)	75	2003 - 2007
Lincoln County	142.6 (123.9 - 163.6)	43	2003 - 2007
Colfax County	129.2 (107.1 - 155.0)	25	2003 - 2007
Los Alamos County	127.4 (105.1 - 153.2)	24	2003 - 2007
Mora County	121.7 (87.2 - 166.8)	8	2003 - 2007
Harding County	*	3 or fewer	2003 - 2007

Notes:

Created by statecancerprofiles.cancer.gov on 02/04/2011 9:16 pm.

[State Cancer Registries](#) may provide more current or more local data.

Data presented on the State Cancer Profiles Web Site may differ from statistics reported by the State Cancer Registries ([for more information](#)).

Source: Death data provided by the [National Vital Statistics System](#) public use data file. Death rates calculated by the National Cancer Institute using [SEER*Stat](#) . Death rates (per 100,000 population per year) are age-adjusted to the [2000 US standard population](#) (19 age groups: <1, 1-4, 5-9, ... , 80-84, 85+). The Healthy People 2010 goals are based on different methods but the differences should be minimal. Population counts for denominators are based on the Census 1969-2006 US Population Data File as modified by NCI. The data release have been adjusted for the population shifts due to hurricanes [Katrina and Rita](#) for 62 counties and parishes in Alabama, Mississippi, Louisiana, and Texas. * Data have been [suppressed](#) to ensure confidentiality and stability of rate estimates. Counts are suppressed if fewer than 16 cases were reported in a specific area-sex-year combination. Healthy People 2010 Goal 03-01 : Reduce the overall cancer death rate to 159.9.

[Healthy People 2010](#) Objectives provided by the [Centers for Disease Control and Prevention](#) .

<http://statecancerprofiles.cancer.gov/map/scpMapDataTable.php?35&001&001&00&0&1&0&2&6&2>

Incidence Rates[†] for New Mexico, 2003 - 2007

**Thyroid
All Races (includes Hispanic), Both Sexes, All Ages**

County △	Annual Incidence Rate over rate period cases per 100,000 (95% Confidence Interval) ▼	Average Cases per Year over rate period ▽	Rate Period	Interval Range
New Mexico ³	12.2 (11.5 - 12.9)	234	2003 - 2007	N/A
US (SEER + NPCR) ¹	10.2 (10.2 - 10.3)	§	2003 - 2007	N/A
Los Alamos County ⁷	33.6 (22.1 - 48.7)	6	2003 - 2007	15.3 - 33.6
Grant County ⁷	15.8 (9.6 - 24.4)	4	2003 - 2007	15.3 - 33.6
Bernalillo County ⁷	15.2 (13.9 - 16.7)	94	2003 - 2007	13.6 - 15.2
Sandoval County ⁷	14.0 (11.0 - 17.5)	15	2003 - 2007	13.6 - 15.2
Santa Fe County ⁷	13.6 (11.1 - 16.6)	21	2003 - 2007	13.6 - 15.2
Rio Arriba County ⁷	13.5 (8.9 - 19.6)	6	2003 - 2007	12.7 - 13.5
Valencia County ⁷	13.1 (9.5 - 17.7)	9	2003 - 2007	12.7 - 13.5
Lea County ⁷	13.0 (9.0 - 18.1)	7	2003 - 2007	12.7 - 13.5
Eddy County ⁷	12.6 (8.6 - 17.9)	7	2003 - 2007	9.9 - 12.6
Curry County ⁷	11.9 (7.7 - 17.5)	5	2003 - 2007	9.9 - 12.6
San Miguel County ⁷	10.3 (5.9 - 16.9)	3	2003 - 2007	9.9 - 12.6
McKinley County ⁷	9.8 (6.5 - 14.3)	6	2003 - 2007	7.0 - 9.8
Chaves County ⁷	8.6 (5.5 - 12.8)	5	2003 - 2007	7.0 - 9.8
Dona Ana County ⁷	6.9 (5.3 - 8.9)	12	2003 - 2007	6.6 - 6.9
Otero County ⁷	6.9 (4.3 - 10.6)	4	2003 - 2007	6.6 - 6.9
San Juan County ⁷	6.6 (4.6 - 9.1)	7	2003 - 2007	6.6 - 6.9
Catron County ⁷	*	3 or fewer	2003 - 2007	*
Cibola County ⁷	*	3 or fewer	2003 - 2007	*
Colfax County ⁷	*	3 or fewer	2003 - 2007	*
De Baca County ⁷	*	3 or fewer	2003 - 2007	*
Guadalupe County ⁷	*	3 or fewer	2003 - 2007	*
Harding County ⁷	*	3 or fewer	2003 - 2007	*
Hidalgo County ⁷	*	3 or fewer	2003 - 2007	*
Lincoln County ⁷	*	3 or fewer	2003 - 2007	*
Luna County ⁷	*	3 or fewer	2003 - 2007	*
Mora County ⁷	*	3 or fewer	2003 - 2007	*
Quay County ⁷	*	3 or fewer	2003 - 2007	*
Roosevelt County ⁷	*	3 or fewer	2003 - 2007	*
Sierra County ⁷	*	3 or fewer	2003 - 2007	*
Socorro County ⁷	*	3 or fewer	2003 - 2007	*
Taos County ⁷	*	3 or fewer	2003 - 2007	*
Torrance County ⁷	*	3 or fewer	2003 - 2007	*

Union County ⁷	*	3 or fewer	2003 - 2007	*
---------------------------	---	------------	-------------	---

Notes:

Created by statecancerprofiles.cancer.gov on 02/04/2011 10:14 pm.

[State Cancer Registries](#) may provide more current or more local data.

Data presented on the State Cancer Profiles Web Site may differ from statistics reported by the State Cancer Registries ([for more information](#)).

[†] Incidence rates (cases per 100,000 population per year) are age-adjusted to the [2000 US standard population](#) (19 age groups: <1, 1-4, 5-9, ..., 80-84, 85+). Rates are (except for bladder which is invasive and in situ) or unless otherwise specified. Rates calculated using SEER*Stat. Population counts for denominators are based on Census data modified by NCI. The US populations included with the data release have been adjusted for the population shifts due to [hurricanes Katrina and Rita](#) for 62 counties and parts of Mississippi, Louisiana, and Texas. The 1969-2007 US Population Data File is used with SEER November 2009 data. The 1969-2006 US Population Data File is used with SEER 2008/January 2009 data.

¹ Source: CDC's National Program of Cancer Registries Cancer Surveillance System (NPCR-CSS) November 2009/January 2010 data submission and [SEER November 2009](#).

³ Source: [SEER November 2009 submission](#). State Cancer Registry also receives funding from CDC's National Program of Cancer Registries.

⁷ Source: SEER November 2009 submission.

[†] Data have been [suppressed](#) to ensure confidentiality and stability of rate estimates. Counts are suppressed if fewer than 16 cases were reported in a specific area-sex-age group.

[§] Data not provided because it did not meet USCS data quality standards for one or more years during the rate period of data collection. While 96% of the US population is covered by population-based cancer registries meeting the registry eligibility criteria for 2004, 4% of the US population was not yet represented in the [United States Cancer Statistics](#). The [United States Cancer Society's Facts & Figures](#) provides estimates of numbers of new cancer cases and deaths.

[▲ return to top](#)

Incidence Rates [†] for New Mexico, 2003 - 2007					
Breast					
All Races (includes Hispanic), Female, All Ages					
County	Annual Incidence Rate over rate period cases per 100,000 (95% Confidence Interval)	Average Cases per Year over rate period	Rate Period	Interval Range	Inte
US (SEER + NPCR) ¹	120.6 (120.4 - 120.9)	§	2003 - 2007	N/A	
New Mexico ³	108.5 (105.7 - 111.4)	1,127	2003 - 2007	N/A	
Los Alamos County ⁷	133.5 (104.3 - 169.0)	16	2003 - 2007	119.7 - 133.5	
Santa Fe County ⁷	131.7 (120.8 - 143.4)	112	2003 - 2007	119.7 - 133.5	
Sandoval County ⁷	131.1 (118.1 - 145.2)	75	2003 - 2007	119.7 - 133.5	
Grant County ⁷	120.4 (99.2 - 145.2)	24	2003 - 2007	119.7 - 133.5	
Torrance County ⁷	119.6 (88.6 - 158.1)	10	2003 - 2007	112.6 - 119.6	
Taos County ⁷	117.7 (97.1 - 141.7)	24	2003 - 2007	112.6 - 119.6	
Union County ⁷	117.2 (67.9 - 194.9)	3	2003 - 2007	112.6 - 119.6	
Bernalillo County ⁷	114.4 (109.2 - 119.7)	374	2003 - 2007	112.6 - 119.6	
Quay County ⁷	112.5 (78.2 - 158.5)	8	2003 - 2007	103.1 - 112.5	
Chaves County ⁷	111.5 (96.1 - 128.7)	39	2003 - 2007	103.1 - 112.5	
Dona Ana County ⁷	108.7 (99.2 - 118.7)	98	2003 - 2007	103.1 - 112.5	
Curry	108.2 (89.6 - 129.5)	24	2003 - 2007	103.1 - 112.5	

County ⁶					
San Miguel County ⁷	103.7 (82.4 - 128.9)	17	2003 - 2007	103.1 - 112.5	
Valencia County ⁷	103.0 (88.5 - 119.3)	36	2003 - 2007	92.6 - 103.0	
Otero County ⁷	102.6 (87.8 - 119.2)	35	2003 - 2007	92.6 - 103.0	
Eddy County ⁷	100.7 (84.9 - 118.5)	30	2003 - 2007	92.6 - 103.0	
Roosevelt County ⁷	98.9 (71.3 - 133.5)	9	2003 - 2007	92.6 - 103.0	
San Juan County ⁷	98.9 (87.7 - 111.1)	57	2003 - 2007	92.6 - 103.0	
Luna County ⁷	92.5 (71.5 - 117.7)	15	2003 - 2007	79.7 - 92.5	
Sierra County ⁷	91.1 (64.0 - 127.5)	9	2003 - 2007	79.7 - 92.5	
Socorro County ⁷	87.5 (62.2 - 119.7)	8	2003 - 2007	79.7 - 92.5	
Lea County ⁷	85.1 (70.5 - 101.8)	24	2003 - 2007	79.7 - 92.5	
Cibola County ⁷	83.6 (63.4 - 108.3)	12	2003 - 2007	79.7 - 92.5	
Rio Arriba County ⁷	79.6 (63.8 - 98.3)	18	2003 - 2007	71.4 - 79.6	
Lincoln County ⁷	78.2 (58.5 - 103.1)	12	2003 - 2007	71.4 - 79.6	
Colfax County ⁷	76.2 (52.4 - 108.2)	7	2003 - 2007	71.4 - 79.6	
McKinley County ⁷	71.4 (58.6 - 86.0)	23	2003 - 2007	71.4 - 79.6	
Catron County ⁷	*	3 or fewer	2003 - 2007	*	
De Baca County ⁷	*	3 or fewer	2003 - 2007	*	
Guadalupe County ⁷	*	3 or fewer	2003 - 2007	*	
Harding County ⁷	*	3 or fewer	2003 - 2007	*	
Hidalgo County ⁷	*	3 or fewer	2003 - 2007	*	
Mora County ⁷	*	3 or fewer	2003 - 2007	*	

Notes:

Created by statecancerprofiles.cancer.gov on 02/04/2011 10:21 pm.

[State Cancer Registries](#) may provide more current or more local data.

Data presented on the State Cancer Profiles Web Site may differ from statistics reported by the State Cancer Registries ([for more information](#)).

[†] Incidence rates (cases per 100,000 population per year) are age-adjusted to the [2000 US standard population](#) (19 age groups: <1, 1-4, 5-9, ..., 80-84, 85+). For cancer only (except for bladder which is invasive and in situ) or unless otherwise specified. Rates calculated using SEER*Stat. Population counts for denominators are based on 2000 Census populations as modified by NCI. The US populations included with the data release have been adjusted for the population shifts due to [hurricanes Katrina](#) and [Rita](#) in Louisiana and [Hurricane Ike](#) in Texas. The 1969-2007 US Population Data File is used with SEER November 2009 data. The 1969-2007 US Population Data File is used with NPCR data November 2008/January 2009 data.

[‡] Source: CDC's National Program of Cancer Registries Cancer Surveillance System (NPCR-CSS) November 2009/January 2010 data submission and [SEER data submission](#).

³ Source: [SEER November 2009 submission](#). State Cancer Registry also receives funding from CDC's National Program of Cancer Registries.

⁷ Source: SEER November 2009 submission.

⁷ Data have been [suppressed](#) to ensure confidentiality and stability of rate estimates. Counts are suppressed if fewer than 16 cases were reported in a specific

⁸ Data not provided because it did not meet USCS data quality standards for one or more years during the rate period of data collection. While 96% of the US population was represented in the geographic areas with population-based cancer registries meeting the registry eligibility criteria for 2004, 4% of the US population was not yet represented in the [Statistics](#). American Cancer Society's [Facts & Figures](#) provides estimates of numbers of new cancer cases and deaths.

Incidence Rates† for New Mexico, 2003 - 2007

Prostate All Races (includes Hispanic), Male, All Ages

County △	Annual Incidence Rate over rate period cases per 100,000 (95% Confidence Interval) ▼	Average Cases per Year over rate period ▽	Rate Period	Interval Range	Interval Color
US (SEER + NPCR) ¹	153.5 (153.2 - 153.8)	§	2003 - 2007	N/A	N/A
New Mexico ³	143.3 (139.8 - 146.8)	1,302	2003 - 2007	N/A	N/A
Los Alamos County ⁷	219.3 (181.0 - 264.0)	25	2003 - 2007	160.4 - 219.3	
Taos County ⁷	179.7 (152.4 - 210.7)	32	2003 - 2007	160.4 - 219.3	
Santa Fe County ⁷	169.8 (156.2 - 184.2)	122	2003 - 2007	160.4 - 219.3	
Catron County ⁷	161.1 (107.2 - 249.2)	6	2003 - 2007	160.4 - 219.3	
Roosevelt County ⁷	161.1 (122.8 - 207.3)	12	2003 - 2007	160.4 - 219.3	
Eddy County ⁷	160.3 (139.2 - 183.7)	42	2003 - 2007	145.3 - 160.3	
Sandoval County ⁷	158.4 (142.3 - 175.8)	73	2003 - 2007	145.3 - 160.3	
Dona Ana County ⁷	150.5 (138.8 - 162.9)	123	2003 - 2007	145.3 - 160.3	
Bernalillo County ⁷	149.0 (142.5 - 155.6)	411	2003 - 2007	145.3 - 160.3	
Valencia County ⁷	146.5 (128.2 - 166.7)	48	2003 - 2007	145.3 - 160.3	
Rio Arriba County ⁷	145.2 (121.8 - 171.8)	28	2003 - 2007	133.0 - 145.2	
Luna County ⁷	141.9 (118.3 - 169.4)	26	2003 - 2007	133.0 - 145.2	
Chaves County ⁷	136.3 (118.5 - 156.0)	42	2003 - 2007	133.0 - 145.2	
Curry County ⁷	134.4 (111.3 - 160.6)	24	2003 - 2007	133.0 - 145.2	
Socorro County ⁷	134.4 (101.4 - 174.7)	12	2003 - 2007	133.0 - 145.2	
Torrance County ⁷	132.9 (97.7 - 176.3)	11	2003 - 2007	126.0 - 132.9	
Sierra County ⁷	131.8 (103.9 - 167.4)	16	2003 - 2007	126.0 - 132.9	
Grant County ⁷	131.5 (110.2 - 156.3)	27	2003 - 2007	126.0 - 132.9	
Cibola County ⁷	130.2 (103.7 - 161.5)	17	2003 - 2007	126.0 - 132.9	
Lincoln County ⁷	128.0 (103.3 - 157.6)	19	2003 - 2007	126.0 - 132.9	
Union County ⁷	125.9 (71.9 - 209.2)	3	2003 - 2007	114.8 - 125.9	
San Miguel County ⁷	125.8 (101.2 - 154.6)	19	2003 - 2007	114.8 - 125.9	
Hidalgo County ⁷	120.2 (70.9 - 192.9)	4	2003 - 2007	114.8 - 125.9	
Mora County ⁷	116.8 (71.1 - 184.5)	4	2003 - 2007	114.8 - 125.9	
Quay County ⁷	115.8 (81.9 - 161.3)	8	2003 - 2007	114.8 - 125.9	
Lea County ⁷	114.7 (96.3 - 135.4)	28	2003 - 2007	101.2 - 114.7	

Otero County ⁷	108.7 (92.9 - 126.3)	35	2003 - 2007	101.2 - 114.7	
Colfax County ⁷	105.2 (78.4 - 139.5)	10	2003 - 2007	101.2 - 114.7	
McKinley County ⁷	101.5 (83.1 - 122.6)	22	2003 - 2007	101.2 - 114.7	
San Juan County ⁷	101.2 (88.5 - 115.2)	48	2003 - 2007	101.2 - 114.7	
De Baca County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Guadalupe County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Harding County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷

Notes:

Created by statecancerprofiles.cancer.gov on 02/04/2011 10:23 pm.

[State Cancer Registries](#) may provide more current or more local data.

Data presented on the State Cancer Profiles Web Site may differ from statistics reported by the State Cancer Registries ([for more information](#)).

¹ Incidence rates (cases per 100,000 population per year) are age-adjusted to the [2000 US standard population](#) (19 age groups: <1, 1-4, 5-9, ... , 80-84, 85+). Rates are for invasive cancer only (except for bladder which is invasive and in situ) or unless otherwise specified. Rates calculated using SEER*Stat. Population counts for denominators are based on Census populations as modified by NCI. The US populations included with the data release have been adjusted for the population shifts due to [hurricanes Katrina and Rita](#) for 62 counties and parishes in Alabama, Mississippi, Louisiana, and Texas. The 1969-2007 US Population Data File is used with SEER November 2009 data. The 1969-2006 US Population Data File is used with NPCR data November 2008/January 2009 data.

² Source: CDC's National Program of Cancer Registries Cancer Surveillance System (NPCR-CSS) November 2009/January 2010 data submission and [SEER November 2009 submission](#).

³ Source: [SEER November 2009 submission](#). State Cancer Registry also receives funding from CDC's National Program of Cancer Registries.

⁴ Source: SEER November 2009 submission.

⁵ Data have been [suppressed](#) to ensure confidentiality and stability of rate estimates. Counts are suppressed if fewer than 16 cases were reported in a specific area-sex-race category.

⁶ Data not provided because it did not meet USCS data quality standards for one or more years during the rate period of data collection. While 96% of the US population resided in geographic areas with population-based cancer registries meeting the registry eligibility criteria for 2004, 4% of the US population was not yet represented in the [United States Cancer Statistics](#). American Cancer Society's [Facts & Figures](#) provides estimates of numbers of new cancer cases and deaths.

Incidence Rates† for New Mexico, 2003 - 2007

Melanoma of the Skin

All Races (includes Hispanic), Male, All Ages

County 	Annual Incidence Rate over rate period cases per 100,000 (95% Confidence Interval) 	Average Cases per Year over rate period 	Rate Period	Interval Range	Interval Color
US (SEER + NPCR) ¹	23.1 (23.0 - 23.2)	§	2003 - 2007	N/A	N/A
New Mexico ³	21.1 (19.7 - 22.5)	189	2003 - 2007	N/A	N/A
Los Alamos County ⁷	38.2 (22.5 - 61.0)	4	2003 - 2007	27.4 - 38.2	
Luna County ⁷	28.1 (17.0 - 44.0)	4	2003 - 2007	27.4 - 38.2	
Chaves County ⁷	27.3 (19.7 - 37.0)	8	2003 - 2007	25.0 - 27.3	
Bernalillo County ⁷	25.5 (22.8 - 28.3)	70	2003 - 2007	25.0 - 27.3	
Sandoval County ⁷	24.9 (18.9 - 32.2)	12	2003 - 2007	22.3 - 24.9	
Valencia County ⁷	23.3 (16.0 - 32.7)	7	2003 - 2007	22.3 - 24.9	
Santa Fe County ⁷	23.0 (18.2 - 28.7)	17	2003 - 2007	22.3 - 24.9	
Lincoln County ⁷	22.2 (12.8 - 37.1)	3	2003 - 2007	20.4 - 22.2	
Grant County ⁷	21.6 (12.9 - 34.3)	4	2003 - 2007	20.4 - 22.2	
Otero County ⁷	21.5 (14.7 - 30.3)	7	2003 - 2007	20.4 - 22.2	
Lea County ⁷	20.3 (13.0 - 29.9)	5	2003 - 2007	17.8 - 20.3	
Dona Ana County ⁷	19.5 (15.4 - 24.3)	16	2003 - 2007	17.8 - 20.3	
Eddy County ⁷	17.7 (11.3 - 26.5)	5	2003 - 2007	12.3 - 17.7	
San Juan County ⁷	12.3 (8.3 - 17.5)	6	2003 - 2007	12.3 - 17.7	

Catron County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁶
Cibola County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁶
Colfax County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁶
Curry County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁶
De Baca County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁶
Guadalupe County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁶
Harding County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁶
Hidalgo County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁶
McKinley County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁶
Mora County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁶
Quay County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁶
Rio Arriba County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁶
Roosevelt County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁶
San Miguel County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁶
Sierra County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁶
Socorro County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁶
Taos County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁶
Torrance County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁶
Union County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁶

Notes:

Created by statecancerprofiles.cancer.gov on 02/04/2011 10:25 pm.

[State Cancer Registries](#) may provide more current or more local data.

Data presented on the State Cancer Profiles Web Site may differ from statistics reported by the State Cancer Registries ([for more information](#)).

¹ Incidence rates (cases per 100,000 population per year) are age-adjusted to the [2000 US standard population](#) (19 age groups: <1, 1-4, 5-9, ... , 80-84, 85+). Rates are for invasive cancer only (except for bladder which is invasive and in situ) or unless otherwise specified. Rates calculated using SEER*Stat. Population counts for denominators are based on Census populations as modified by NCI. The US populations included with the data release have been adjusted for the population shifts due to [hurricanes Katrina and Rita](#) for 62 counties and parishes in Alabama, Mississippi, Louisiana, and Texas. The 1969-2007 US Population Data File is used with SEER November 2009 data. The 1969-2006 US Population Data File is used with NPCR data November 2008/January 2009 data.

¹ Source: CDC's National Program of Cancer Registries Cancer Surveillance System (NPCR-CSS) November 2009/January 2010 data submission and [SEER November 2009 submission](#).

³ Source: [SEER November 2009 submission](#). State Cancer Registry also receives funding from CDC's National Program of Cancer Registries.

⁷ Source: SEER November 2009 submission.

⁶ Data have been [suppressed](#) to ensure confidentiality and stability of rate estimates. Counts are suppressed if fewer than 16 cases were reported in a specific area-sex-race category.

⁹ Data not provided because it did not meet USCS data quality standards for one or more years during the rate period of data collection. While 96% of the US population resided in geographic areas with population-based cancer registries meeting the registry eligibility criteria for 2004, 4% of the US population was not yet represented in the [United States Cancer Statistics](#). American Cancer Society's [Facts & Figures](#) provides estimates of numbers of new cancer cases and deaths.

Incidence Rates† for New Mexico, 2003 - 2007

Colon & Rectum

All Races (includes Hispanic), Male, All Ages

County ▲	Annual Incidence Rate over rate period cases per 100,000 (95% Confidence Interval) ▼	Average Cases per Year over rate period ▽	Rate Period	Interval Range	Interval Color
US (SEER + NPCR) ¹	57.0 (56.9 - 57.2)	§	2003 - 2007	N/A	N/A
New Mexico ³	48.0 (45.9 - 50.1)	425	2003 - 2007	N/A	N/A
Quay County ⁷	90.6 (60.4 - 132.6)	6	2003 - 2007	59.9 - 90.6	
Eddy County ⁷	67.1 (53.7 - 83.0)	17	2003 - 2007	59.9 - 90.6	
Sierra County ⁷	63.0 (43.8 - 90.4)	7	2003 - 2007	59.9 - 90.6	
Rio Arriba County ⁷	61.5 (46.5 - 79.7)	12	2003 - 2007	59.9 - 90.6	
Grant County ⁷	59.8 (44.9 - 78.4)	11	2003 - 2007	54.3 - 59.8	

Roosevelt County ¹	58.5 (36.9 - 87.8)	5	2003 - 2007	54.3 - 59.8	
Socorro County ¹	58.2 (37.7 - 86.0)	5	2003 - 2007	54.3 - 59.8	
San Miguel County ¹	57.2 (41.2 - 77.5)	9	2003 - 2007	54.3 - 59.8	
Taos County ¹	54.2 (39.8 - 72.4)	10	2003 - 2007	49.4 - 54.2	
Lea County ¹	53.1 (41.0 - 67.6)	13	2003 - 2007	49.4 - 54.2	
McKinley County ¹	49.8 (37.9 - 64.0)	13	2003 - 2007	49.4 - 54.2	
Sandoval County ¹	49.5 (40.6 - 59.6)	23	2003 - 2007	49.4 - 54.2	
Curry County ¹	49.3 (35.8 - 66.1)	9	2003 - 2007	44.1 - 49.3	
Otero County ¹	47.3 (36.9 - 59.8)	15	2003 - 2007	44.1 - 49.3	
Dona Ana County ¹	46.0 (39.6 - 53.1)	37	2003 - 2007	44.1 - 49.3	
Bernalillo County ¹	45.7 (42.0 - 49.5)	120	2003 - 2007	44.1 - 49.3	
Santa Fe County ¹	44.9 (37.8 - 53.0)	30	2003 - 2007	44.1 - 49.3	
Lincoln County ¹	44.0 (29.6 - 63.8)	6	2003 - 2007	38.8 - 44.0	
Chaves County ¹	43.6 (33.7 - 55.6)	13	2003 - 2007	38.8 - 44.0	
Valencia County ¹	43.1 (33.2 - 55.0)	13	2003 - 2007	38.8 - 44.0	
San Juan County ¹	42.4 (34.4 - 51.6)	21	2003 - 2007	38.8 - 44.0	
Colfax County ¹	38.7 (22.9 - 62.8)	4	2003 - 2007	34.6 - 38.7	
Los Alamos County ¹	37.8 (22.8 - 59.8)	4	2003 - 2007	34.6 - 38.7	
Cibola County ¹	37.3 (23.8 - 55.7)	5	2003 - 2007	34.6 - 38.7	
Luna County ¹	34.6 (23.1 - 50.4)	6	2003 - 2007	34.6 - 38.7	
Catron County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
De Baca County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Guadalupe County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Harding County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Hidalgo County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Mora County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Torrance County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Union County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ⁷

Notes:

Created by statecancerprofiles.cancer.gov on 02/04/2011 10:27 pm.

[State Cancer Registries](#) may provide more current or more local data.

Data presented on the State Cancer Profiles Web Site may differ from statistics reported by the State Cancer Registries ([for more information](#)).

¹ Incidence rates (cases per 100,000 population per year) are age-adjusted to the [2000 US standard population](#) (19 age groups: <1, 1-4, 5-9, ... , 80-84, 85+). Rates are for invasive cancer only (except for bladder which is invasive and in situ) or unless otherwise specified. Rates calculated using SEER*Stat. Population counts for denominators are based on Census populations as modified by NCI. The US populations included with the data release have been adjusted for the population shifts due to [hurricanes Katrina and Rita](#) for 62 counties and parishes in Alabama, Mississippi, Louisiana, and Texas. The 1969-2007 US Population Data File is used with SEER November 2009 data. The 1969-2006 US Population Data File is used with NPCR data November 2008/January 2009 data.

¹ Source: CDC's National Program of Cancer Registries Cancer Surveillance System (NPCR-CSS) November 2009/January 2010 data submission and [SEER November 2009 submission](#).

³ Source: [SEER November 2009 submission](#). State Cancer Registry also receives funding from CDC's National Program of Cancer Registries.

⁷ Source: SEER November 2009 submission.

* Data have been [suppressed](#) to ensure confidentiality and stability of rate estimates. Counts are suppressed if fewer than 16 cases were reported in a specific area-sex-race category.

⁵ Data not provided because it did not meet USCS data quality standards for one or more years during the rate period of data collection. While 96% of the US population resided in geographic areas with population-based cancer registries meeting the registry eligibility criteria for 2004, 4% of the US population was not yet represented in the [United States Cancer Statistics](#). American Cancer Society's [Facts & Figures](#) provides estimates of numbers of new cancer cases and deaths.

Incidence Rates† for New Mexico, 2003 - 2007

Lung & Bronchus

All Races (includes Hispanic), Male, All Ages

County △	Annual Incidence Rate over rate period cases per 100,000 (95% Confidence Interval)	Average Cases per Year over rate period ▽	Rate Period	Interval Range	Interval Color
-------------	---	--	-------------	----------------	----------------

US (SEER + NPCR) ¹	84.9 (84.7 - 85.1)	§	2003 - 2007	N/A	N/A
New Mexico ³	55.5 (53.3 - 57.8)	485	2003 - 2007	N/A	N/A
Roosevelt County ⁷	106.6 (75.7 - 145.5)	8	2003 - 2007	82.8 - 106.6	
Quay County ⁷	106.5 (74.0 - 150.6)	7	2003 - 2007	82.8 - 106.6	
Sierra County ⁷	87.1 (64.4 - 117.7)	10	2003 - 2007	82.8 - 106.6	
Lea County ⁷	82.8 (67.3 - 100.6)	20	2003 - 2007	82.8 - 106.6	
Chaves County ⁷	82.7 (68.9 - 98.4)	25	2003 - 2007	71.9 - 82.7	
Valencia County ⁷	75.4 (61.9 - 91.0)	23	2003 - 2007	71.9 - 82.7	
Otero County ⁷	75.1 (62.2 - 90.1)	24	2003 - 2007	71.9 - 82.7	
Curry County ⁷	74.5 (57.5 - 94.7)	13	2003 - 2007	71.9 - 82.7	
Eddy County ⁷	71.8 (58.0 - 88.0)	19	2003 - 2007	52.1 - 71.8	
Luna County ⁷	64.7 (49.1 - 84.4)	12	2003 - 2007	52.1 - 71.8	
San Juan County ⁷	62.8 (52.5 - 74.3)	28	2003 - 2007	52.1 - 71.8	
Cibola County ⁷	53.2 (36.2 - 75.2)	6	2003 - 2007	52.1 - 71.8	
Bernalillo County ⁷	52.0 (48.1 - 56.1)	134	2003 - 2007	48.0 - 52.0	
Dona Ana County ⁷	51.1 (44.4 - 58.6)	41	2003 - 2007	48.0 - 52.0	
Grant County ⁷	50.3 (37.3 - 67.0)	10	2003 - 2007	48.0 - 52.0	
Sandoval County ⁷	49.7 (40.7 - 60.0)	22	2003 - 2007	48.0 - 52.0	
San Miguel County ⁷	49.0 (34.0 - 68.4)	7	2003 - 2007	48.0 - 52.0	
Lincoln County ⁷	47.9 (33.4 - 67.6)	7	2003 - 2007	40.4 - 47.9	
Torrance County ⁷	47.1 (28.6 - 73.5)	4	2003 - 2007	40.4 - 47.9	
Taos County ⁷	45.5 (32.3 - 62.7)	8	2003 - 2007	40.4 - 47.9	
Colfax County ⁷	44.0 (26.8 - 69.4)	4	2003 - 2007	40.4 - 47.9	
Santa Fe County ⁷	40.3 (33.4 - 48.1)	26	2003 - 2007	28.2 - 40.3	
Socorro County ⁷	40.2 (23.0 - 65.0)	3	2003 - 2007	28.2 - 40.3	
Rio Arriba County ⁷	28.6 (18.5 - 42.0)	5	2003 - 2007	28.2 - 40.3	
McKinley County ⁷	28.2 (18.9 - 40.3)	6	2003 - 2007	28.2 - 40.3	
Catron County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ¹
De Baca County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ¹
Guadalupe County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ¹
Harding County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ¹
Hidalgo County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ¹
Los Alamos County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ¹
Mora County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ¹
Union County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ¹

Notes:

Created by statecancerprofiles.cancer.gov on 02/04/2011 10:29 pm.

[State Cancer Registries](#) may provide more current or more local data.

Data presented on the State Cancer Profiles Web Site may differ from statistics reported by the State Cancer Registries ([for more information](#)).

¹ Incidence rates (cases per 100,000 population per year) are age-adjusted to the [2000 US standard population](#) (19 age groups: <1, 1-4, 5-9, ... , 80-84, 85+). Rates are for invasive cancer only (except for bladder which is invasive and in situ) or unless otherwise specified. Rates calculated using SEER*Stat. Population counts for denominators are based on Census populations as modified by NCI. The US populations included with the data release have been adjusted for the population shifts due to [hurricanes Katrina and Rita](#) for 62 counties and parishes in Alabama, Mississippi, Louisiana, and Texas The 1969-2007 US Population Data File is used with SEER November 2009 data. The 1969-2006 US Population Data File is used with NPCR data November 2008/January 2009 data.

¹ Source: CDC's National Program of Cancer Registries Cancer Surveillance System (NPCR-CSS) November 2009/January 2010 data submission and [SEER November 2009 submission](#).

³ Source: [SEER November 2009 submission](#). State Cancer Registry also receives funding from CDC's National Program of Cancer Registries.

¹ Source: SEER November 2009 submission.

² Data have been [suppressed](#) to ensure confidentiality and stability of rate estimates. Counts are suppressed if fewer than 16 cases were reported in a specific area-sex-race category.

³ Data not provided because it did not meet USCS data quality standards for one or more years during the rate period of data collection. While 96% of the US population resided in geographic areas with population-based cancer registries meeting the registry eligibility criteria for 2004, 4% of the US population was not yet represented in the [United States Cancer Statistics](#). American Cancer Society's [Facts & Figures](#) provides estimates of numbers of new cancer cases and deaths.

Incidence Rates[†] for New Mexico, 2003 - 2007

Non-Hodgkin Lymphoma

All Races (includes Hispanic), Male, All Ages

County ▲	Annual Incidence Rate over rate period cases per 100,000 (95% Confidence Interval) ▼	Average Cases per Year over rate period ▽	Rate Period	Interval Range	Interval Color
US (SEER + NPCR) ¹	23.1 (23.0 - 23.3)	§	2003 - 2007	N/A	N/A
New Mexico ³	18.1 (16.9 - 19.4)	160	2003 - 2007	N/A	N/A
Lincoln County [†]	26.8 (15.3 - 44.0)	4	2003 - 2007	24.1 - 26.8	
Santa Fe County [†]	24.0 (19.0 - 30.0)	16	2003 - 2007	20.9 - 24.0	
Dona Ana County [†]	20.9 (16.6 - 25.9)	17	2003 - 2007	20.9 - 24.0	
San Juan County [†]	20.8 (15.3 - 27.6)	10	2003 - 2007	19.9 - 20.8	
Chaves County [†]	20.5 (13.9 - 29.2)	6	2003 - 2007	19.9 - 20.8	
Bernalillo County [†]	19.8 (17.4 - 22.3)	53	2003 - 2007	14.9 - 19.8	
Eddy County [†]	16.6 (10.4 - 25.3)	4	2003 - 2007	14.9 - 19.8	
Sandoval County [†]	14.8 (10.1 - 20.8)	7	2003 - 2007	14.5 - 14.8	
Lea County [†]	14.5 (8.5 - 23.0)	4	2003 - 2007	14.5 - 14.8	
Otero County [†]	14.4 (9.1 - 21.8)	5	2003 - 2007	13.1 - 14.4	
Valencia County [†]	13.1 (7.9 - 20.4)	4	2003 - 2007	13.1 - 14.4	
Catron County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Cibola County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Colfax County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Curry County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
De Baca County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Grant County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Guadalupe County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Harding County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Hidalgo County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Los Alamos County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Luna County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
McKinley County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Mora County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Quay County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Rio Arriba County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Roosevelt County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
San Miguel County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Sierra County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Socorro County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Taos County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]

Torrance County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Union County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]

Notes:

Created by statecancerprofiles.cancer.gov on 02/04/2011 10:31 pm.

[State Cancer Registries](#) may provide more current or more local data.

Data presented on the State Cancer Profiles Web Site may differ from statistics reported by the State Cancer Registries ([for more information](#)).

[†] Incidence rates (cases per 100,000 population per year) are age-adjusted to the [2000 US standard population](#) (19 age groups: <1, 1-4, 5-9, ... , 80-84, 85+). Rates are for invasive cancer only (except for bladder which is invasive and in situ) or unless otherwise specified. Rates calculated using SEER*Stat. Population counts for denominators are based on Census populations as modified by NCI. The US populations included with the data release have been adjusted for the population shifts due to [hurricanes Katrina and Rita](#) for 62 counties and parishes in Alabama, Mississippi, Louisiana, and Texas The 1969-2007 US Population Data File is used with SEER November 2009 data. The 1969-2006 US Population Data File is used with NPCR data November 2008/January 2009 data.

¹ Source: CDC's National Program of Cancer Registries Cancer Surveillance System (NPCR-CSS) November 2009/January 2010 data submission and [SEER November 2009 submission](#).

³ Source: [SEER November 2009 submission](#). State Cancer Registry also receives funding from CDC's National Program of Cancer Registries.

⁷ Source: SEER November 2009 submission.

[†] Data have been [suppressed](#) to ensure confidentiality and stability of rate estimates. Counts are suppressed if fewer than 16 cases were reported in a specific area-sex-race category.

[§] Data not provided because it did not meet USCS data quality standards for one or more years during the rate period of data collection. While 96% of the US population resided in geographic areas with population-based cancer registries meeting the registry eligibility criteria for 2004, 4% of the US population was not yet represented in the [United States Cancer Statistics](#). American Cancer Society's [Facts & Figures](#) provides estimates of numbers of new cancer cases and deaths.

Incidence Rates[†] for New Mexico, 2003 - 2007

Ovary

All Races (includes Hispanic), Female, All Ages

County 	Annual Incidence Rate over rate period cases per 100,000 (95% Confidence Interval) 	Average Cases per Year over rate period 	Rate Period	Interval Range	Interval Color
US (SEER + NPCR) ¹	12.8 (12.8 - 12.9)	§	2003 - 2007	N/A	N/A
New Mexico ³	12.2 (11.3 - 13.2)	127	2003 - 2007	N/A	N/A
Sandoval County [†]	17.1 (12.5 - 22.8)	9	2003 - 2007	15.6 - 17.1	
Otero County [†]	15.5 (10.2 - 22.7)	5	2003 - 2007	14.9 - 15.5	
Santa Fe County [†]	15.5 (11.9 - 19.8)	13	2003 - 2007	14.9 - 15.5	
Valencia County [†]	14.8 (9.7 - 21.8)	5	2003 - 2007	13.2 - 14.8	
Eddy County [†]	14.5 (8.9 - 22.4)	4	2003 - 2007	13.2 - 14.8	
Dona Ana County [†]	13.1 (10.0 - 16.9)	12	2003 - 2007	11.5 - 13.1	
Chaves County [†]	12.4 (7.4 - 19.3)	4	2003 - 2007	11.5 - 13.1	
McKinley County [†]	11.4 (6.8 - 17.8)	4	2003 - 2007	10.1 - 11.4	
Bernalillo County [†]	10.8 (9.3 - 12.6)	36	2003 - 2007	10.1 - 11.4	
San Juan County [†]	10.0 (6.7 - 14.5)	6	2003 - 2007	10.0 - 10.0	
Catron County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Cibola County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Colfax County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Curry County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
De Baca County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Grant County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Guadalupe County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Harding County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Hidalgo County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Lea County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Lincoln County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]
Los Alamos County [†]	*	3 or fewer	2003 - 2007	*	Suppressed [†]

Luna County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Mora County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Quay County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Rio Arriba County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Roosevelt County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
San Miguel County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Sierra County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Socorro County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Taos County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Torrance County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Union County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷

Notes:

Created by statecancerprofiles.cancer.gov on 02/04/2011 10:33 pm.

[State Cancer Registries](#) may provide more current or more local data.

Data presented on the State Cancer Profiles Web Site may differ from statistics reported by the State Cancer Registries ([for more information](#)).

¹ Incidence rates (cases per 100,000 population per year) are age-adjusted to the [2000 US standard population](#) (19 age groups: <1, 1-4, 5-9, ... , 80-84, 85+). Rates are for invasive cancer only (except for bladder which is invasive and in situ) or unless otherwise specified. Rates calculated using SEER*Stat. Population counts for denominators are based on Census populations as modified by NCI. The US populations included with the data release have been adjusted for the population shifts due to [hurricanes Katrina and Rita](#) for 62 counties and parishes in Alabama, Mississippi, Louisiana, and Texas. The 1969-2007 US Population Data File is used with SEER November 2009 data. The 1969-2006 US Population Data File is used with NPCR data November 2008/January 2009 data.

² Source: CDC's National Program of Cancer Registries Cancer Surveillance System (NPCR-CSS) November 2009/January 2010 data submission and [SEER November 2009 submission](#).

³ Source: [SEER November 2009 submission](#). State Cancer Registry also receives funding from CDC's National Program of Cancer Registries.

⁷ Source: SEER November 2009 submission.

* Data have been [suppressed](#) to ensure confidentiality and stability of rate estimates. Counts are suppressed if fewer than 16 cases were reported in a specific area-sex-race category.

⁸ Data not provided because it did not meet USCS data quality standards for one or more years during the rate period of data collection. While 96% of the US population resided in geographic areas with population-based cancer registries meeting the registry eligibility criteria for 2004, 4% of the US population was not yet represented in the [United States Cancer Statistics](#). American Cancer Society's [Facts & Figures](#) provides estimates of numbers of new cancer cases and deaths.

Incidence Rates† for New Mexico, 2003 - 2007

Brain & ONS

All Races (includes Hispanic), Female, All Ages

County △	Annual Incidence Rate over rate period cases per 100,000 (95% Confidence Interval) ▼	Average Cases per Year over rate period ▽	Rate Period	Interval Range	Interval Color
US (SEER + NPCR) ¹	5.7 (5.7 - 5.8)	§	2003 - 2007	N/A	N/A
New Mexico ³	4.3 (3.8 - 5.0)	44	2003 - 2007	N/A	N/A
Santa Fe County ⁷	7.2 (4.8 - 10.5)	6	2003 - 2007	6.6 - 7.2	
Dona Ana County ⁷	6.5 (4.4 - 9.3)	6	2003 - 2007	3.9 - 6.5	
Bernalillo County ⁷	3.8 (2.9 - 4.9)	12	2003 - 2007	3.8 - 3.8	
Catron County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Chaves County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Cibola County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Colfax County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Curry County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
De Baca County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Eddy County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Grant County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Guadalupe County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷
Harding County ⁷	*	3 or fewer	2003 - 2007	*	Suppressed ⁷

Hidalgo County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ¹
Lea County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ¹
Lincoln County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ¹
Los Alamos County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ¹
Luna County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ¹
McKinley County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ¹
Mora County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ¹
Otero County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ¹
Quay County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ¹
Rio Arriba County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ¹
Roosevelt County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ¹
San Juan County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ¹
San Miguel County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ¹
Sandoval County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ¹
Sierra County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ¹
Socorro County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ¹
Taos County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ¹
Torrance County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ¹
Union County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ¹
Valencia County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ¹

Notes:

Created by statecancerprofiles.cancer.gov on 02/04/2011 10:34 pm.

[State Cancer Registries](#) may provide more current or more local data.

Data presented on the State Cancer Profiles Web Site may differ from statistics reported by the State Cancer Registries ([for more information](#)).

¹ Incidence rates (cases per 100,000 population per year) are age-adjusted to the [2000 US standard population](#) (19 age groups: <1, 1-4, 5-9, ... , 80-84, 85+). Rates are for invasive cancer only (except for bladder which is invasive and in situ) or unless otherwise specified. Rates calculated using SEER*Stat. Population counts for denominators are based on Census populations as modified by NCI. The US populations included with the data release have been adjusted for the population shifts due to [hurricanes Katrina and Rita](#) for 62 counties and parishes in Alabama, Mississippi, Louisiana, and Texas. The 1969-2007 US Population Data File is used with SEER November 2009 data. The 1969-2006 US Population Data File is used with NPCR data November 2008/January 2009 data.

¹ Source: CDC's National Program of Cancer Registries Cancer Surveillance System (NPCR-CSS) November 2009/January 2010 data submission and [SEER November 2009 submission](#).

³ Source: [SEER November 2009 submission](#). State Cancer Registry also receives funding from CDC's National Program of Cancer Registries.

⁷ Source: SEER November 2009 submission.

^{*} Data have been [suppressed](#) to ensure confidentiality and stability of rate estimates. Counts are suppressed if fewer than 16 cases were reported in a specific area-sex-race category.

[§] Data not provided because it did not meet USCS data quality standards for one or more years during the rate period of data collection. While 96% of the US population resided in geographic areas with population-based cancer registries meeting the registry eligibility criteria for 2004, 4% of the US population was not yet represented in the [United States Cancer Statistics](#). American Cancer Society's [Facts & Figures](#) provides estimates of numbers of new cancer cases and deaths.

Incidence Rates[†] for New Mexico, 2003 - 2007

Leukemia

All Races (includes Hispanic), Female, All Ages

County 	Annual Incidence Rate over rate period cases per 100,000 (95% Confidence Interval) 	Average Cases per Year over rate period 	Rate Period	Interval Range	Interval Color
New Mexico ³	10.1 (9.3 - 11.0)	104	2003 - 2007	N/A	N/A
US (SEER + NPCR) ¹	9.6 (9.6 - 9.7)	§	2003 - 2007	N/A	N/A
Chaves County ¹	12.7 (8.1 - 19.0)	5	2003 - 2007	12.2 - 12.7	
Santa Fe County ¹	12.1 (8.8 - 16.2)	9	2003 - 2007	11.2 - 12.1	
Bernalillo County ¹	11.1 (9.5 - 12.9)	37	2003 - 2007	11.0 - 11.1	
Dona Ana County ¹	10.9 (8.1 - 14.3)	10	2003 - 2007	10.5 - 10.9	

Lea County ¹	10.9 (6.2 - 17.7)	3	2003 - 2007	10.5 - 10.9	
Sandoval County ¹	10.4 (7.0 - 15.0)	6	2003 - 2007	9.3 - 10.4	
San Juan County ¹	9.2 (6.1 - 13.4)	5	2003 - 2007	9.2 - 9.2	
Catron County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
Cibola County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
Colfax County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
Curry County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
De Baca County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
Eddy County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
Grant County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
Guadalupe County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
Harding County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
Hidalgo County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
Lincoln County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
Los Alamos County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
Luna County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
McKinley County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
Mora County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
Otero County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
Quay County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
Rio Arriba County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
Roosevelt County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
San Miguel County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
Sierra County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
Socorro County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
Taos County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
Torrance County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
Union County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²
Valencia County ¹	*	3 or fewer	2003 - 2007	*	Suppressed ²

Notes:

Created by statecancerprofiles.cancer.gov on 02/04/2011 10:35 pm.

[State Cancer Registries](#) may provide more current or more local data.

Data presented on the State Cancer Profiles Web Site may differ from statistics reported by the State Cancer Registries ([for more information](#)).

[†] Incidence rates (cases per 100,000 population per year) are age-adjusted to the [2000 US standard population](#) (19 age groups: <1, 1-4, 5-9, ... , 80-84, 85+). Rates are for invasive cancer only (except for bladder which is invasive and in situ) or unless otherwise specified. Rates calculated using SEER*Stat. Population counts for denominators are based on Census populations as modified by NCI. The US populations included with the data release have been adjusted for the population shifts due to [hurricanes Katrina and Rita](#) for 62 counties and parishes in Alabama, Mississippi, Louisiana, and Texas The 1969-2007 US Population Data File is used with SEER November 2009 data. The 1969-2006 US Population Data File is used with NPCR data November 2008/January 2009 data.

¹ Source: CDC's National Program of Cancer Registries Cancer Surveillance System (NPCR-CSS) November 2009/January 2010 data submission and [SEER November 2009 submission](#).

³ Source: [SEER November 2009 submission](#). State Cancer Registry also receives funding from CDC's National Program of Cancer Registries.

⁷ Source: SEER November 2009 submission.

^{*} Data have been [suppressed](#) to ensure confidentiality and stability of rate estimates. Counts are suppressed if fewer than 16 cases were reported in a specific area-sex-race category.

[§] Data not provided because it did not meet USCS data quality standards for one or more years during the rate period of data collection. While 96% of the US population resided in geographic areas with population-based cancer registries meeting the registry eligibility criteria for 2004, 4% of the US population was not yet represented in the [United States Cancer Statistics](#). American Cancer Society's [Facts & Figures](#) provides estimates of numbers of new cancer cases and deaths.

Incidence Rates[†] for New Mexico, 2003 - 2007

Stomach

All Races (includes Hispanic), Female, All Ages

County 	Annual Incidence Rate over rate period	Average Cases per Year over rate period	Rate Period	Interval Range	Interval Color
---	---	--	-------------	----------------	----------------

	cases per 100,000 (95% Confidence Interval)				
New Mexico³	5.2 (4.6 - 5.9)		55	2003 - 2007	N/A
US (SEER + NPCR)¹	4.8 (4.7 - 4.8)	§		2003 - 2007	N/A
Dona Ana County⁷	6.9 (4.7 - 9.8)		6	2003 - 2007	4.9 - 6.9
Santa Fe County⁷	4.8 (2.9 - 7.6)		4	2003 - 2007	4.6 - 4.8
Bernalillo County⁷	4.5 (3.6 - 5.7)		15	2003 - 2007	4.5 - 4.5
Catron County⁷	*	3 or fewer		2003 - 2007	*
Chaves County⁷	*	3 or fewer		2003 - 2007	*
Cibola County⁷	*	3 or fewer		2003 - 2007	*
Colfax County⁷	*	3 or fewer		2003 - 2007	*
Curry County⁷	*	3 or fewer		2003 - 2007	*
De Baca County⁷	*	3 or fewer		2003 - 2007	*
Eddy County⁷	*	3 or fewer		2003 - 2007	*
Grant County⁷	*	3 or fewer		2003 - 2007	*
Guadalupe County⁷	*	3 or fewer		2003 - 2007	*
Harding County⁷	*	3 or fewer		2003 - 2007	*
Hidalgo County⁷	*	3 or fewer		2003 - 2007	*
Lea County⁷	*	3 or fewer		2003 - 2007	*
Lincoln County⁷	*	3 or fewer		2003 - 2007	*
Los Alamos County⁷	*	3 or fewer		2003 - 2007	*
Luna County⁷	*	3 or fewer		2003 - 2007	*
McKinley County⁷	*	3 or fewer		2003 - 2007	*
Mora County⁷	*	3 or fewer		2003 - 2007	*
Otero County⁷	*	3 or fewer		2003 - 2007	*
Quay County⁷	*	3 or fewer		2003 - 2007	*
Rio Arriba County⁷	*	3 or fewer		2003 - 2007	*
Roosevelt County⁷	*	3 or fewer		2003 - 2007	*
San Juan County⁷	*	3 or fewer		2003 - 2007	*
San Miguel County⁷	*	3 or fewer		2003 - 2007	*
Sandoval County⁷	*	3 or fewer		2003 - 2007	*
Sierra County⁷	*	3 or fewer		2003 - 2007	*
Socorro County⁷	*	3 or fewer		2003 - 2007	*
Taos County⁷	*	3 or fewer		2003 - 2007	*
Torrance County⁷	*	3 or fewer		2003 - 2007	*
Union County⁷	*	3 or fewer		2003 - 2007	*
Valencia County⁷	*	3 or fewer		2003 - 2007	*

Notes:

Created by statecancerprofiles.cancer.gov on 02/04/2011 10:36 pm.

[State Cancer Registries](#) may provide more current or more local data.

Data presented on the State Cancer Profiles Web Site may differ from statistics reported by the State Cancer Registries ([for more information](#)).

[†] Incidence rates (cases per 100,000 population per year) are age-adjusted to the [2000 US standard population](#) (19 age groups: <1, 1-4, 5-9, ... , 80-84, 85+). Rates are for invasive cancer only (except for bladder which is invasive and in situ) or unless otherwise specified. Rates calculated using SEER*Stat. Population counts for denominators are based on Census populations as modified by NCI. The US populations included with the data release have been adjusted for the population shifts due to [hurricanes Katrina and Rita](#) for 62 counties and parishes in Alabama, Mississippi, Louisiana, and Texas. The 1969-2007 US Population Data File is used with SEER November 2009 data. The 1969-2006 US Population Data File is used with NPCR data November 2008/January 2009 data.

¹ Source: CDC's National Program of Cancer Registries Cancer Surveillance System (NPCR-CSS) November 2009/January 2010 data submission and [SEER November 2009 submission](#).

³ Source: [SEER November 2009 submission](#). State Cancer Registry also receives funding from CDC's National Program of Cancer Registries.

⁷ Source: SEER November 2009 submission.

¹ Data have been [suppressed](#) to ensure confidentiality and stability of rate estimates. Counts are suppressed if fewer than 16 cases were reported in a specific area-sex-race category.

⁸ Data not provided because it did not meet USCS data quality standards for one or more years during the rate period of data collection. While 96% of the US population resided in geographic areas with population-based cancer registries meeting the registry eligibility criteria for 2004, 4% of the US population was not yet represented in the [United States Cancer Statistics](#). American Cancer Society's [Facts & Figures](#) provides estimates of numbers of new cancer cases and deaths.